


3.

6.

L. R. 1.









THE  
LONDON, EDINBURGH, AND DUBLIN  
PHILOSOPHICAL MAGAZINE  
AND  
JOURNAL OF SCIENCE.

CONDUCTED BY

SIR DAVID BREWSTER, K.H. LL.D. F.R.S.L.&E. &c.  
RICHARD TAYLOR, F.L.S. G.S. Astr.S. Nat.H.Mosc. &c.  
RICHARD PHILLIPS, F.R.S.L.&E. F.G.S. &c.  
SIR ROBERT KANE, M.D. M.R.I.A.

---

“Nec araneorum sane textus ideo melior quia ex se fila gignunt, nec noster vilior quia ex alienis libamus ut apes.” JUST. LIPS. *Polit. lib. i. cap. 1. Not.*

---

VOL. XXVIII.

NEW AND UNITED SERIES OF THE PHILOSOPHICAL MAGAZINE,  
ANNALS OF PHILOSOPHY, AND JOURNAL OF SCIENCE.

JANUARY—JUNE, 1846.

---

LONDON:

RICHARD AND JOHN E. TAYLOR, RED LION COURT, FLEET STREET,  
*Printers and Publishers to the University of London;*

SOLD BY LONGMAN, BROWN, GREEN, AND LONGMANS; CADELL; SIMPKIN,  
MARSHALL AND CO.; S. HIGHLEY; WHITTAKER AND CO.; AND  
SHERWOOD, GILBERT, AND PIPER, LONDON: — BY ADAM AND  
CHARLES BLACK, AND THOMAS CLARK, EDINBURGH; SMITH  
AND SON, GLASGOW; HODGES AND SMITH, DUBLIN:  
AND G. W. M. REYNOLDS, PARIS.

"Meditationis est perscrutari occulta; contemplationis est admirari  
perspicua . . . . . Admiratio generat quæstionem, quæstio investigationem,  
investigatio inventionem."—*Hugo de S. Victore.*





# CONTENTS OF VOL. XXVIII.

(THIRD SERIES.)

## NUMBER CLXXXIV.—JANUARY, 1846.

	Page
Mr. R. Hunt on the Influence of Magnetism on Molecular Arrangement (with a Plate) .....	1
Mr. R. W. Fox on certain Pseudomorphous Crystals of Quartz .....	5
Prof. J. R. Young on the General Expression for the Sum of an Infinite Geometrical Series .....	10
Drs. T. Tilley and D. Maclagan on the Conversion of Cane-sugar into a substance isomeric with Cellulose and Inuline .	12
Mr. G. G. Stokes's Remarks on Professor Challis's Theoretical Explanation of the Aberration of Light .....	15
Lieut.-Col. P. Yorke on the Solubility of Oxide of Lead in Pure Water .....	17
The Rev. B. Bronwin's Equations for the Determination of the Motion of a Disturbed Planet by means of M. Hansen's Altered Time.....	20
Lieut.-Col. Sabine on some Points in the Meteorology of Bombay (with a Plate).....	24
Mr. T. Taylor on some New Species of Animal Concretions ..	36
Mr. C. B. Cayley's Inquiries in the Elements of Phonetics ..	47
Mr. A. Smith on Fresnel's Theory of Double Refraction ....	48
Mr. J. D. Dana on the Origin of the constituent and adventitious Minerals of Trap and the allied Rocks .....	49
Mr. G. B. Jerrard's Reflections on the Resolution of Algebraic Equations of the Fifth Degree .....	63
Proceedings of the Royal Society.....	64
Action of Nitric Acid on Wax .....	66
Dry Distillation of Wax .....	67
Analysis of Phosphate of Alumina, by M. A. Delesse.....	68
A New Planet .....	69
Notice of an Aurora Borealis seen at Manchester .....	70
Meteorological Observations for November 1845 .....	71
Meteorological Observations made by Mr. Thompson at the Garden of the Horticultural Society at Chiswick, near London; by Mr. Veall at Boston; by the Rev. W. Dunbar at Applegarth Manse, Dumfries-shire; and by the Rev. C. Clouston at Sandwick Manse, Orkney.....	72

## NUMBER CLXXXV.—FEBRUARY.

Mr. H. Collen on the Application of the Photographic Camera to Meteorological Registration (with a Plate).....	73
--	----

	Page
Mr. G. G. Stokes on Fresnel's Theory of the Aberration of Light .....	76
Mr. E. Wilson's Observations on the Development and Growth of the Epidermis .....	82
The Rev. J. Challis on the Aberration of Light, in Reply to Mr. Stokes. ....	90
Dr. A. Waller's Observations on certain Molecular Actions of Crystalline Particles, &c.; and on the Cause of the Fixation of Mercurial Vapours in the Daguerreotype Process (with a Plate) .....	94
Note to Mr. Hennessy's Paper on the Connexion between the Rotation of the Earth and the Geological Changes of its Surface .....	106
The Rev. W. V. Harcourt's Letter to Henry Lord Brougham, F.R.S. &c., containing Remarks on certain Statements in his Lives of Black, Watt, and Cavendish .....	106
Mr. J. Cockle on a Proposition relating to the Theory of Equations .....	132
Mr. R. Moon on Fresnel's Theory of Double Refraction ....	134
Mr. R. Moon's Reply to some Remarks contained in Prof. Young's recent paper "On the Evaluation of the Sums of Neutral Series" .....	136
Jesuiticus's Remarks on a Paper by Mr. Moon on Fresnel's Theory of Double Refraction .....	144
The Editor's Observations on the subject of the preceding Communications .....	146
Proceedings of the Royal Society. ....	147
Analysis of a Substance occurring with Disthene, by M. A. Delesse .....	150
Hydrated Silicate of Magnesia, by M. A. Delesse .....	152
Analysis of the Elie Pyrope or Garnet, by Prof. Connell. ....	152
Analysis of Meteoric Iron from Burlington, Ostego County, New York, by Mr. C. H. Rockwell .....	154
Preparation of Chloro-acetic Acid .....	154
Composition of Phosphate of Ammonia and Magnesia .....	155
Composition of Common Phosphate of Soda .....	155
On several New Series of Double Oxalates, by M. Rees Heece ..	156
Reaction for the Discovery of Sulphurous Acid, by M. Heintz. .	157
Analysis of the Molares of a Fossil Rhinoceros .....	158
Experiments on the Yolk of Eggs, by M. Gobley .....	158
Meteorological Observations for December 1845. ....	159
Table .....	160

---

NUMBER CLXXXVI.—MARCH.

Mr. C. Langberg on the Determination of the Temperature and Conducting Power of Solid Bodies .....	161
--	-----

	Page
Mr. T. Hopkins on the Causes of the Semi-diurnal Fluctuations of the Barometer . . . . .	166
The Rev. J. Challis on the Principles to be applied in explaining the Aberration of Light. . . . .	176
Prof. J. W. Draper on the Cause of the Circulation of the Blood . . . . .	178
Mr. J. Cockle on the Existence of Finite Algebraic Solutions of the general Equations of the Fifth, Sixth, and Higher Degrees . . . . .	190
Mr. T. Taylor on some New Species of Animal Concretions . . . . .	192
Lieut.-Col. Sabine on the Winter Storms of the United States . . . . .	200
Mr. R. C. Taylor on the Anthracite and Bituminous Coal-Fields in China. . . . .	204
Prof. C. F. Schönbein on the Conversion of the solid Ferrocyanide of Potassium into the Sesqui-ferrocyanide . . . . .	211
Prof. C. F. Schönbein on the Decomposition of the Yellow and Red Ferrocyanides of Potassium by Solar Light . . . . .	211
Prof. Potter's Reference to former Contributions to the Philosophical Magazine, on Physical Optics . . . . .	212
Prof. J. R. Young on Differentiation as applied to Periodic Series: with a few Remarks in Reply to Mr. Moon . . . . .	213
Mr. Moon in Reply to Jesuiticus. . . . .	215
Proceedings of the Royal Society. . . . .	219
————— Royal Astronomical Society. . . . .	223
Experiments on the Spots on the Sun, by Prof. Henry . . . . .	230
Method of Purifying Oxide of Uranium from Nickel, Cobalt and Zinc, by Prof. Wöhler . . . . .	232
On some New Double Haloid Salts, by M. Poggiale . . . . .	232
On the Volatile Acids of Cheese, by MM. Iljenko and Laskowski . . . . .	234
On the Double Salts of the Magnesian Group . . . . .	235
Preparation of Hypophosphites. . . . .	236
Biela's Comet. . . . .	238
Meteorological Observations for January 1846 . . . . .	239
————— Table. . . . .	240

---

NUMBER CLXXXVII.—APRIL.

Mr. W. Brown, Jun., on the Oscillations of the Barometer, with particular reference to the Meteorological Phænomena of November 1842 (with Six Plates) . . . . .	241
Prof. De Morgan on the Derivation of the Word Theodolite. . . . .	287
Mr. T. Graham's Reply to the Observations of M. Pierre, on the Proportion of Water in the Magnesian Sulphates and Double Sulphates . . . . .	289
M. F. Donny on the Cohesion of Liquids and their Adhesion to Solid Bodies . . . . .	291

	Page
Dr. Faraday's Experimental Researches in Electricity.—Nineteenth Series. On the Magnetization of Light and the Illumination of Magnetic Lines of Force . . . . .	294
Lieut.-Col. Sabine on the Cause of remarkably Mild Winters which occasionally occur in England . . . . .	317
M. Pouillet's Observations on the Recent Researches of Prof. Faraday . . . . .	324
Mr. G. G. Stokes on the Aberration of Light . . . . .	335
Analysis of Diaspore from Siberia, by M. A. Damour . . . . .	336
On Boracic Æther . . . . .	337
Action of Boracic Acid on Pyroxylic Spirit . . . . .	339
On a Simple Method of Protecting from Lightning, Buildings with Metallic Roofs, by Prof. Henry . . . . .	340
Observations on Capillarity, by Prof. Henry . . . . .	341
Obituary . . . . .	343
Meteorological Observations for February 1846 . . . . .	343
————— Table . . . . .	344

---

NUMBER CLXXXVIII.—MAY.

Dr. Faraday's Thoughts on Ray-vibrations . . . . .	345
Mr. J. E. Teschemacher on the Wax of the Chamærops . . . . .	350
Mr. J. Middleton's Analysis of a Cobalt Ore found in Western India . . . . .	352
Mr. H. E. Strickland on the Structural Relations of Organized Beings . . . . .	354
Mr. W. J. Henwood's Abstract of Meteorological Observations made during the year 1845 at Gongo Soco, in the interior of Brazil . . . . .	364
Dr. R. D. Thomson on Pegmine and Pyropine, animal substances allied to Albumen . . . . .	368
The Rev. B. Bronwin on certain Definite Multiple Integrals . . . . .	373
Mr. W. R. Birt on the Storm-Paths of the Eastern Portion of the North American Continent . . . . .	379
Prof. De Morgan on the first introduction of the words <i>Tangent</i> and <i>Secant</i> . . . . .	382
Dr. J. Lhotsky's Complete Collection of Kepler's Works . . . . .	387
The Rev. J. Challis on the Aberration of Light, in Reply to Mr. Stokes . . . . .	393
Mr. J. Cockle on the Finite Solution of Equations . . . . .	395
Dr. Faraday's Experimental Researches in Electricity.—Twentieth Series. On new Magnetic Actions, and on the Magnetic Condition of all Matter . . . . .	396
Prof. Louyet's Description of a new Mercurial Trough . . . . .	406
Proceedings of the Royal Society . . . . .	408

	Page
Note by Mr. T. Hopkins on his Paper on the Semi-diurnal Fluctuations of the Barometer . . . . .	416
On some new Compounds of Perchloride of Tin, by M. Lewy . .	416
Analysis of two species of Epiphytes, or Air Plants, by John Thomson, A.M. . . . .	420
Analysis of <i>Ceradia furcata</i> Resin, by Robert D. Thomson, M.D.	422
Meteorological Observations for March 1846 . . . . .	423
Table . . . . .	424

---

NUMBER CLXXXIX.—JUNE.

Dr. D. P. Gardner's Researches on the Functions of Plants, with a view of showing that they obey the Physical Laws of Diffusion in the Absorption and Evolution of Gases by their Leaves and Roots . . . . .	425
Dr. C. F. Schœnbein on the relation of Ozone to Hyponitric Acid . . . . .	432
Mr. T. Graham on the Composition of the Fire-Damp of the Newcastle Coal Mines . . . . .	437
Dr. J. Stenhouse's Observations on the Resin of the <i>Xanthorœa hastilis</i> , or Yellow Gum-resin of New Holland . . . . .	440
Mr. H. Sloggett on the Constitution of Matter . . . . .	443
Messrs. Scoresby and Joule's Experiments and Observations on the Mechanical Powers of Electro-Magnetism, Steam, and Horses. . . . .	448
Dr. Faraday's Experimental Researches in Electricity.—Twentieth Series.—Action of Magnets on Metals generally ( <i>concluded</i> ) . . . . .	455
The Astronomer Royal on the Equations applying to Light under the action of Magnetism. . . . .	469
The Rev. W. V. Harcourt's Letter to Henry Lord Brougham, F.R.S. &c., containing Remarks on certain Statements in his Lives of Black, Watt and Cavendish . . . . .	478

---

NUMBER CXC.—SUPPLEMENT TO VOL. XXVIII.

The Rev. W. V. Harcourt's Letter to Henry Lord Brougham, F.R.S. &c., containing Remarks on certain Statements in his Lives of Black, Watt, and Cavendish ( <i>concluded</i> ) . . . . .	505
Prof. Owen's Observations on Mr. Strickland's Article on the Structural Relations of Organized Beings . . . . .	525
Prof. Marignac's Observations on Messrs. Lyon Playfair and Joule's Memoir on Atomic Volume and Specific Gravity . .	527

	Page
The Astronomer Royal's Remarks on Dr. Faraday's Paper on Ray-vibrations .....	532
Mr. R. Mallet's Explanation of the Vorticose Movement, assumed to accompany Earthquakes .....	537
Prof. E. Wartmann on the Causes to which Musical Sounds produced in Metals by discontinuous Electric Currents are attributable .....	544
Mr. E. F. Teschemacher's Account of various Substances found in the Guano Deposits and in their Vicinity .....	546
Dr. Gregory's Notes on the Preparation of Alloxan .....	550
On Chloroazotic Acid .....	555
Notices of New Localities of Rare Minerals, and Reasons for uniting several supposed Distinct Species, by Francis Alger.	557
Notice on certain Impurities in Commercial Sulphate of Copper, by Mr. S. Piesse .....	565
On a New Eudiometric Process, by Prof. Graham .....	566
Equivalent of Chlorine .....	566
On Hippuric Acid, Benzoic Acid, and the Sugar of Gelatine..	567
Comparative Analyses of Oriental Jade and Tremolite, by M. Damour .....	568
Meteorological Observations for April 1846 .....	569
————— Table .....	570
Index .....	571

## PLATES.

- I. Illustrative of Lieut.-Col. Sabine's paper on the Meteorology of Bombay.
- II. Illustrative of Mr. Hunt's paper on the Influence of Magnetism on Molecular Arrangement.
- III. Illustrative of Dr. Waller's paper on the Molecular Actions of Crystalline Particles.—Mr. Collen's paper on the Application of Photography to Meteorological Registration.
- IV. } Illustrative of Mr. Brown's paper on the Meteorological Phænomena of November 1842.
- V. }
- VI. }
- VII. }
- VIII. }
- IX. }

## ERRATA AND ADDENDA.

- Page 190, Note \*\*, between "3." and "p." add vol. xxvii.  
 ... 391, for *Salis deliquio* read *Solis*.  
 ... .., for *Bontschii* read *Bartschii*.  
 ... 393, for *Saganensibus* read *Saganensibus*.

THE  
LONDON, EDINBURGH AND DUBLIN  
PHILOSOPHICAL MAGAZINE  
AND  
JOURNAL OF SCIENCE.

[THIRD SERIES.]

JANUARY 1846.

- I. *The Influence of Magnetism on Molecular Arrangement.*  
By ROBERT HUNT, *Keeper of Mining Records, Museum of Economic Geology.*

[With a Plate.] *l. 75*

To Richard Phillips, Esq., F.R.S.

DEAR SIR,

HAVING been engaged some time since in investigating the influences of bodies on each other in the dark, the results of which investigations were published under the title of "Thermography," I then observed many peculiar effects which led me to believe that magnetic electricity had some influence in determining the arrangements of molecules. From that time until a few days since, the subject has rested with me without any further research. Having however put the subject to the test of experimental examination, I am induced, the results being of great interest, to transmit to you an account of my experiments. In doing this, I shall, for the present, confine myself strictly to a description of the arrangements used and the results obtained, reserving any theoretical views for some future period, when by a greater number and variety of experiments it appears probable some general law of action may be satisfactorily deduced.

1. I placed a concentrated solution of nitrate of silver in a test-tube, against the poles of a permanent horse-shoe magnet, having another tube containing a similar solution not in contact with it. The crystallization commenced first in the tube connected with the magnet, immediately at the point opposite the upper surface of the metal (Plate II. fig. 1); a large tabular crystal shot off from this point towards the bottom of the glass, dividing the lower portion of the fluid in two parts. Other crystals sprung off from different points above and be-

*Phil. Mag. S. 3. Vol. 28. No. 184. Jan. 1846.* B

low this crystalline plate, but all of them arranged themselves at angles inclining towards the magnet; no crystallization taking place in the upper stratum of the fluid. In the other tube, crystals formed irregularly throughout the fluid, but in no part were the crystals so dense as in the tube which I suppose to be under the influence of magnetism.

2. With a view of determining if the cooling influence of the metal had anything to do with the crystalline arrangement, portions of the same solution of nitrate of silver were put into glass capsules. One of these was placed against the poles of the magnet, and the other in contact with a mass of brass of the same weight. In the first, crystallization commenced opposite the north pole of the magnet, and proceeded slowly in regular lines to crystallize over every part; all these lines have a tendency towards the poles of the magnet. In the capsule in contact with the brass, crystallization commenced at a point furthest from the metal, and even when the fluid had become quite cold, nearly one quarter of it, which was nearest the mass of metal, remained quite free from any crystalline formation.

3. To exhibit this in a more striking manner, a capsule was placed between the mass of brass and the magnet, in contact with each, as shown in fig. 2; the solution of nitrate of silver in this case, not being so concentrated as that previously used, the arrangement was allowed to remain at rest for some hours. It was then found that crystallization had taken place only over one portion of the fluid, and that immediately in connexion with the north pole of the magnet, except three long crystals which sprung from the fluid opposite the south pole, and were directed towards those springing from the north pole. This experiment was repeated four times, and, except when the solution was so concentrated as to crystallize almost immediately, the same result was obtained.

4. The phænomenon of molecular disposition under magnetic influence is pleasingly seen by a modification of the arrangement described. The two glass capsules with their solutions are placed on a plate of glass blackened on its under surface, one glass being put in contact with the brass and the other with the magnet. Their images are to be observed in the black mirror on which they rest, the light falling upon them at an angle of about  $25^{\circ}$ . As the fluids cool, the circulating currents coloured by their refracting powers are seen in the mirror. In the image of the capsule in contact with the brass, no regularity of circulatory movement is observable; but in that under magnetic influence, a series of perfectly regular curved lines proceed from the circumference to the



centre; and these are crossed by small streamers springing laterally from these primary curves, presenting an appearance similar to that shown in fig. 3. These curves are constantly varying in position, but they uniformly preserve the utmost regularity.

5. The magnet was suspended from a tripod, and two steel needles attached to its poles; these needles were made to dip into a solution of nitrate of silver in a watch-glass. As the pellicle formed over the surface, it arranged itself in a series of curved lines, as represented in fig. 4, which are strikingly similar to those produced by sprinkling iron-filing on stretched paper placed over a magnet. That these curves are due to magnetic influence there can be no doubt, as no such effect could be produced by any cooling influence, independent of magnetic excitation.

6. A similar arrangement was allowed to remain in action for twelve hours. At the end of this time crystallization had taken place in every part of the fluid, but there was an evident tendency to a curvilinear arrangement of the crystals. Around the wire depending from the north pole of the magnet, some revived silver had made its appearance: no such change was discovered at the south pole.

7. Wires similarly suspended were dipped into a solution of sulphate of iron. Crystallization commenced around the wire at the north pole, but after a few hours crystals had formed around both of the wires, but in the greatest quantity around the north pole wire. On removing them from the solution, the crystals were found to present an arrangement similar to that represented in fig. 5, showing obviously a tendency to arrange themselves along lines of magnetic direction.

8. A solution of protonitrate of mercury was placed under similar circumstances; crystallization commenced at the wire suspended from the north pole, and proceeded rapidly to a line midway between the two wires; one-half of the fluid being crystallized and the other remaining fluid. At length a few crystals formed around the wire hanging from the south pole, which all took a direction towards the opposite arrangement of crystals.

9. With a more dilute solution, the crystallization of the nitrate of mercury took place only around the wire at the north pole, and immediately at the central point between the two wires, from which small needle-shaped crystals radiated towards either pole.

10. A plate of glass, with an edge of clay, forming a shallow trough, was placed upon the poles of an electro-magnet,

#### 4 *On the Influence of Magnetism on Molecular Arrangement.*

capable of supporting fifty pounds when connected with a single galvanic pair excited by water acidulated with sulphuric acid. On pouring a warm and tolerably strong solution of nitrate of mercury into the trough, there was immediately formed over the surface a series of beautifully regular curves from pole to pole, as shown in fig. 6, which also represents the arrangement.

11. A similar glass trough, filled with a moderately strong solution of the nitrate of mercury, was supported on the poles of the same electro-magnet, connected with a small battery of a more permanent, but less powerful arrangement, and all was allowed to remain at rest until crystallization had taken place. The result was similar to that already described (9.), but much more strikingly shown. The order of arrangement taken by the crystals is shown in Plate II. fig. 7.

12. A plate of copper with an edging of wax was placed on the electro-magnet in the same manner as the glass plate; over it a very weak solution of nitrate of silver was quickly poured; the plate immediately blackened from the decomposition of the silver salt by the copper. In about a minute the finely divided silver arranged itself into curves, as represented in fig. 8, which were after a few minutes again destroyed. By using a sheet of chemically-pure copper, obtained by electrotype deposit, I found a permanent impression of these curves could be obtained, owing to the oxidation of the copper along the spaces, which the finely divided silver, when distributed in curve-lines, did not cover.

13. A plate of hard copper, such as is used by engravers, was placed in precisely the same circumstances, and covered with a tolerably strong solution of nitrate of silver. It was left in contact with the electro-magnet for a night. On washing off the deposit of silver which covered it, it was found that the acid of the silver salt had bitten deeply into the plate over an oval space around the poles, leaving a small space between them quite bright. The copper over this etched space was covered with an immense number of minute holes; and beyond this the oxidation of the surface had proceeded in curved lines, as represented in fig. 9. We thus have permanent evidence of the influence of magnetic force in determining chemical action.

14. Into one of the glass troughs before named, placed on the electro-magnet, a weak solution of nitrate of silver was poured, and into this an equally weak solution of sulphate of iron. In about five minutes precipitation of silver commenced; this precipitate arranged itself over the glass in curves proceeding from and around the poles in the same manner as it

distributed itself over the copper plate. In a short time, precipitation increasing, two curious curved spaces were formed by the fine deposit, proceeding from one pole towards the other in opposite directions, increasing in width as they proceeded, until they were abruptly checked at a little distance from the poles towards which they were directed; these spaces being very distinct from the first formed curved lines. Fig. 10 represents this very interesting arrangement.

These experiments are sufficient to show that magnetism exerts a powerful influence on molecular arrangements, and that it regulates the direction of crystalline formations. I hope to be enabled to pursue this interesting inquiry still further; it has most important bearings on many of the great phenomena of nature, and I am therefore anxious thus early in my inquiry to call attention to the singular and conclusive results which I have obtained.

I have the pleasure of remaining,

Dear Sir, yours truly,

6 Craig's Court, Dec. 10, 1845.

ROBERT HUNT.

---

## II. *On certain Pseudomorphous Crystals of Quartz.*

By ROBERT WERE FOX, *Esq.*\*

I SUBMIT to the Society's notice some specimens of quartz, with pseudomorphous octahedral crystals of the same substance, which appear to me to possess a sort of historical interest, or at least to indicate that a succession of changes must have occurred in the condition of the mineral vein from which they were taken. They were found by S. Peters (dealer in minerals) in one of the heaps of vein stones, at the Consolidated Mines, and I understand were broken from a copper vein in "*killas*," at the depth of about 160 fathoms below the surface. He observed that many of the crystals contained water, and he secured some of it for me, by carefully breaking some of them. This he did mostly in my presence, and we had considerable difficulty in collecting even very small portions of the liquid in different phials. Two of these portions were nearly tasteless, or saline in a very slight degree, as far as I could judge from a single drop of each. In both common salt was detected, and nothing else in one of the portions; but the other, when evaporated, left minute needle-formed crystals, which I was prevented by an accident from examining. The third portion of water was much more in

\* Read at a meeting of the Cornwall Polytechnic Society, on the 8th of October, 1845, and communicated by the Author.

quantity than both the others—nearly a tea-spoonful, and obtained from only one crystal. It was very acrid to the taste, and gave very copious precipitates when tested by muriate of barytes and hydrocyanate of potash, showing the presence of much sulphuric acid and iron. Oxalate of ammonia and nitrate of silver, indicated, moreover, the presence of lime and muriatic acid. The saline matter in this water (mostly sulphate of iron) was equal to one-tenth of its weight; and if it contained any common salt, of which I am not positive, the proportion was very small indeed. Litmus paper showed an excess of acid, the nature of which was not ascertained.

Many of the pseudomorphous crystals are more than an inch in diameter, and are partly or entirely filled with crystalline quartz, whilst others are empty, or partly filled with more or less numerous fragments of disintegrated fluor. I counted nearly a hundred of such fragments taken from one of the crystals or cavities, exclusive of many other very small pieces. All the fragments are corroded, and indicate, by their rounded edges and indented surfaces, the action of a solvent which penetrated most readily between the *planes of cleavage*\*. Besides this disintegrated fluor, perfect octahedrons of fluor occur in the same specimens; but they were rather more imbedded in the quartz and more protected from injury than the others. Water was found alone in some of the pseudomorphous crystals or cavities, and in others it was found with fragments of fluor, or with crystalline quartz.

The most perfect pseudomorphous octahedrons occur within large cavities of quartz. Some of the latter are more than two inches in diameter, having the same form, and their sides generally parallel to those of the former.

The quartz specimens to which the crystals are attached, present, when broken, the appearance of fortification agate, having lines parallel to their structure of transparent and milk-white quartz, differing in thickness; these seem to indicate that the siliceous matter had been deposited at intervals of greater or less duration, or at least under different circumstances. After a time an entire change of conditions apparently occurred in the vein, and octahedral crystals of fluor were formed on the quartz; then silex was deposited either in a compact form, or in minute crystals, and coated the crystals of fluor; afterwards fluor again appeared, forming octahedrons over the others, and mostly with sides and angles parallel to them. These processes appear from some of the

\* When crystals of alum were kept for a time in water, the planes of cleavage were first acted on, and fragments were separated from the crystals, resembling those of the disintegrated fluor.

crystals to have been again repeated: then came a coat of silex over the fluor, or judging from the lines, many coats of it, forming a thick crust, having a surface of small quartz crystals. Some specimens were found at the same time with one or more layers of quartz between two or more portions of fluor, which tend to confirm these views.

I think it may be inferred, from the well-defined and smooth impressions which the octahedrons of fluor have left in the quartz, and the general parallelism of the sides and angles of the outer cavities to those of the smaller pseudomorphous crystals inclosed in them\*, that the inner and outer crystals of fluor were perfect and uninjured until after the whole series of them were coated with quartz. At some subsequent period then it would appear that other changes occurred in the vein, and that the solution or destruction of the fluor commenced. Some of the cavities which were found to contain water only, as well as those which contained water together with disintegrated fluor, have the appearance of having been so hermetically sealed, that it is difficult to understand how the liquid solvent could have obtained access to the fluor and abstracted it from its case. It cannot be supposed that the pressure of the column of water above it, although equal to more than half a ton on some of the larger crystals, could alone have produced the effects; for not only must the solvent have been continually admitted through the crusts of the quartz, but the salts resulting from the solution of the fluor must, at the same time, have passed through them in the opposite direction,—a sort of *endosmose* and *exosmose* must have existed, as I conceive, to produce the phænomena; whilst in other instances, the thick envelopes of quartz were impervious and protected the fluor from injury. The salts resulting from the solution of the fluor must have been soluble, although this condition seems to present some difficulties under the circumstances of the case; and doubtless the destruction of the fluor was very slowly effected in many instances, and in others it was begun, but never completed. The differences in the saline contents of the water obtained from some of the crystals is another circumstance of some interest, indicating the existence of different conditions in the vein when the water was last admitted into the respective crystals.

The phænomena exhibited by these minerals cannot, I con-

\* How are such coincidences to be accounted for? Are we to assume that *polarising forces* have determined the arrangement? In many instances the layers of quartz which were *interposed* between the crystals are very thin, imperfect, and pervious to water; but in others they are not so, and some of the inner crystals now contain water.

ceive, be accounted for but by supposing the water existing in the fissures of the earth to have been changed by circulation from time to time, and to have been charged with different ingredients at different periods.

I have on former occasions alluded to various causes which would produce circulation in the subterranean waters, such as the opening or closing of any portions of fissures; the ascent of warm and the descent of cooler currents of water, in consequence of the differences in their specific gravities; or in some instances by the pressure of the sea-water acting on the fresh\*. Nearly two years ago I stated in this room my views in reference to the operation of this latter cause on land springs, and at the same time I attempted to show the possibility, not to say probability, of steam existing in fissures *below* the water at a very great depth. I may perhaps be permitted to refer again to this subject, because it appears to me to be one of some interest. I then took it for granted that the temperature of the earth increases in some proportion to the increase of depth below its surface, and that if the ratio be taken at  $1^{\circ}$  Fahr. for every forty-eight feet, as found in our deep mines, and if Le Roche's data for calculating the elastic force and density of steam be adopted, the forces of steam and of water pressure would balance each other at rather more than nine miles deep, each being equal to the pressure of more than 1400 atmospheres. The density of the steam would there be about one-fourth that of water at  $60^{\circ}$  Fahr., and its temperature above  $1050^{\circ}$  Fahr. But the temperature may probably not increase so rapidly as this at great depths, and the equilibrium in the pressures of the column of water and of steam may occur much further below the surface, where the density of steam under an augmented pressure of water would, of course, be still greater. However this may be, it would seem that, under any probable circumstances in regard to the ratio of increase in the earth's temperature, the increase in the pressure of the lengthened column of water would not keep pace with the rapidly increasing tendency of the water in descending into more heated parts of the earth to expand into steam, the elasticity of which at very high temperatures, when confined and in contact with water, is greatly augmented by very small increments of sensible heat.

No water could long remain unchanged into steam below the line of division between them, and there the steam would

\* Columns of *sea and spring water*, about five feet high, balanced against each other in a U-shaped tube, more than a year ago, still remain unmixed, showing nearly the same difference of level as at first (exceeding an inch).

be denser than at any deeper station, for it would be continually diminishing in density in descending further, from the augmentation of the temperature of the earth, because the *expanding* influence of the increasing heat would much exceed the *condensing* influence of the extended column of steam, added to that of the nearly constant column of water.

The line of demarcation between the water and steam would, doubtless, conform in some degree to the inequalities of the surface. It may be difficult at first to conceive the steam capable of supporting the water, or rather of existing permanently under it; but this difficulty will, I think, be obviated by the consideration, that the points of contact may be, for the most part, in very narrow fissures, or mere cracks in the rocks; and that the water being greatly heated, may be much less than *four times* the density of the steam in immediate contact with it. A continual struggle would, no doubt, exist between the water and steam under such circumstances, so that in many places they would alternately encroach beyond the line of demarcation; but as the checks on both would increase in proportion to the extent of their encroachments from the diminution of the temperature above and its augmentation below, such encroachments would probably not be very extensive, or of long duration under ordinary circumstances. Suppose a temporary encroachment of the water on the limits of the steam to occur at one point, the steam would probably encroach on the water at another at the same time, and then, reactions taking place, the effects would be reversed. Thus, assuming what indeed would appear to follow from *admitted data as necessary consequences*, steam would not only exist below the water, but such oscillations would tend to give motion and activity to the water in the neighbouring fissures, causing it to circulate in the earth more or less freely and extensively according to circumstances. In volcanic districts, where the heat may be great at comparatively small depths, analogous phænomena sometimes occur at the surface, which are probably caused by the action and reaction of steam and water. Amongst these may be included the intermitting Geyser springs in Iceland, as well as some of the mud volcanoes found in Sicily, and in Asia, and America.

It seems probable that earthquakes may be produced by the action of highly elastic vapour rapidly generated at great depths, in consequence perhaps of copious and sudden influxes of water into intensely heated parts of the earth; and their lines of direction are doubtless influenced by those of the fissures or veins of the districts in which they occur. But these are phænomena of comparatively rare occurrence, and

it is no wonder that they should be so, when we consider how vastly greater must be the force required to uplift the rocky crust of the earth and wrench it asunder, than that which will support a column of water equal to the thickness of that crust.

Since the foregoing paper was read, I have rather hastily examined some other portions of water taken from different pseudomorphous crystals. One of those portions contained muriatic and sulphuric acids, iron, a trace of lime, and of common salt. Acid was a little in excess, and some peroxide of iron was left in the cavity from which the water was taken. In another the same acids were detected and some iron. In the third portion there seemed to be nothing besides a little common salt. In many of the octahedral cavities, oxide of iron was found, and sometimes iron pyrites or copper pyrites adhering to the sides; these were apparently deposited from some of the water which had entered the crystals in some instances, but in others they were evidently imbedded in the fluor, and, adhering to the deposit of quartz, were not dissolved with the former.

Earthy carbonate of iron occurs in some cavities mixed with very minute crystals of quartz; and I have one pseudomorphous quartz crystal which is filled with fragments of fluor, intermixed with translucent fragments of carbonate of iron and earthy carbonate of iron, all curiously cemented together into one mass; the iron ore being rather in excess.

I have also some hollow pseudomorphous crystals of quartz formed originally on carbonate of iron, which appear to be water-tight, and yet the latter substance has, like the fluor, been abstracted.

III. *On the General Expression for the Sum of an Infinite Geometrical Series.* By J. R. YOUNG, Professor of Mathematics in Belfast College\*.

THE general expression for the sum of the infinite series  
 $1 - x + x^2 - x^3 + x^4 - \&c.$   
 is

$$S = \frac{1}{1+x} - \frac{x^\infty}{1+x},$$

which reduces to  $\frac{1}{1+x}$  when  $x$  is a proper fraction, either po-

\* Communicated by the Author.



sitive or negative, on account of the evanescence of  $x^{\infty}$ . It is usual to consider the infinite exponent in this expression as invariable throughout all the changes of  $x$  within the limits 0 and 1; although it is known that for any fixed exponent short of infinite, however great it may be, the expression into which it enters becomes more and more considerable as  $x$  advances from 0 towards 1; and notwithstanding the additional fact, that when this exponent is actually infinite, the expression referred to becomes ultimately equal to  $\frac{1}{e}$ .

But it is evident—due weight being given to the circumstances here mentioned—that this assumption, as to the invariability of the infinite exponent, is unwarrantable and erroneous; and that the exponent must follow some law of variation exactly fitted to counteract and neutralize the tendency which, as  $x$  approaches to 1, the expression  $x^{\infty}$  would otherwise have to depart from zero, and ultimately to become  $\frac{1}{e}$ .

If  $x$ , at any stage of its approach to 1, be generally represented by  $1 - \frac{1}{k}$ , then the law of variation alluded to will be expressed by  $\infty'' = k \infty'$ : that is, the exponent must vary as  $k$ . For it is a remarkable fact that, commencing with the exponent 4 and proceeding onwards to infinity, we shall invariably have

$$\left(\frac{3}{4}\right)^4 = \cdot 3 \dots, \left(\frac{4}{5}\right)^5 = \cdot 3 \dots, \left(\frac{5}{6}\right)^6 = \cdot 3 \dots, \left(\frac{6}{7}\right)^7 = \cdot 3 \dots, \\ \left(\frac{16801}{16802}\right)^{16802} = \cdot 3 \dots, \dots \left(\frac{25684}{25685}\right)^{25685} = \cdot 3 \dots, \dots \left(1 - \frac{1}{\infty}\right)^{\infty} = \cdot 3 \dots$$

And since  $(\cdot 3 \dots)^{\infty}$  is necessarily zero, and no power short of infinite can give zero, it follows that in order that  $\left(1 - \frac{1}{k}\right)^{\infty''}$  may be uniformly zero, and that all tendency to depart from zero may be counteracted,  $\infty''$  must be  $k \infty'$ ; so that the strictly accurate form for S is

$$S = \frac{1}{1 + \left(1 - \frac{1}{k}\right)} - \frac{\left(1 - \frac{1}{k}\right)^{k \infty'}}{1 + \left(1 - \frac{1}{k}\right)},$$

which is equal to  $\frac{1}{2}$  when  $k$  is infinite. And in this manner is the formula, employed in my paper (p. 363, last vol.), established.

In the same way that it has now been proved that  $\left(1 - \frac{1}{k}\right)$  is always equal to  $\cdot 3 \dots$ , whatever be  $k$ , above 3, may it be further shown that  $\left(1 + \frac{1}{k}\right)^k$  is always equal to  $2 \dots$ ; and thence that  $\left(1 + \frac{1}{k}\right)^{k\infty}$  is necessarily infinite when  $k$  is: so that it is indisputably true that the extreme of the *convergent* cases of the above series S, usually written in the form

$$1 - 1 + 1 - 1 + 1 - 1 + \&c.$$

is  $\frac{1}{2}$ , and that the extreme of the *divergent* cases, usually written in the same form, is really infinite, as stated in my former paper; which last conclusion could never have been anticipated from the theory hitherto prevalent. The views now developed are only the continuation and completion of those exhibited in my paper on Series submitted to the British Association in June 1845. If I have been anticipated in any of these views, which are doubtless calculated to produce a reform in the existing theory, I hope to be informed of the circumstance through the medium of this Journal. I have only further to add, that when an expression for the convergent cases of a series is found—as it often may be by aid of the differential theorem—then the general equivalent of the series may afterwards be ascertained by developing this expression sufficiently far to unfold to us the general form of the remainder. The expression for the convergent cases of the general series, discussed at page 439 of the last volume, may in this manner be determined; and the development of this expression by common division, as there proposed, furnishes the formula by which that expression must be corrected, in order that the algebraical equivalent of the series may be exhibited in its utmost generality.

Belfast, November 21, 1845.

IV. *On the Conversion of Cane-sugar into a substance isomeric with Cellulose and Inuline.* By THOMAS TILLEY, Esq., Ph.D., and DOUGLAS MACLAGAN, M.D., F.R.S. Edin.\*

WHEN the juice of beetroot undergoes fermentation at temperatures varying from  $30^{\circ}$  to  $40^{\circ}$  C., the cane-sugar which it contains is at first converted into sugar of grapes, and after some time into mannite, lactic acid and a

\* Communicated by the Chemical Society; having been read April 21, 1845.

gummy substance, having a composition identical with that of gum-arabic. This is remarkable, inasmuch as it affords an instance of what may be called a retrograde chemical action, the sugar being converted into dextrine,—a change similar to that which occurs in fruits when they lose their sweetness, and assume that condition commonly called “sleepy.” The conversion of cellulose into dextrine and sugar seems to be a process of continual occurrence and great importance in the vegetable œconomy, but we are not aware of any example of the reverse of this action, except those instances mentioned above; in the former of which sugar is converted by fermentation into a body having all the properties and composition of gum; in the latter, the sugar being changed into cellulose\*. We therefore consider the observation we are about to describe to be possessed of some interest, as affording another case of a similar retrograde action. It has been observed that the effervescing drinks known as lemonade, gingerade, &c., made by forcing carbonic acid gas into solutions of sugar variously flavoured with tartaric acid and essential oils, in certain cases lose their fluidity, and assume a thick, slimy consistence. When the bottles containing these thickened liquids are opened, the expansion of the carbonic acid expels their contents with difficulty, owing to their extreme tenacity. Instances of this change are of continual occurrence, all the manufacturers of whom we have inquired having observed it when the bottles had been kept for some time. Various opinions have been expressed by them as to the cause of the conversion, but it seems to occur invariably when the liquor is kept long enough. We are indebted to Mr. Baildon of this city (Edinburgh) for an opportunity of examining a sample of gingerade, in which this thickening had occurred. This liquid is made by sweetening an infusion of ginger-roots with cane-sugar, and flavouring it with oil of lemons and tartaric acid; this is then placed in bottles, and carbonic acid forced by pressure into the fluid. Another manufacturer uses the following ingredients in the preparation of effervescing lemonade:—2 ounces of honey, 4 pounds of sugar, 2 ounces of citric acid, 2 drachms of oil of lemons and  $1\frac{1}{2}$  ounce of bicarbonate of soda. According to the opinion of this manufacturer, the change occurs chiefly in winter, when the liquid is exposed to cold, and he thinks that the addition of a double proportion of honey tends to prevent it. To separate the substance to which the viscosity was owing, the contents of a bottle were digested with six or seven parts of alcohol, under the action of which the gummy matter consolidated, and when dried became so hard

\* See Mulder's *All. Phys. Chem.*, p. 243 *et seq.*

as to be pulverizable. After being powdered, it was again digested and washed with alcohol until nothing more was dissolved. When dried at  $100^{\circ}$  C. it had the appearance of a semi-transparent horny substance, and was sufficiently elastic to render pulverization difficult. The alcohol contained in solution a quantity of sugar of a brownish colour, quite uncrystallizable, and rendered sour by the presence of the acid used in the manufacture.

The gummy substance treated with cold water slowly re-assumes its original appearance. When treated with a large quantity of boiling water it forms a mucilage, which filters with difficulty. Iodine produces no effect on the solution. Subjected to Trommer's test for dextrine, sugar and gum, this did not indicate the presence of any of these substances. With nitric acid it produces oxalic acid. It gives a precipitate with diacetate of lead. It contains, after having been washed with alcohol, a small quantity of ashes, amounting to 1.37 per cent. It was analysed in the usual manner.

I. 0.746 of substance gave, with oxide of copper and chlorate of potash, 4.070 HO and  $1.1735 \text{ CO}_2 = 0.04727 \text{ H}$  and 0.32448 C.

II. 0.1525 of substance gave 0.092 HO and  $0.232 \text{ CO}_2 = 0.010222 \text{ H}$  and 0.06474 C.

These numbers, allowance being made of the ashes, give the following proportions:—

	I.	II.	Atoms.	Calculated.
Carbon . .	43.80	43.31	24	43.71
Hydrogen . .	6.14	6.80	21	6.25
Oxygen . .	50.06	49.89	21	50.04

From this it would appear that this gummy substance is isomeric with cellulose and inuline\*.

This substance, which has a composition similar to cellulose and inuline, is evidently formed from the cane-sugar in the lemonade, as all its other constituents exist in too small quantity to admit the idea of their having been its origin.

* Cellulose, Payen. Endine.	From turnip. Fromberg.
Carbon . 43.40	Carbon . 43.95
Hydrogen . 6.12	Hydrogen . 6.13
Oxygen . 50.38 <sup>a</sup>	Oxygen . 49.66 <sup>b</sup>
Inuline. Parnell.	From Dahlia root. Payen.
Carbon . 43.95	Carbon . 44.19
Hydrogen . 6.30	Hydrogen . 6.17
Oxygen . 49.75 <sup>c</sup>	Oxygen . 49.64 <sup>d</sup>

<sup>a</sup> *Ann. des Sc. Nat.*, 1840, p. 73. Bot.

<sup>b</sup> Mulder, *Op. cit.*, p. 203.

<sup>c</sup> *Phil. Mag.*, vol. xvii. p. 126.

<sup>d</sup> *Op. cit.*, p. 91.

2 atoms sugar . . . .	C 24	H 22	O 22
1 ... water . . . .		H	O
1 ... gummy substance	C 24	H 21	O 21

This substance is formed then from 2 atoms of sugar by the abstraction of 1 atom of water.

As a solution of the gummy substance gave a compound with lead, we endeavoured to obtain by its aid its atomic weight. 0.260 of the precipitate gave of lead and oxide of lead quantities equal to 0.316 oxide of lead, which, when allowance is made for ashes, is equal to 55.8 per cent. of oxide of lead. We had not enough of the salt to enable us to make the combustion, but have calculated the formula from the quantity of lead.

		Atoms.	Calculated.
Carbon . . .	19.31	24	1834.4 = 18.7
Hydrogen . . .	2.76	21	260.0 = 2.7
Oxygen . . .	22.11	21	2100.0 = 21.4
PbO . . .	55.80 found	4	5578.0 = 57.1

From 55.8 per cent. oxide of lead the atomic weight found is 4400.0. The calculated one is 4198.4.

We had imagined that this curious change in sugar might have been the effect of organization, but our friend Mr. John Goodsir was kind enough to examine the substance, and informed us that he could discover no trace of organization.

V. *Remarks on Professor Challis's Theoretical Explanation of the Aberration of Light.* By G. G. STOKES, M.A., Fellow of Pembroke College, Cambridge\*.

**T**HERE are a few points connected with Prof. Challis's paper on the Aberration of Light, published in the number of this Magazine for November 1845, respecting which I wish to offer a few remarks.

In the first place I perfectly agree with Prof. Challis, that the explanation of aberration is really independent of the manner in which light may pass through the eye; but I cannot agree with him that it is necessary to suppose that we see a star in its true place, and that it is the wire of the telescope with which it is observed that is affected by aberration. The following mode of viewing the subject, due to Boscovich, will perhaps put the matter in a clearer point of view.


If we wish to determine the real or apparent direction of an object, we may, theoretically speaking, adopt the following plan:—Let two small circular holes be so adjusted that the

\* Communicated by the Author.

light from the object which passes through the centre of the one shall also pass through the centre of the other. The line joining the centres of the holes will then determine the direction of the object. Now this is, in principle, just what is done in the case of an astronomical instrument, only, the fixed points are replaced by the optical centre of the object-glass of the telescope with which the object is viewed, and by the wire to which it is referred. When the image of a star is bisected by the wire, we define the apparent direction of the star to be that of the line joining the optical centre of the object-glass with the bisecting wire. Whether it is the wire or the star which is seen out of its true place, is a question with which we have no concern. The answer which we shall be disposed to give to it depends on the theory of aberration which we adopt. According to the theory of aberration which I explained in the July number of this Magazine, the answer would of course be, that it is the wire which is seen in its true place.

The principal thing, however, to which I object in Prof. Challis's paper, is the reasoning by which he establishes his equation (5.). In the figure,  $ab$  is a very small portion of a wave of light, which in the small time  $t$  would be propagated to  $cd$  if the æther from  $a$  to  $b$  were moving with the velocity of the æther at  $b$ , while, in consequence of the difference in the velocity of the æther at  $a$  and  $b$ , the disturbance at  $a$  is propagated to  $e$ . Now Prof. Challis takes  $cae$  for the angle through which the normal to the wave's front is displaced as the wave passes from  $ab$  to  $ed$ . But  $ae$  is only the direction in space along which the disturbance at  $a$  is propagated, a direction which has no immediate relation to the normal to the wave, inasmuch as it differs from it by an angle which is of the order of the aberration, the very order of quantities that we are considering. In fact, according to the reasoning in my paper, to which Prof. Challis does not appear to object, I found that the law of aberration does not result from supposing the waves of light to be carried by the moving æther, so long as its motion is taken arbitrary; and in order to explain aberration, I was *compelled* to suppose  $udx + vdy + wdz$  to be an exact differential, at least when the square of the aberration is neglected.

It is evidently immaterial whether we make the construction that Prof. Challis has given, or suppose  $ef$  to be the position into which the wave  $ab$  would come at the end of the



time  $t$ , in consequence of the velocity of propagation combined with the velocity of the æther at  $a$ , and suppose that  $f$  is brought to  $d$  in consequence of the difference of velocity of the æther at  $a$  and  $b$ . It is easy to show that  $df$  is equal and parallel to  $ce$ ; so that, according to *this* construction, the normal to the wave ought to be displaced by the motion of the æther through the angle  $fb d$  from  $fb$  to  $db$ , which is just the contrary direction to that given by Prof. Challis's construction.

Prof. Challis seems to think that the undulatory theory of light cannot be maintained unless it can be shown that the law of aberration ought to be the actual law, *whatever* may be the motion of the æther. But it is surely sufficient to show that a conceivable kind of motion exists which would lead to the observed law of aberration, provided we have no reason for regarding that sort of motion as improbable. Now even were I to allow that  $u dx + v dy + w dz$  cannot, in the case of ordinary fluids, be an exact differential unless the motion is rectilinear, that would not be a fatal objection. For the equations of motion of fluids commonly employed are formed on the hypothesis that the mutual action of two elements of the fluid is normal to the surface which separates them, whereas one of the most remarkable properties of the æther with which we are acquainted, is the great tangential force which it is capable of exerting, in consequence of which the transversal vibrations which constitute light are propagated with such an immense velocity.

## VI. On the Solubility of Oxide of Lead in Pure Water.

By Lieut.-Col. PHILIP YORKE\*.

**I**N the Philosophical Magazine for August 1834, I published a paper on the action of water and air on lead. Some of the principal results contained in it were confirmed by Bonsdorff in two papers; he found that 7000 parts of pure water free from access of carbonic acid dissolved one of oxide of lead; my experiments gave  $\frac{1}{12,000}$ th to  $\frac{1}{10,000}$ th. Since that time two papers have appeared on the same subject, one by Dr. Christison†, and one by Mr. R. Phillips, Jun.‡ The last-named chemist considers that the oxide of lead is not dissolved, but merely mechanically suspended in the water, because the liquid is deprived of the lead by passing it through

\* Communicated by the Chemical Society; having been read May 17, 1845.

† Transactions of the Royal Society of Edinburgh.

‡ Chemical Gazette for Jan. 1, 1845.

a paper filter. It is to this opinion that I propose to direct attention in the present notice.

The fact that the aqueous solution of oxide of lead would not pass through a filter was noticed by me in the paper already referred to; but as the action of tests on the liquid was just what one observes with solutions; as no time allowed for subsidence made any difference in these appearances; as the liquid deposited crystals of oxide of lead not only on the lead but on other bodies; as when decomposed by the voltaic battery it gave metallic lead at the negative pole, and peroxide at the positive; I did not consider that the stoppage of the oxide of lead by the filter was any proof of its not being dissolved. There still, however, remains this question to be answered,—In what way does the paper act in retaining the oxide? and I think that the following experiments afford an answer to the question.

I placed some clean rods of lead in bottles of distilled water loosely stoppered; in this way, after removing the rods of lead, I obtained a clear liquid, which, when tested by sulphuretted hydrogen, gave a deep brown colour. On passing this liquid through a double filter, which had been previously washed with hot distilled water, it appeared to be very nearly deprived of lead: when two or three fluid ounces had passed through, the filters were removed, washed, then immersed in a solution of sulphuretted hydrogen, again washed and dried. Some torn fragments of the filters were then mounted in Canada balsam for examination by the microscope. On examination with powers of from 150 to 400, the fibres of the flax composing the paper were seen to be browned, and in many instances it could be distinctly observed that the colouring substance occupied the interior of the tubular fibre. Now, it is stated by Mr. Crum, in the *Philosophical Magazine* for April 1844, that cotton wool possesses the power of abstracting the oxide of lead from its solution in lime-water, and that this property is made available in the processes for dyeing cotton with the chromates. I found that on filtering a solution of oxide of lead in lime water through a triple filter, that whereas the original solution gave a deep black when tested by sulphuretted hydrogen, the filtered liquid gave but a pale brown; and it required that the unfiltered liquid should be diluted with thirty times its volume of water to produce the same test as the filtered.

I then tried the effect of mere immersion of the paper in the aqueous solutions before used. A bit of filtering-paper ten inches by two inches was boiled in distilled water and then put into an ounce phial filled with the aqueous solution; after



remaining six hours the liquid was poured off and tested: it gave a pale brown, and it required that the liquid which had not been in contact with the paper should be diluted with ten times its volume of water to produce the same tint. This experiment was repeated with a stronger solution of oxide of lead in water, the water was poured off at the end of four hours; it then gave a pale brown, and it required that the original liquid should be diluted with four times its bulk of water to produce the same tint. A fresh portion of the same solution was then poured on the same paper and left for a night; then, on testing, the liquid gave a brown tint, barely perceptible, and it required that the original liquid should be diluted with from fifteen to twenty times its volume of water to produce the same.

From these experiments it is clear that the effect in question is dependent on a power possessed by the paper in common with several other porous bodies and organised fibres, of separating certain substances from their solutions, a power sufficiently well known, though little understood\*. In considering this view of the subject in the present instance, there is a circumstance of some practical importance which it would appear ought to follow, viz. that after the fibres of the paper had been saturated with the oxide of lead, then this substance should pass through in solution. To ascertain whether this was the case I made the following experiments.

I obtained a strong aqueous solution of oxide of lead by immersing slips of clean lead in about three quarts of distilled water, contained in a two-necked bottle, through which oxygen gas was passed and maintained in contact with, under a slight pressure. In this manner I procured a solution which when quite clear yielded  $\frac{1}{7500}$ th of ignited oxide of lead. A filter of paper rather less than  $\frac{1}{200}$ th of an inch thick and four inches in diameter was prepared and washed; then, by fitting into one of the two necks of the bottle a siphon with equal legs, so as to resemble Gay-Lussac's apparatus for washing filters (except that I used a contrivance to prevent the necessity of the air supplied to the bottle from bubbling through the solution), I was enabled to allow the filtration to go on with considerable regularity for many hours. The first portion of liquid which passed through gave a pale brown when tested; when nine fluid ounces had passed through the effect was the same as at first, and a portion (*a*) was reserved for future comparison. When forty fluid ounces had passed through, the

\* The effective filter mentioned by Dr. Clark is formed of well-washed sand, and has been in use during twelve months without any apparent diminution of power.

liquid, which was quite clear, gave a much darker tint with the test than any which had previously been obtained in the experiment. It gave a tint about equal to that given with the unfiltered liquid when diluted with its own volume of water; while it (*i. e.* the last filtered portion) required to be diluted with twice its volume of water to produce the same tint as that given by the reserved filtered portion (*a*). The liquid now passed through the filter very slowly; it was tested again, when eight more fluid ounces had passed through, with the same result as before, except that the tint was a trifle darker.

This experiment sufficiently shows that the effect contemplated does occur, and that it would be unsafe to trust to the action of a filter to separate oxide of lead from water for an unlimited time.

VII. *Equations for the Determination of the Motion of a Disturbed Planet by means of M. Hansen's Altered Time.*  
By the Rev. BRICE BRONWIN\*.

THE theory of M. Hansen on Lunar and Planetary Perturbations, owing to the two times  $\tau$  and  $t$  which it contains, is attended with many difficulties, and is very perplexing. His results I think are in an advantageous form; but perhaps they might be obtained more easily by the equations given in this paper, which are referred to the plane of the orbit as if it were a fixed plane, because I have proved that so referred they are true. [See this Magazine for November 1844, and also the Cambridge Mathematical Journal, No. 24.]

The equation

$$\frac{h^2}{\mu r} = 1 + e \cos(v - \pi) = 1 + e \cos \pi \cos v + e \sin \pi \sin v$$

is true for the disturbed orbit;  $h$ ,  $e$ , and  $\pi$  having their known variable values depending on the disturbing force. If  $h_0$ ,  $e_0$ , and  $\pi_0$  be the values of these quantities when the disturbing force is made to vanish, then

$$e \cos \pi = e_0 \cos \pi_0 + \int (\cos \pi de - e \sin \pi d\pi),$$

$$e \sin \pi = e_0 \sin \pi_0 + \int (\sin \pi de + e \cos \pi d\pi),$$

$$h^2 = h_0^2 + 2 \int h dh.$$

These values substituted in the above equation give

\* Communicated by the Author.

$$\frac{h_0^2}{\mu r} = 1 + e_0 \cos(v - \pi_0) - \frac{2}{\mu r} \int h \, dh$$

$$+ \cos v \int (\cos \pi \, de - e \sin \pi \, d\pi) + \sin v \int (\sin \pi \, de + e \cos \pi \, d\pi).$$

Let the constant quantities  $\lambda$  and  $\varrho$  be the same functions of a constant time  $\tau$  which  $v$  and  $r$  are of  $t$ ; then putting the former in place of the latter, we may put them under the sign of integration, changing  $\tau$  into  $t$  after the integrations are performed. This will change the last equation into

$$\frac{h_0^2}{\mu r} = 1 + e_0 \cos(v - \pi_0) - \frac{2}{\mu} \int \frac{h \, dh}{\varrho} + \int \cos(\lambda - \pi) \, de$$

$$+ \int e \sin(\lambda - \pi) \, d\pi.$$

But 
$$\int h \, dh = - \int \frac{dR}{dv} h \, dt,$$

$$de = - \frac{h \, dt}{\mu} \sin(v - \pi) \frac{dR}{dr} - \left( \frac{2h \, dt}{\mu r} \cos(v - \pi) + \frac{dr}{\mu} \sin(v - \pi) \right) \frac{dR}{dv},$$

$$d\pi = \frac{h \, dt}{\mu e} \cos(v - \pi) \frac{dR}{dr} - \left( \frac{2h \, dt}{\mu r e} \sin(v - \pi) - \frac{dr}{\mu e} \cos(v - \pi) \right) \frac{dR}{dv}.$$

The coefficients of  $\frac{dR}{dv}$  are put under the above form for convenience. Substituting these values, we find

$$\frac{1}{r} = \frac{\mu}{h_0^2} + \frac{\mu e_0}{h_0^2} \cos(v - \pi_0) + \frac{1}{h_0^2} \int \frac{dR}{dr} \sin(\lambda - v) h \, dt$$

$$+ \frac{1}{h_0^2} \int \frac{dR}{dv} \left\{ \sin(\lambda - v) dr - \frac{2h \, dt}{r} \cos(\lambda - v) + \frac{2h \, dt}{\varrho} \right\}.$$

To abridge, let this be written

$$\frac{1}{r} = \frac{\mu}{h_0^2} + \frac{\mu e_0}{h_0^2} \cos(v - \pi_0) + P.$$

But if  $\varrho t$  be the progression of the apse,

$$\cos(v - \pi_0) = \cos(v - \varrho t - \pi_0 + \varrho t) = \cos(v - \varrho t - \pi_0)$$

$$- \varrho t \sin(v - \varrho t - \pi_0),$$

neglecting higher powers of  $\varrho t$ . Therefore

$$\frac{1}{r} = \frac{\mu}{h_0^2} \{ 1 + e_0 \cos(v - \varrho t - \pi_0) - e_0 \varrho t \sin(v - \varrho t - \pi_0) \} + P.$$

Terms similar to the above, containing  $t$  in their coefficients, will arise from the development of  $P$ ; and  $\varrho$  must be so determined as to drive them out, which will be easily done. We may always neglect terms involving the higher powers of  $t$ .

Let  $r_1$  and  $v_1$  be the same functions of the new time  $\zeta$ , and the constants  $h_1, e_1, \pi_0, \epsilon_0$  which  $r$  and  $v$  are, when there is no disturbing force, of the time  $t$  and the constants  $h_0, e_0, \pi_0$ , and  $\epsilon_0$ . Also let  $v = v_1 + \epsilon t$ . We shall have

$$\frac{1}{r_1} = \frac{\mu}{h_1^2} \{1 + e_1 \cos(v_1 - \pi_0)\}.$$

We shall not with M. Hansen find the log of  $r$ , and therefore shall make  $\frac{1}{r} = \frac{1}{r_1} + p$ . Substituting this value, we shall easily find

$$p = \mu \left( \frac{1}{h_0^2} - \frac{1}{h_1^2} \right) + \mu \left( \frac{e_0}{h_0^2} - \frac{e_1}{h_1^2} \right) \cos(v_1 - \pi_0) - \frac{e_0 \epsilon t}{h_0^2} \sin(v_1 - \pi_0) + P.$$

Whence  $p$  is of the order of the disturbing force, and it has the advantage of requiring only one integration.

In virtue of the supposed equation

$$r_1^2 dv_1 = h_1 d\zeta, \text{ or } \frac{dv_1}{d\zeta} = \frac{h_1}{r_1^2},$$

we have

$$\frac{dv}{dt} = \frac{dv_1}{dt} + \epsilon = \frac{dv_1}{d\zeta} \frac{d\zeta}{dt} + \epsilon = \frac{h_1}{r_1^2} \frac{d\zeta}{dt} + \epsilon.$$

This value, substituted in the known equation

$$\frac{dv}{dt} = \frac{1}{r^2} \left( h_0 - \int \frac{dR}{dv} dt \right)$$

gives

$$\frac{d\zeta}{dt} = (1 + r_1 p)^2 \left( \frac{h_0}{h_1} - \frac{1}{h_1} \int \frac{dR}{dv} dt \right) - \frac{\epsilon r_1^2}{h_1}.$$

Of the four quantities  $h_0, h_1, e_0, e_1$ , two are to be found in terms of the others, which will be arbitraries of the theory; and the mode of determining them will be obvious after the development is effected.

Putting  $\phi$  for the latitude,  $i$  for the inclination,  $\vartheta$  and  $\theta$  for the longitude of the node on the plane of the orbit and on the fixed plane, we have

$$\begin{aligned} \vartheta &= \int \cos i d\theta, \quad \sin \phi = \sin i \sin(v - \vartheta) \\ &= \sin i (\cos \vartheta \sin v - \sin \vartheta \cos v), \end{aligned}$$

$$\sin i \cos \vartheta = \sin i_0 \cos \vartheta_0 + \int (\cos i \cos \vartheta di - \sin i \sin \vartheta d\vartheta),$$

$$\sin i \sin \vartheta = \sin i_0 \sin \vartheta_0 + \int (\cos i \sin \vartheta di + \sin i \cos \vartheta d\vartheta).$$

Substituting these values, and changing  $v$  into  $\lambda$ , and putting it under the sign of integration, we obtain

$$\begin{aligned} \sin \phi &= \sin i_0 \sin (v - \vartheta_0) + \int \sin (\lambda - \vartheta) \cos i di \\ &\quad - \int \cos (\lambda - \vartheta) \sin i d\vartheta. \end{aligned}$$

But  $di = \frac{dt}{h \sin i} \frac{dR}{d\theta}$ ,  $d\vartheta = -\frac{\cos i dt dR}{h \sin i di}$ .

These values being put in the above, it will become

$$\begin{aligned} \sin \phi &= \sin i_0 \sin (v - \vartheta_0) + \int \frac{\cos i dt dR}{h \sin i di} \sin (\lambda - \vartheta) \\ &\quad + \int \frac{\cos i dt dR}{h di} \cos (\lambda - \vartheta). \end{aligned}$$

To abridge, we may write this

$$\sin \phi = \sin i_0 \sin (v - \vartheta_0) + Q,$$

or  $\sin \phi = \sin i_0 \sin (v + \gamma t - \vartheta_0) - \gamma t \sin i \cos (v + \gamma t - \vartheta_0) + Q.$

Here  $\gamma t$  is the regression of the node, and  $\gamma$  is to be so determined as to take away from  $Q$  the terms having  $t$  in their coefficients.

If in the equations

$$\begin{aligned} \sin \phi &= \sin i \sin (v - \vartheta), \\ \frac{d \sin \phi}{dt} &= \sin i \cos (v - \vartheta) \frac{dv}{dt}, \end{aligned}$$

we change  $\phi$ ,  $v$ ,  $i$ , and  $\vartheta$  into  $\phi_0 + \delta \phi$ ,  $v_0 + \delta v$ ,  $i_0 + \delta i$ , and  $\vartheta_0 + \delta \vartheta$ ,  $\delta \phi$ , &c. being the parts depending on the disturbing force; and if we expand, taking account of the first power only of  $\delta \phi$ , &c., we shall find equations of the form

$$\begin{aligned} \delta i &= A \delta \phi + B \delta v, \\ \delta \vartheta &= C \delta \phi + D \delta v. \end{aligned}$$

From these we may correct the values of  $i$  and  $\vartheta$  or  $\theta$  by means of the corrections of  $\phi$  and  $v$  due to the disturbing force, and in doing so we may take account of higher powers of  $\delta \phi$ , &c.

I think the development of the preceding equations would be attended with much less difficulty and perplexity than the development of M. Hansen's. I have not noticed the reduction to a fixed plane, but must refer for that to the Number of this Magazine for February 1844, where I have given equations particularly adapted to the lunar theory, and leading to results expressed in terms of the true elliptic longitude.

I see Mr. Cayley has amended his paper of November 1844. If he would amend it a little further, it would not be amiss. He has now made  $p$  a prime number instead of any odd one:  $\theta$  is made of the second instead of the general form. In the expression of  $\phi, x$ , or rather  $\phi x'$ , he should have had the transformed function  $x' = \frac{x}{\beta}$ , not  $x, \beta$  a function of  $\theta$ . Moreover,

$$\omega' = \frac{1}{p\beta} (\alpha\omega + \beta v), \quad v' = \frac{1}{p\beta} \alpha'\omega + \beta'v.$$

Some other amendments are greatly wanted. When  $x$  has a determinate value,  $x'$  should have one also, since  $\beta$  is a known function of  $\theta$ . And if we know what value to assign to  $x$ , we should have the value of  $x'$ , which is the complete function. I have no room to enlarge, and as Mr. Cayley has done nothing which invalidates what I have done, it is unnecessary.

Gunthwaite Hall, Dec. 12, 1845.

VIII. *On some Points in the Meteorology of Bombay.*  
By Lieut.-Colonel SABINE, R.A., F.R.S.\*

[With a Plate.]

**I**N a communication which I had the honour to make to the Section at the York meeting of the British Association, on the subject of the meteorological observations made at Toronto in Canada in the years 1840 to 1842†, I noticed some of the advantages which were likely to result to the science of meteorology, from the resolution of the barometric pressure into its two constituents of aqueous and of gaseous pressure. It was shown that when the constituents of the barometric pressure at Toronto were thus disengaged from each other and presented separately, their annual and diurnal variations exhibited a very striking and instructive accordance with the annual and diurnal variations of the temperature. The characteristic features of the several variations when projected in curves were seen to be the same, consisting in all cases of a single progression, having one ascending and one descending branch; the epochs of maxima and minima of the pressures being the same, or very nearly the same, with those of the maxima and minima of temperature; and the correspondence in other respects being such as to manifest the existence of a very intimate connexion between the periodical variations of the temperature, and those of the elastic forces of the air and vapour. The curve of gaseous pressure was inverse in respect to the other two; that is to say, as the temperature increased the elastic force of the vapour increased also, but that of the air diminished, and *vice versa*; and this was the case both in the annual and the diurnal variations.

\* Communicated to the Mathematical and Physical Section of the British Association for the Advancement of Science, at the Cambridge Meeting, 1845.

† See Phil. Mag., vol. xxvi. p. 94.

Such being the facts, I endeavoured to show, in the case of the diurnal variations, that the correspondence of the phænomena of the temperature and gaseous pressure might be explained, in accordance with principles which had been long and universally admitted in the interpretation of other meteorological phænomena, by the suppositions,—of an extension in height and consequent overflow in the higher regions of the atmosphere of the column of air over the place of observation, during the hours of the day when the surface of the earth was gaining heat by radiation,—and of a contraction of the column during the hours of diminishing temperature, and consequent reception of the overflow from other portions of the atmosphere, which in their turn had become heated and elongated.

According to this explanation there should exist, during the hours of the day when the temperature is increasing,—1st, an *ascending current* of air at the place of observation, of which the strength should be measured by the amount of the increments of temperature corresponding to given intervals of time; and 2nd, a *lateral influx of air at the lower parts of the column*, of proportionate velocity, constituting a *diurnal variation in the force of the wind* at the place of observation, which should also correspond with the variations of the temperature in the epochs of its maximum and minimum, and intermediate gradation of strength. The anemometrical observations at Toronto were shown to be in agreement with the view which had been then taken, confirming the existence of a diurnal variation in the force of the wind, corresponding in all respects with the variation of the temperature.

Admitting the explanation thus offered to be satisfactory in regard to the diurnal variations, it was obvious that the correspondence of the annual variations of the temperature and pressures might receive an analogous explanation.

A comparison of the results of the observations at Toronto with those of the observations of M. Kreil at Prague in Bohemia, (published in the *Mag. und Met. Beob, zu Prag*, and in the *Jahrbuch für Prag*. 1843,) showed that the characteristic features of the periodical variations at Toronto were not peculiar to that locality, but might rather be considered as belonging to stations situated in the temperate zone and in the interior of a continent. The annual and diurnal variations at Prague were also single progressions, and the same correspondence was observable between the variations of the temperature and of the gaseous pressure.

The publication of the volume of magnetical and meteorological observations made at Greenwich in 1842, which took place shortly after the meeting of the Association at York, enabled me to add a postscript to the printed statement of my communication in the annual volume of the Association Reports, showing the correspondence of the results at Greenwich with the relations which had been found to exist in the periodical march of the phænomena at Toronto and at Prague.

From the concurrence of these three stations, it was obvious that a considerable insight had been obtained into the laws which regulate

the periodical variations in the temperate zone, and into the sequence of natural causes and effects, in accordance with which the annual and diurnal fluctuations of the elastic forces of air and vapour at the surface of the earth depend on the variations of temperature: and from these premises it was inferred, that the normal state of the diurnal variations of the pressures of the air and vapour and of the force of the wind, in the temperate zone, might be regarded as that of a single progression with one maximum and one minimum, the epochs of which should nearly coincide with those of the maximum and minimum of temperature\*.

That exceptions should be found to this state of things in particular localities in the temperate zone was far from being impro-

\* Since this communication was read at Cambridge I have received from M. Dove a copy of a paper read to the Academy of Berlin, entitled 'Ueber die periodischen Aenderungen des Druckes der Atmosphäre im Innern der Continente,' in which the remarkable facts are stated, that at Catherinenbourg and Nertchinsk (on the mean of several years), and at Barnaoul (in the years 1838 and 1840), the mean diurnal *barometric* curve itself exhibits but one maximum and one minimum in the twenty-four hours; the maximum coinciding nearly with the coldest, and the minimum with the hottest hours of the day. At these stations therefore, and in the years referred to, the forenoon maximum disappeared, and the barometric curve assimilated in character to the curve of the dry air in other places in the temperate zone. These stations are situated far in the interior of the greatest extent of dry land on the surface of our globe, and at a very great distance from an expanse of water, from whence vapour can be supplied. The diminished pressure of the dry air produced by the ascending current and overflow as the temperature of the day increases, is not therefore compensated by an increased elasticity of vapour, and the curve of the diurnal variation of the barometer approximates to the form assumed when the elasticities of the vapour at the several hours of observation are abstracted. This assimilation in character of the barometric and (inferred) gaseous curves, which is thus found to take place in cases where, from natural causes, the influence of the vapour is greatly lessened, appears a confirmation of the propriety of separating the effects of the elastic forces of the dry air and vapour in their action on the barometer.

M. Dove considers that the single progression of the diurnal barometric curve, which takes place at the three Asiatic stations referred to in this note, is characteristic of a true continental climate. It is, without doubt, characteristic of an extreme climate, and as such is highly instructive. There appears reason to doubt whether an extreme climate of corresponding character exist at all in the temperate latitudes of the continent of America.

If, however, we examine the record of the observations made hourly in the year 1842 at Catherinenbourg, Barnaoul and Nertchinsk, in the 'Annuaire Magnétique et Météorologique de Russie,' we find that at Catherinenbourg in that year the barometer exhibits a double progression, but that the morning maximum, which occurs at the observation hour of 8<sup>h</sup> 22<sup>m</sup> A.M., exceeds the antecedent minimum only by a quantity less than 0.003 in. At Barnaoul there is also a double progression in the barometric mean in that year, the morning maximum being still small, and taking place between the observation hours of 9<sup>h</sup> 54<sup>m</sup> and 10<sup>h</sup> 54<sup>m</sup> A.M. At Nertchinsk also there is a morning maximum occurring at the observation hour of 9<sup>h</sup> 17<sup>m</sup> A.M. In all the three cases the double progression shown by the barometer disappears wholly in the curve of the dry air, which curve exhibits at these three stations, as well as at Toronto, Prague and Greenwich, but one maximum and one minimum in the twenty-four hours. At the three stations of extreme dryness cited by M. Dove, therefore the vapour was still sufficient to impart, in the year 1842 at least, a double progression to the diurnal variation of the barometer; but the hour of the morning maximum was earlier than where the increase of vapour, as the day advances, is greater.



bable ; it could not be expected that the influences of temperature should always be so simple and direct as they appeared to be at Toronto ; and a more complex aspect of the phænomena might particularly be looked for, where a juxtaposition should exist of columns of air resting on surfaces differently affected by heat (as those of land and sea), and possessing different retaining and radiating properties. In such localities *within the tropics*, the well-known regular occurrence of land and sea breezes for many months of the year made it obvious that a double progression in the diurnal variation of the force of the wind must exist, and rendered it highly probable that a double progression of the gaseous pressure would also be found. It was therefore with great pleasure that I received, through the kindness of Dr. Buist, a copy of the monthly abstracts of the two-hourly meteorological observations, made under that gentleman's superintendence at the observatory at Bombay in the year 1843 ; accompanied by a copy of his meteorological report for that year, possessing a particular value, in the full account which it gives of the periodical variations of the wind, and in the means which it thereby affords of explaining the diurnal variation of the gaseous pressure. This pressure presents at Bombay an aspect at first sight more complex than at the three above-named stations in the temperate zone, but I believe it to be equally traceable to variations of the temperature, and to furnish a probable type of the variations at inter-tropical stations similarly circumstanced in regard to the vicinity of the sea.

The observatory at Bombay is situated on the island of Colabah, in N. lat.  $18^{\circ} 54'$  and E. long.  $72^{\circ} 50'$  at an elevation of thirty-five feet above the level of the sea. In the copy of the observations received from Dr. Buist, the monthly abstracts are given separately for each month, of the standard thermometer,—of the wet thermometer, and of its depression below the dry,—and of the barometer. In Table I. I have brought in one view the thermometrical and barometrical means at every second hour, and the mean tension of the vapour and mean gaseous pressure at the same hours. The tension of the vapour at the several observation hours has been computed from the *monthly means*, at the same hours, of the wet thermometer and of its depression below the dry thermometer. The values are consequently somewhat less than they would have been, had the tension been computed from each individual observation of the wet and dry thermometers, and had the mean of the tensions thus obtained been taken as the value corresponding to the hour. The difference is however so small, that for the present purpose it may be regarded as quite insignificant. It would not amount in a single instance to the hundredth part of an inch ; and as in every instance the difference would be in the same direction, the *relative* values, which are those with which we are at present concerned, would be scarcely sensibly affected. The pressures of the dry air (or the gaseous pressures) are obtained by deducting the tension of the vapour from the whole barometric pressure.

TABLE I.

Bombay, 1843.—Mean Temperature, Mean Barometric Pressure, Mean Tension of Vapour, and Mean Gaseous Pressure at every second hour.

Hours of Mean Bombay Time. Astronomical Reckoning.	Temperature.	Barometer.	Tension of Vapour.	Gaseous Pressure.
	°	in.	in.	in.
18	78·4	29·805	0·750	29·055
20	79·6	29·840	0·766	29·074
22	81·8	29·852	0·771	29·081
0	83·2	29·817	0·768	29·049
2	84·1	29·776	0·795	28·981
4	83·9	29·755	0·800	28·955
6	82·3	29·774	0·802	28·972
8	81·2	29·806	0·801	29·005
10	80·3	29·825	0·780	29·045
12	79·8	29·809	0·775	29·034
14	79·4	29·786	0·766	29·020
16	78·9	29·778	0·761	29·017
Mean of the year ...	81·1	29·802	0·780	29·022

The sun is vertical at Bombay twice in the year, viz. in the middle of May and towards the end of July. The rainy season sets in about the commencement of June (in 1843 on the 2nd of June), and terminates in August, but with heavy showers of no long duration continuing into September. During the rainy season, and in the month of May which immediately precedes it, the sky is most commonly covered with clouds, by which the heating of the earth by day, and its cooling at night by radiation, are impeded, and the range of the diurnal variation of the temperature is greatly lessened in comparison with what takes place at other times in the year. The strength of the land and the sea breezes in those months is also comparatively feeble, and on many days the alternation of land and sea breeze is wholly wanting. During the months of November, December, January and February, the diurnal range of the temperature is more than twice as great as in the rainy season, and the land and sea breezes prevail with the greatest regularity and force.

In addition to the monthly tables, we may therefore advantageously collect in one view, for purposes of contrast, the means of the months of May, June, July and August, as the season when the sky is generally clouded,—and of the months of November, December, January and February, as the season of opposite character, when the range of the diurnal temperature is greatest, and the land and sea breezes alternate regularly, and blow with considerable strength. These seasons are contrasted in Table II.

If we direct our attention to the diurnal variations, commencing with those of the temperature, we find them exhibiting a single progression, having a minimum at 18<sup>h</sup> and a maximum at 2<sup>h</sup>; the average difference between the temperature at 18<sup>h</sup> and 2<sup>h</sup> being 7°·77

in the clear season,  $3^{\circ}\cdot71$  in the clouded season, and  $5^{\circ}\cdot7$  on the mean of the whole year.

When however we direct our attention to the gaseous pressure, we perceive, very distinctly marked, the characters of a double progression, having one maximum at  $10^{\text{h}}$  and another at  $22^{\text{h}}$ ; one minimum at  $4^{\text{h}}$  and another at  $16^{\text{h}}$ . The double progression is exhibited both in the clouded and in the clear seasons, with a slight difference only in the hours of maxima; the principal maximum in the cloudy season being at  $20^{\text{h}}$  instead of  $22^{\text{h}}$ , and the inferior maximum in the clear season being at  $12^{\text{h}}$  instead of  $10^{\text{h}}$ . The range of the diurnal variation, like that of the temperature, is more than twice as great in the clear as in the clouded season, marking distinctly the connexion subsisting between the phænomena of the temperature and of the gaseous pressure.

TABLE II.

Bombay, 1843.—Comparison of the Temperature and of the Gaseous Pressure in the months of May, June, July and August, when the sky is usually covered with clouds; and in November, December, January and February, when the sky is usually clear.

Hours of Mean Time at Bombay. Astronomical Reckoning.	Temperature.		Gaseous Pressure.	
	November, December, January and February.	May, June, July and August.	November, December, January and February.	May, June, July and August.
			in.	in.
18	74·1	81·9	29·344	28·782
20	75·3	83·1	29·368	28·806
22	78·1	84·3	29·391	28·798
0	80·8	85·1	29·353	28·782
2	81·9	85·6	29·230	28·746
4	81·7	85·4	29·195	28·724
6	79·6	84·3	29·199	28·740
8	78·4	83·4	29·248	28·754
10	76·9	83·0	29·308	28·800
12	76·2	82·7	29·316	28·775
14	75·7	82·6	29·295	28·754
16	74·9	82·2	29·285	28·753
Means .....	77·8	83·7	29·298	28·763

If we now turn our attention to the phænomena of the direction and force of the wind, we find by Dr. Buist's report, that for 200 days in the year there is a regular alternation of land and sea breezes. The land breeze springs up usually about  $10^{\text{h}}$ , or between  $10^{\text{h}}$  and  $14^{\text{h}}$ , blows strongest and freshest towards daybreak, and gradually declines until about  $22^{\text{h}}$ , at which time the direction of the aerial currents changes, and there is generally a lull of an hour or an hour and a half's duration. The sea breeze then sets in, the ripple on the surface of the water indicating its commencement being first observed close in shore, and extending itself gradually out to sea. The sea breeze is freshest from  $2^{\text{h}}$  to  $4^{\text{h}}$ , and progressively declines in the evening hours.

The diurnal variation in the force of the wind during these 200 days is therefore obviously a double progression, having two maxima and two minima; one maximum at or near the hottest, and the other at or near the coldest hour of the day,—being the hours when the difference of temperature is greatest between the columns of air which rest respectively on the surfaces of land and sea; and two minima coinciding with the hours, when the surface temperature over the land and over the sea approaches nearly to an equality.

In the remaining portion of the year the diurnal range of the temperature is most frequently insufficient to produce that alternation in the direction of the wind, which prevails uninterruptedly during the larger portion. There appears however to have been only one month, viz. July, in the year 1843, in which there were not some days in which the alternation of land and sea breezes was perceptible. The causes which produce the alternation are not therefore wholly inoperative, though the effects are comparatively feeble during the clouded weather which accompanies the south-west monsoon\*.

If we now view together the diurnal variations of the wind and gaseous pressure, as shown in Plate I., we find a minimum of pressure coinciding with the greatest strength of the sea breeze; a second minimum of pressure coinciding with the greatest strength of the land breeze; and a maximum of pressure at each of the periods when a change takes place in the direction of the aerial currents; or, otherwise stated, we find a decrease of pressure coincident with the increase of strength both of the land and sea breezes, and an increase of pressure coincident with their decline in strength.

The facts thus stated appear to me to admit of the following explanation:—the diminution of pressure which precedes the minimum at 4<sup>h</sup> is occasioned by the rarefaction and ascent of the column during the heat of the day, and its consequent overflow in the higher regions of the atmosphere, which is but partially counterbalanced in the forenoon by the influx of the sea breeze at the lower part of the column. Shortly after the hottest hour is passed, the overflow above and the supply below become equal in amount, and the diminution of pressure ceases. As the temperature falls towards evening, the column progressively contracts, when the influx from the sea breeze more than counterbalances the overflow, and the pressure again increases until a temporary equilibrium is restored, when the sea breeze ceases and the pressure is stationary.

As the night advances, the air over the land becomes colder than over the sea; the length of the column over the land contracts, and the air in its lower part becomes denser than in that over the sea: an interchange then commences of an opposite character to that which prevailed during the day. The outward flow is now from the *lower* part of the column, or from the land towards the sea,

\* There are no data in Dr. Buist's report from which the diurnal variation in the force of the wind may be judged of in the days during the south-west monsoon, when no alternation takes place in its direction. It would seem probable that on such days the variation should be a single progression, weakest towards daybreak, and strongest about the hottest hour of the day.

causing the pressure to diminish over the land; it continues to do so until towards daybreak, when the strength of the land breeze is greatest, because the air over the land is then coldest in comparison with that over the sea. As the sun gains in altitude and the temperature of the day advances, the land heats rapidly; the density of the air over the land and sea returns towards an equality; the land breeze declines in strength, and the drain from the lower part of the column ceases to counterbalance the overflow which the land column is at the same time receiving in the higher regions; the pressure consequently having attained a second minimum at or near the hour of the greatest disproportion of temperature, again increases until the temperature and height of the column over the sea and land are the same, and the pressure again becomes stationary. But now the rarefaction of the column over the land continuing, its increase in height above the less heated column with which it is in juxtaposition, and its consequent overflow, occasion the pressure to decrease until the minimum at 4 o'clock is reached.

We have thus therefore at Bombay a *double progression of the diurnal variation of the gaseous pressure*; the principal minimum occurring at 4 o'clock in the afternoon, occasioned by an overflow from the column in the higher regions of the atmosphere; and the second minimum occurring at 18<sup>h</sup>, occasioned by an efflux from the lower part of the column. The first minimum is similar to that which has been shown to take place at Toronto, Prague and Greenwich, and is similarly explained: the second minimum, which does not take place at the three above-named stations, is owing to the juxtaposition of the columns of air over the sea and land, which differ in temperature, and therefore in density and height, in consequence of their resting respectively on surfaces which are differently affected by heat.

Plate I. shows the curve of the gaseous pressure, and the curve of the elastic force of the vapour; and between them is placed a diagram illustrating the hours of prevalence and of the greatest strength of the land and sea breezes. At Toronto and at Greenwich the diurnal curve of the vapour is a single progression, having its maximum at or near the hottest hour of the day, and its minimum at or near the coldest hour. We perceive in the Plate which represents the phenomena at Bombay, the modification which takes place in consequence of the supply of vapour brought in by the sea breeze continuing until a late hour in the evening, and prolonging the period during which the tension is at or near its maximum. The minimum occurs as usual at or near the hour of the coldest temperature.

If, then, the explanation which I have offered to the Section, of the physical causes which produce the diurnal variation of the gaseous pressure at Bombay, be correct, the diurnal variation of the barometric pressure occurring there is also explained, since it is simply the combination of the two elastic forces of the air and of the vapour.

The solution of the problem of the diurnal variation of the barometer is therefore obtained by the resolution of the barometric pres-

sure into its constituent pressures of vapour and air; since the physical causes of the diurnal variation of the component pressures have been respectively traced to the variations of temperature produced in the twenty-four hours by the earth's revolution on its axis, and to the different properties possessed by the material bodies at the surface of the globe in respect to the reception, conveyance, and radiation of heat.

*Annual variation.*—We now proceed to the annual variations, which are shown in the subjoined table.

TABLE III.

1843.	Tempera- ture.	Vapour Pressure.	Gaseous Pressure.	Barometer.	Humidity.	Monthly Means greater (+) or less (-) than the Annual Means.		
						Tempera- ture.	Vapour Pressure.	Gaseous Pressure.
January ...	76·4	0·578	29·352	29·930	67	-0·47	-0·202	+0·329
February ...	77·7	0·648	29·246	29·894	71	-3·4	-0·132	+0·223
March .....	79·7	0·710	29·128	29·838	74	-1·4	-0·070	+0·105
April .....	84·2	0·853	28·961	29·814	76	+3·1	+0·073	-0·062
May .....	85·9	0·921	28·743	29·664	78	+4·8	+0·141	-0·280
June.....	85·4	0·935	28·718	29·653	80	+4·3	+0·155	-0·305
July .....	82·1	0·896	28·737	29·633	85	+1·0	+0·116	-0·286
August .....	81·2	0·859	28·869	29·728	84	+0·1	+0·079	-0·154
September..	81·1	0·859	28·920	29·779	84	0·0	+0·079	-0·103
October ...	82·2	0·819	29·026	29·845	78	+1·1	+0·039	+0·003
November ..	80·5	0·675	29·213	29·888	67	-0·6	-0·105	+0·190
December ..	76·6	0·592	29·368	29·960	67	-4·5	-0·188	+0·345
	81·1	0·780	29·023	29·803	76			

We here perceive that the leading features of the phenomena are clearly analogous to those which have been seen to present themselves at Toronto, Prague and Greenwich; viz. a correspondence of the maximum of vapour pressure and minimum of gaseous pressure with the maximum of temperature,—and of the minimum of vapour pressure and maximum of gaseous pressure with the minimum of temperature; and a progressive march of the three variations from the minimum to the maximum, and back to the minimum again. The epochs, or turning-points of the respective phenomena, are not in every case strictly identical; but their connexion, which is the subject immediately before us, is most obvious.

We have thus a further illustration of the universality of the principle of the dependence of the regular periodical variations, annual as well as diurnal, of the pressures of the dry air and of the vapour, on those of the temperature\*.

\* In the tropics and in the temperate zone the heat of summer produces and accompanies a low gaseous pressure, and the cold of winter a high gaseous pressure. When we consider how large a portion of the northern hemisphere is occupied by land, which becoming greatly heated in summer rarefies the superincumbent atmosphere, causing it to overtop the adjacent less heated masses, and to overflow them, we should be led to expect that in parts of the Arctic Circle situ-

The humidity exhibits also a single progression; but may perhaps be rather characterized as evidencing a very dry season from November to February, and a very humid one from June to September.

ated to the north of the great continents, the gaseous pressure should be *increased* in summer, and that the curve of annual variation should become the converse of what it is in the lower latitudes. It appears from the meteorological observations made in 1843 by Messrs. Grewe and Cole, and presented to the British Association at the York meeting by Dr. Lee, that such is the case at Alten, near the north cape of Europe. The barometer and thermometer were observed three times a day, from October 1842 to December 1843 inclusive. The hours of observation were 9 A.M., 3 P.M. and 9 P.M. No hygrometric observations were made, but we are able to infer the approximate tension of the vapour from the record of the thermometer. The quarterly means of the barometer and thermometer in 1843 are as follows; the barometer being reduced to the level of the sea, and corrected for gravity:—

	Barometer. in.	Thermometer. ° F.
December, January, February.....	29·375	24
March, April, May .....	29·948	27·7
June, July, August .....	29·905	52·4
September, October, November ...	29·716	34·2
	<hr/>	<hr/>
Mean of the year .....	29·736	34·6

Assuming the humidity in each quarter of the year to be 75, or the vapour to be in each case three-fourths of that required for saturation at the respective temperatures, we should have the following gaseous pressures:—

December, January, February .....	29·257
March, April, May .....	29·804
June, July, August .....	29·616
September, October, November, December ...	29·566
	<hr/>
	29·561

It appears therefore that in the six summer months the mean barometric pressure exceeded that of the winter months by 0·381 inch; and the mean gaseous pressure of summer exceeded that of winter by about 0·3 inch. As in this case the curve of the gaseous pressure, as well as that of the aqueous vapour, accords in character with the curve of temperature, i. e. ascends with ascending temperature, and descends with descending temperature,—the barometric annual range is greater than the gaseous annual range, which is contrary to what takes place in the temperate and equatorial zones. It is not improbable that in the Antarctic Circle the phenomenon which we have just noticed as taking place in the Arctic Circle, viz. the summer increase of the gaseous pressure,—may not be found in the same degree, if at all; for the two hemispheres present a remarkable contrast in their respective proportions of sea and land, and the rarefaction of the atmosphere over the middle latitudes of the southern hemisphere during its summer must be greatly less than in the same latitudes of the northern hemisphere in the corresponding season. The barometrical observations made by Sir James Ross in summer in the Antarctic Circle accord with this inference; since after correcting them for the shortening of the column of mercury by the increased force of gravity in the high latitudes, and abstracting the small tension of vapour corresponding to the temperature, the mean gaseous pressure deduced from them, though nearly equal to the mean gaseous pressure of the year at Bombay, does not exceed it; whereas at Alten it is only in the winter months that the gaseous pressure descends so low as to approximate to the usual mean gaseous pressure of the tropical regions.

It is much to be desired that the zealous observers at Alten should observe the  
*Phil. Mag.* S. 3. Vol. 28. No. 184. Jan. 1846. D

ber, the latter season being that of the rains. The average degree of humidity in the year is very slightly lower than either at Toronto or at Greenwich, but is still closely approaching to a value expressing the presence of three-fourths of the quantity of vapour required for saturation.

The mean gaseous pressure in 1843, derived from the two-hourly observations, appears to have been ( $29\cdot023 + 0\cdot025$ , an index correction which Dr. Buist gives as that of the barometer with which the observations were made =)  $29\cdot048$  English inches; or, measured by the height of a mercurial column in the latitude of  $45^{\circ}$ ,  $28\cdot988$ . The height above the sea is thirty-five feet, and the latitude  $19^{\circ}$  N.

The mean height of the barometer in the year 1843, derived from observations at every second hour, appears to have been ( $29\cdot803 + 0\cdot025$  =)  $29\cdot828$ , or, with the correction applied for gravity,  $29\cdot768$ , the elevation being thirty-five feet above the sea. This is less than what is generally received as the average height of the barometer in the same latitude. From the careful comparison described in Dr. Buist's report of the standard barometer with several other barometers, there seems great reason to believe that the mean height shown by it must be a very near approximation at least to the true mean atmospheric pressure in the year 1843 at Bombay.

The mean height of the barometer in the four clouded months of May, June, July and August, is  $29\cdot667$ ; and in the four clear months of November, December, January and February,  $29\cdot921$ . The mean vapour pressure in the same seasons is respectively  $0\cdot904$  and  $0\cdot623$ , and the gaseous pressure consequently  $28\cdot763$  and  $29\cdot298$ . There is therefore between the two seasons a difference of  $0\cdot535$  in. of gaseous pressure, and of  $5^{\circ}\cdot84$  of temperature; the lowest pressure corresponding to the highest temperature, and *vice versa*. If we may allow ourselves to make a rough proportion drawn from a single case, we may estimate a decrement of  $0\cdot1$  in. of pressure to an increment of  $1^{\circ}$  F. The highest temperature and lowest pressure are accompanied for nearly the whole of the period by the southern monsoon; the lowest temperature and the highest pressure are accompanied by the northern monsoon.

The curves of the annual variation of the gaseous, barometric, and vapour pressures, which are represented in Plate I., show how much of the influence produced on the gaseous pressure, by the alternation of the overflow in the high regions of the atmosphere as either side of the equator becomes heated in its turn, is masked in the barometric curve by the combination in the latter of the vapour pressure, the variations of which take place throughout the year

wet thermometer at the same time as the barometer; the register would also be rendered much more complete by the addition of another observation-hour, about 6 A.M., which might not perhaps be inconvenient. The atmospheric pressure and the tension of vapour at or near the coldest hour of the twenty-four, are important data in meteorological discussions.



in the opposite direction to those of the gaseous pressure. From this cause the range of the barometric curve during the year is only 0·327 inch, whilst that of the gaseous pressure is 0·650 inch.

The analogy of the annual and diurnal variations, considered in respect to the explanation which has been attempted of the latter, is too obvious to be dwelt upon. The decreased gaseous pressure in the hot season is occasioned by the rarefaction of the air over the land whilst the sun is in the northern signs, and its consequent overflow in the higher regions, producing a return current in the lower strata; and the increased pressure in the cold season is occasioned by the cooling and condensation of the air, whilst the sun is on the south side of the equinoctial, and its consequent reception of the overflow in the upper strata from the regions which are then more powerfully warmed, and which is but partially counteracted by the opposite current in the lower strata.

In concluding this communication, I beg respectfully to submit to the consideration of the eminent meteorologists here present, that it is very important towards the progress of this science, that the propriety (in such discussions as the present) of separating the effect of the two elastic forces which are considered to unite in forming the barometric pressure, should be speedily admitted or disproved. The very remarkable fact recently brought to our notice by Sir James Ross, as one of the results of his memorable voyage, that the mean height of the barometer is full an inch less in the latitude of 75° S. than in the tropics, and that it diminishes progressively from the tropics to the high latitudes, presses the consideration of this point upon our notice; for it is either explained wholly or in greater part by the diminution of the vapour constituent in the higher latitudes, which diminution appears nearly to correspond throughout to the decrease of barometric pressure observed by Sir James Ross; or it is a fact unexplained, and I believe hitherto unattempted to be explained, on any other hypothesis, and of so startling a character as to call for immediate attention.

If, by deducting the tension of the vapour from the barometric pressure, we do indeed obtain a true measure of the pressure of the gaseous portion of the atmosphere, the variations of the mean annual gaseous pressure, which will thus be obtained in different parts of the globe,—and the differences of pressure in different seasons at individual stations,—may be expected to throw a much clearer light than we have hitherto possessed on those great aërial currents, which owe their origin to variations of temperature proceeding partly from the different angles of inclination at which the sun's rays are received, and partly from the nature and configuration of the material bodies at the surface of the earth: and a field of research appears to be thus opened by which our knowledge of both the persistent and the periodical disturbances of the equilibrium of the atmosphere may be greatly extended.

IX. *On some New Species of Animal Concretions.*

By THOMAS TAYLOR, Surgeon.

*To Richard Taylor, Esq.*

DEAR SIR,

AS the Catalogue of the Calculi belonging to the Royal College of Surgeons has now been published some months, and there consequently remains no further necessity for silence, I purpose in the following paper to redeem the promise I formerly made, of describing some of the more remarkable of the concretions which have been discovered during the examination of that very large collection; and also to detail the experimental proofs on which the assertions as to their composition were founded in the short notice which you did me the favour of inserting in this Journal in May 1844.

I do this the more willingly, as it was considered advisable to omit the details of the analyses in the Catalogue. Moreover, the Catalogue having but a limited circulation, many of the new facts that have been elicited would not otherwise be generally known. I shall, however, confine myself in this paper to the notice only of such concretions as are entirely new, or whose composition has been either imperfectly or incorrectly described. For the historical account of the successive steps by which our present knowledge of these bodies has been obtained, and for the description of the more common species of calculi, I must refer to the Catalogue itself.

*Urinary Calculus from the Iguana, consisting of Urate of Potass.*

Small and unimportant quantities of urate of potass may occasionally be detected in human urinary calculi, but no instance of this salt constituting an entire calculus has hitherto been described. There are three specimens of this description in the College collection, which resemble each other in every respect save in size. Two of them were described in the M.S. Catalogue of Sir Hans Sloane's collection as "*Piedra de Yguana*," and there is little doubt but that they were taken from the urinary bladder of some of the large Iguanas or tree lizards of South America. The other concretion had no history, but had been described as "*a mixed calculus in which uric acid predominates.*" Although much larger, it was so similar in composition and general appearance to the others, that there does not appear any reason to doubt its having a similar origin. In their external characters these concretions resembled calculi composed of the mixed phosphates, being made up of

concentric layers of a dirty white colour with a shade of pink. They were of an ovoid figure, but all of them were remarkable for being flattened in a peculiar manner on one side.

When heated before the blowpipe they consumed like a uric acid calculus, but left a fusible salt which spread over the platina foil, and tinged the outer flame violet. When heated in a test-tube, carbonate and hydrocyanate of ammonia with a little empyreumatic oil and water were given off. The carbonaceous residue, when treated with water, gave a solution which had a strong alkaline reaction, effervesced with acids, and emitted a slight odour of prussic acid. Tartaric acid and chloride of platina produced in the solution precipitates indicative of the presence of potass.

Water digested upon the powdered calculus afforded a solution which deposited small scales of suburate of potass upon being evaporated; the liquor gave no precipitate with a salt of lime, consequently no soluble oxalate was present.

When digested with caustic potass, ammonia was freely evolved, and the whole dissolved with the exception of a little flocculent matter. The alkaline solution, when mixed with muriatic acid, gave a copious precipitate of uric acid. It was therefore evident that these concretions consisted chiefly of urate of potass mixed with urate of ammonia.

The relative proportion of their constituents was estimated in the following manner:—

19·10 grs., when submitted to a current of dry air at  $212^{\circ}$  Fahr., lost 0·32 water, = 1·67 per cent. The dried powder was digested in boiling acetic acid, which decomposed the urate of potass and left a residue which weighed 15·02 grs., = 78·64 per cent. This residue was found to consist of uric acid, containing a small trace of oxalate of lime.

The acetic solution being evaporated to dryness was boiled with proof spirit; the whole dissolved with the exception of some light yellow flocks of animal matter, which amounted to 0·52 gr., = 2·73 per cent.

The spirituous solution being evaporated to dryness, the earthy and alkaline acetates it had contained were decomposed by the addition of muriatic acid; the mixture was evaporated to dryness and heated red-hot in a platina crucible. The residue dissolved entirely in water, with the exception of 0·06 gr. of phosphate of lime. The aqueous solution was mixed with carbonate of ammonia, a precipitate fell, which when dried and ignited was equivalent to 0·36 gr. of pure lime. It contained however a trace of phosphate of lime.

The solution from which the above precipitates had been separated was evaporated to dryness and the residue heated

red-hot. It was redissolved in water, and mixed with alcohol and chloride of platina. Chloride of platinum and potassium was thrown down, which when washed with alcohol and carefully dried weighed 10·32 grs., = 1·99 PO.

The quantity of ammonia was estimated by boiling the powdered calculus in a solution of potass, transmitting the ammonia evolved through diluted muriatic acid, and precipitating it in the ordinary manner by chloride of platina. 11·40 grs. yielded 4·57 of chloride of platina and ammonia, = 3·10 per cent. I do not, however, place much confidence in this mode of estimating the ammonia. The result of the analysis of this calculus, calculated in 100 parts, is therefore as follows: it is compared with an analysis of the calculus which had no history, in which the quantity of potass is rather greater:—

Uric acid mixed with a trace of oxalate of lime	78·64	78·36
Potass . . . . .	10·42	13·19
Ammonia . . . . .	3·10	3·09
Lime . . . . .	1·89	1·49
Magnesia . . . . .	0·00	0·29
Phosphate of lime . . . . .	0·32	0·02
Animal matter . . . . .	2·73	0·43
Water . . . . .	1·67	1·80
Sulphate of soda with chloride of sodium . . . . .		traces
	98·77	98·67

By another analysis, in which the quantity of potass was alone estimated by calcining the calculus until nothing but carbonate of potass was left, and precipitating the dissolved salt by chloride of platina, 10·72 per cent. of potass was found in the first, and 13·07 in the second calculus.

The potass in these concretions is probably derived from the leaves and other vegetable matter on which the Iguana partly subsists, while the carnivorous or insectivorous habits of the reptile are indicated by the large quantity of uric acid they contain.

*Urinary Calculus from the Sturgeon, consisting of Diphosphate of Lime. Beluga stones.*

These calculi are found by the fishermen of the Caspian Sea and of the Volga in a species of Sturgeon (*Acipenser Huso*, Linn.). The statements of different authors as to the situation of the stone in the fish, are very conflicting, some describing it as occurring in the air-bladder, others in the head and stomach. In Schrober's *Memorabilia Russico-Asiatica*, as quoted by Klaproth, it is said to be most frequently found in a small pouch communicating with the pancreatic

duct; his description is however confused and anatomically incorrect. The subjoined extracts from the works of Pallas\* leave no doubt as to these concretions being taken from the

\* “Les pêcheurs rencontrent assez souvent dans les gros biélougas, la pierre dont j’ai parlé, qui est encore un problème. Ils la vendent à un prix assez modique, de deux à trois roubles. Tous les pêcheurs à qui j’en ai parlé, m’ont assuré qu’on la trouve dans le gros boyau, qui leur sert à se vider et à jeter leurs œufs. On rencontre quelquefois des pierres dans les gros esturgeons ordinaires; elles sont semblables à celles des biélougas. On en trouve aussi dans les gros barbeaux, mais elles sont d’une espèce différente. Les pierres de biélouga sont ovales, unies, et quelques-unes grumelées assez grossièrement; d’autres sont triangulaires et toutes plates. Cette variété, dans la forme et la place qu’elles occupent, prouve que c’est une vraie pierre, et non une arête. Elles ont toutes la couleur et la texture de l’arête. Lorsqu’on les brise, on trouve dans leur substance des rayons luisans spathiques qui tendent de la circonférence au centre; outre la texture écailleuse qu’on distingue à la première superficie, il se détache de l’intérieur de quelques-unes de ces pierres un noyau; il a la même substance que la pierre, mais une autre forme; il ne se trouve pas toujours au centre. J’en ai vu plusieurs qui pesoient jusqu’à trois onces; je les croyois plus pesantes d’après leur grosseur. On peut en raper avec la lame d’un couteau, mais avec peine. J’ai essayé d’en mettre dans des acides et je n’y ai aperçu aucune marque d’effervescence. En Russie, on se sert de cette pierre comme remède domestique, dans les accouchemens laborieux, pour les maladies de l’urètre et celles des enfans; il est très en vogue, et l’on a grand tort. On en fait prendre dans de l’eau à très-petite dose. On attribue les mêmes vertus, et nombre d’autres, à la pierre qu’on rencontre quelquefois dans la vessie urinaire des sangliers, qu’on appelle *Kabannoï Kamen*, pierre de sanglier; elle est beaucoup plus chère que celle du biélouga.”—*Voyages de Pallas*, tom. i. p. 683.

“On fend le cartilage du dos pour en retirer les nerfs; on les lave et étend sur des perches pour les faire sécher.

“C’est en partageant ce cartilage dans toute sa longueur que l’on trouve quelquefois dans les plus gros ichthyocolles cette pierre si vantée. On ne l’aperçoit que lorsque le couteau s’arrête au moment où il la touche. Cette pierre est renfermée dans la chair rouge glanduleuse, qui est adhérente à la partie postérieure de l’épine du dos, et elle tient lieu de rognons. Elle est dans une petite peau particulière, qui remplit l’intérieur de cette espèce de glande. Je rapporte ici ce que M. Sokolof a pu apprendre de plus certain sur sa vraie position, des pêcheurs les plus instruits, qui assurent en avoir trouvé quelques-unes. A l’extérieur, elle est un peu molle et humide lorsqu’elle est fraîchement tirée, mais elle durcit aussitôt qu’elle est à l’air. C’est dans les pêches qui se font près d’Astrakan qu’on la rencontre le plus souvent. Elle n’est jamais plus grosse qu’un œuf de poule. Elle est ovale et assez plate un peu concave; où elle a l’angle qui adhère au cartilage un peu courbé.”—*Voyages de Pallas*, 1789, vol. ii. p. 486.

“In visceribus uropœis Husenum maximorum et ætate provectorum sæpius reperitur *Calculus* ovalis, depressus, hinc concavus, solidus, albus, intus *Zeolithi* fere instar a centro radiatus, nitidusque, cujus chemica analysis adhuc deest. Hunc plebs Rossica, et honoratiores etiam, pro magno medicamento uragogo et partum promovente æstimant atque infantibus propinquant, unde a piscatoribus pretio non exiguo redimuntur, *Calculi Husonis* (*Bjelushie Kamen*) nomine.”—*Zoographia Rosso-Asiatica*, vol. iii. p. 87.

dilated ureter or from the common cloacal termination of the gut of the fish.

These concretions have generally a flattened oval figure, their centre being often depressed or slightly concave. They vary considerably in size, but are usually about that of a hen's egg. Their surface is unequal but quite smooth, and of a yellowish-white colour. When broken they present a highly crystalline structure, consisting of fine plates or needles radiating from the centre to the circumference, but which are made up of very thin concentric layers adhering firmly together. Fragments of these calculi are translucent, and their interior is of a pure white colour. They are exceedingly scarce, and are highly esteemed for their supposed medicinal virtues. Dr. Cook informs us that the powder is highly commended as a diuretic and lithontriptic, and that the common people in the neighbourhood of the Volga take from ten to sixty grains, scraped fine in a little water, three or four times a day when the case is dangerous.

The composition of these calculi was first determined by Klapproth, but the earliest description of them is to be found in the Philosophical Transactions for 1748.

The specimen analysed by Klapproth had been received from Prof. Pallas. It weighed above seven ounces troy, and consisted of albumen 1, water 24, phosphate of lime 71.50, sulphate of lime 0.50.

17.13 grs. of one of the specimens in this collection, previously calcined, gave by solution in dilute muriatic acid and precipitation by oxalate of ammonia, 13.87 grs. of carbonate of lime, which is = 17.54 of the diphosphate of lime; 100 grs. of the same calculus gave—

		By calculation.
Water . . . . .	26.33	25.60 = 5 atoms.
Organic matter . . .	0.40	1.13
Diphosphate of lime .	73.27	73.27 = 1 atom.
	<u>100.00</u>	<u>100.00</u>

The Beluga stones therefore consist of an atom of diphosphate of lime combined with 5 atoms of water. The water is necessarily over-estimated in the analysis, on account of the organic matter being partially soluble in the diluted acid.

By another and more rigid analysis, I found the calculus to consist of 32.21 CaO, 40.33 PO<sub>5</sub>, and 26.33 HO, which would give 72.54 per cent. of diphosphate of lime. This calculus has also been analysed by Prof. Wöhler, who ascertained that four of the five atoms of water are driven off at 392° Fahr., while the last atom is expelled by a red heat.

The phosphoric acid is therefore in the tribasic state, and the formula of the salt will be  $\text{PO}_5 \cdot 2(\text{CaO}), \text{HO}, + 4 \text{aq}$ . I think it right to state that the analysis of this concretion had been printed some months previous to the publication of Prof. Wöhler's paper, and its analysis was made in 1843.

*Intestinal Concretions.*

The composition of the different kinds of intestinal concretions has been very little studied by chemists; a circumstance the more remarkable, as they present many points of interest both to the chemist and physiologist. The only description of these bodies to be found in the systematic works on chemistry, is almost wholly derived from the paper of Messrs. Fourcroy and Vauquelin, published in the 4th volume of the *Annales du Muséum National*.

By these chemists intestinal concretions were divided into the following species:—1, Calculi consisting of superphosphate of lime; 2, of phosphate of magnesia; 3, of phosphate of magnesia and ammonia; 4, of a biliary matter analogous to the colouring matter of the bile; 5, resinous concretions; 6, fungous concretions; and lastly, hair-balls. Their description of these bodies is however exceedingly slight and imperfect, and much inferior in accuracy to their previous researches on urinary calculi. In no instance did they determine the relative proportion of the constituents of the earthy concretions, and under the head of resinous concretions two essentially distinct species were included.

In the Catalogue I have endeavoured to supply these deficiencies by submitting most of these calculi to quantitative analysis, and by the addition of some new species. The following list includes all the intestinal concretions with which I am at present acquainted:—1, Calculi consisting of animal hairs; 2, of vegetable hairs; 3, of ellagic acid—the *oriental bezoar*; 4, of resino-bezoardic acid—the *occidental bezoar*; 5, of phosphate of magnesia and ammonia; 6, of diphosphate of magnesia; 7, of diphosphate of lime; 8, of oxalate of lime; 9, of ambergris.

*The Ellagic Acid Calculus.—The Oriental Bezoar.*

The composition of this species of calculus was described in a report to the Museum Committee in 1841, and in May 1843 I was permitted to insert a short notice as to its composition in the London and Edinburgh Philosophical Magazine. Since that period the ellagic acid calculus has been examined by MM. Merklein and Wöhler, who have confirmed my state-

ments as to its nature\*. The result of a careful comparison of the chemical characters of these concretions with those of ellagic acid prepared from the gall-nut† so fully established their identity, that I did not think it necessary to corroborate my statement by an ultimate analysis, especially as from the limited quantities of the calculus on which I could operate, and the facility with which ellagic acid becomes oxidized when dissolved in an alkali, I did not feel certain that I could ensure that perfect purity without which an organic analysis is wholly valueless. Its ultimate analysis has however been made by MM. Merklein and Wöhler, who have found it to agree with the analysis of ellagic acid by M. Pelouze, *minus* one atom of hydrogen.

The following description of the ellagic acid calculus, its properties and history, I shall quote verbatim from the College Catalogue published in July 1845, as it conveys in a condensed form the results of a long and troublesome series of experiments.

“Ellagic acid calculi are generally of an ovoid figure; their outer surface is smooth, polished, and of a deep olive or greenish brown colour; internally they are brown; they are made up of thin concentric layers, which in some cases adhere so slightly together, as to cause the calculus to fall to pieces on attempting to divide it with a saw. When any of the outer layers of these calculi are removed, the exposed surface readily acquires a high polish by slight friction, and when cut or scraped they assume a waxy lustre. These calculi invariably contain some foreign body as their nucleus, which is generally a small twig or seed.

\* *Ann. der Chemie und Pharm.*, August 1845. If by the following passage, “Aus dieser Zusammensetzung und den oben angegebenen Eigenschaften der Bezoarsäure folgt ferner der merkwürdige Umstand, dass diese Substanz, wie bereits von Th. Taylor vermuthet wurde, in der That nichts Anderes ist als Ellagsäure oder die Säure, die zuerst von Chevreul aus den Galläpfeln dargestellt und von Braconnot näher untersucht worden ist. Um nicht den geringsten Zweifel hierüber zu lassen, haben wir selbst Ellagsäure aus Galläpfeln dargestellt und ihre Eigenschaften mit denen der Bezoarsäure verglichen; sie zeigten sich vollkommen identisch,” MM. Merklein and Wöhler intend to imply that some doubt existed in my mind as to the composition of this concretion, I beg to state that such was not the case; and it is difficult to conceive on what grounds they could form such an opinion, as in the notice alluded to I simply stated the fact in the most concise and positive terms that could be made use of.

† For the opportunity of doing this on rather a large scale, I am indebted to my friend Mr. T. Morson, who kindly placed at my disposal a large quantity of the residue left in the preparation of gallic acid, and it gives me much pleasure to have an opportunity of acknowledging this and similar favours.



“The chemical characters of the constituent of these calculi agree so exactly with those of ellagic acid procured from the infusion of gall-nuts, as to leave no doubt of their being composed principally of that substance. When heated they do not fuse, but emit a slight balsamic odour and partially sublime; if more highly heated they catch fire, burn with a low flame, give off the smell of burning wood, and leave behind a carbonaceous ash. If the powder of the calculus be heated in a glass tube a yellow sublimate is produced, which condenses in the form of long spear-shaped crystals of a yellow colour, with a shade of green. These crystals do not differ in their chemical habitudes from the powder of the calculus, and they are identical in shape and appearance with those procured from the ellagic acid of the gall-nut when similarly treated. When the calculus is reduced to powder and diffused through water, several days elapse before the whole of the powder is deposited, and the water remains opalescent even for weeks. It is also difficult to separate the powder by filtration, as the liquid passes turbid for some time.

“Ellagic acid calculi easily dissolve, with the exception of a few flocks, in a cold solution of caustic potass or soda. The solution is of a deep brownish red colour, with a shade of green; when the ellagic acid is, however, freed from some extractive or colouring matter with which it is generally mixed in the calculus, the solution is of so deep a yellow as to appear red when viewed in bulk. Muriatic acid throws down from the potass solution a greenish, buff-coloured powder, while the supernatant liquor is of a light red colour. If the precipitate be examined by the microscope, it is seen to consist of small thread-like particles, generally blunt, but sometimes tapering at their extremities, and which are occasionally twisted or curved, especially if the solution from which they were thrown down was hot: they are not transparent, and can scarcely be termed crystals.

“When the potass solution is exposed to the air, oxygen and carbonic acid are absorbed, the solution becomes much darker coloured, and a silky greenish yellow precipitate is deposited, consisting of ellagate of potass. This precipitate appears under the microscope as thin rectangular plates, frequently arranged in stellate groups. If a current of carbonic acid is passed through the solution, a buff-coloured precipitate of ellagate of potass is thrown down, while the supernatant liquid remains of a dark reddish colour.

“Ellagic acid calculi are very sparingly soluble in solution of ammonia; the liquid acquires a yellow colour, which on ex-

posure to the air becomes brown and turbid. The small quantity of ellagic acid dissolved is precipitated by an acid.

“Concentrated sulphuric acid readily dissolves these calculi when assisted by a gentle heat. The solution is of a greenish brown colour, and is precipitated by dilution with water. The precipitate has the form of minute prisms arranged in stellate groups; the extremities of some of the prisms are blunt, others are pointed.

“When mixed with nitric acid, the ellagic acid calculus dissolves. If the acid be strong or slightly warmed, effervescence takes place, nitrous fumes are given off, and a solution is produced of a beautiful pink-red colour, similar to that produced by the action of nitric acid upon uric acid. The red colour quickly disappears upon standing; on being heated, a deep yellow solution remains, from which crystals of oxalic acid may be obtained by evaporation. Ammonia added to the solution causes it to assume a red colour, but does not render it turbid.

“The ellagic acid is best obtained from these calculi by dissolving the powdered calculus in a weak solution of caustic potass, and transmitting through it a current of carbonic acid. The precipitate which falls is to be digested in diluted muriatic acid, by which the potass is removed, and tolerably pure ellagic acid remains. During the whole of the operation great care must be taken to prevent the contact of atmospheric air; for when dissolved in alkaline liquids, ellagic is quickly converted into a species of ulmic acid. It is not improbable that catechuic acid is sometimes present in these calculi.

“This species of intestinal concretion appears to have been first examined by Fourcroy and Vauquelin, and is included in their class of *resinous Bezoars*\*. It was shortly afterwards examined by Berthollet, and subsequently by other chemists, all of whom failed in deciding upon its true nature; even so recently as 1843 this calculus was described by M. Lippowitz as consisting of a peculiar organic acid, for which he proposed the name of *Bezoaric acid*†.

“The concretions analysed by Berthollet, and of the properties of which he has given a very accurate account, had been

\* “La seconde variété d’une couleur brune ou violacée, sans saveur amère, presque insoluble dans l’alcool, entièrement soluble dans les alcalis, donnant dans cette dernière dissolution une liqueur qui devient rouge purpurine, lorsqu’elle s’épaissit et se sèche à l’air: fournissant à la distillation un sublimé concret, jaune, d’une saveur et d’une couleur de suie, insoluble dans l’eau et dans l’alcool.”—*Annales du Muséum National*, tom. iv. 334.

† Simon’s *Beitrag zur Phys. und Pathol. Chemie und Mikroskopie*, B. i. 463.

presented to the Emperor Napoleon by the Shah of Persia. They were of a greenish brown colour externally, and brown within; they had an oval figure, and their surface was highly polished; they were formed of irregular concentric layers, and in the centre of all of them was some vegetable matter; their sp. gr. = 1.463. They were regarded by Berthollet as consisting of the woody fibre (*lignin*) of the food of the animal, and he conjectures that they must have been taken from the stomach, on account of the little alteration which the vegetable matters that formed their nucleus\* had undergone.

“The constituent of the ellagic acid calculus is likewise described by John under the name of *Bezoarstoff*†; and Leopold Gmelin thinks it probable that the calculi examined by John were identical in composition with those analysed by Berthollet‡, and that they consisted of a species of ulmin arising from the decomposition of woody fibre or lignin.

“From the descriptions which Tavernier, Kæmpfer, and other Oriental travellers have given of the Oriental Bezoar, corroborated by the analyses of Fourcroy and Berthollet, there is no doubt that it is identical with the ellagic acid concretion above described. The signs by which a true Oriental Bezoar might be distinguished were, according to Tavernier, by steeping it in hot water, and observing whether the liquid became coloured, or the stone lost in weight. If either of these occurred, the stone was to be regarded as fictitious: but the best test was to apply a red-hot iron wire to the calculus, when, if it melted and permitted the iron to enter, it was certainly fictitious. Another test consisted in smearing a piece of paper with chalk, and rubbing the calculus over it. The genuine stone always left a greenish mark. All these criteria would be fulfilled by the ellagic acid calculus, but by none of the other species§.

“This species of concretion was the most valued of the Bezoars, and is denominated by Kæmpfer the ‘*verus et pretiosus Pasahr*,’ from which word, by a corruption of sound, he believes the word Bezoar to have been derived.

“With regard to the origin of this concretion, we have the fullest and most satisfactory evidence. W. Methold, Fryer, Tavernier and Kæmpfer all agree that it is taken most frequently from the alimentary canal of a species of wild goat termed *Pasen* by the Persians, which inhabits the mountainous ridges in Persia, particularly in the province of Chorasaaan or

\* *Mémoires de la Société d'Arcueil*, tom. ii. p. 448.

† *Chem. Schr.* iii. 38.

‡ *Handbuch der Chemic*, B. ii. S. 828, 1488.

§ In the Sloanian MS. Catalogue all the ellagic acid calculi were termed East Indian or Oriental Bezoars.

Chorasmia. Tavernier states that they likewise come from a province of the kingdom of Golconda. The account as to the exact situation of the stone is however not so clear. Most writers indicate the maw or stomach: Kæmpfer says it is found in the pylorus, 'sive productior quarti, quem vocant ventriculi fundus\*,' and that the natives are in the habit of ascertaining how many stones are contained in the stomach by feeling through the parietes of the abdomen, the value of the animal being considerably enhanced by their presence. When recently taken from the animal, they are said to be somewhat soft, or of the consistence of a hard-boiled egg, and that in order to preserve them it was customary to place them in the mouth, and retain them there until they acquired greater hardness.

"The Oriental Bezoar was not however confined to the wild goats, or to the ruminant tribes, as the *Pedra Bugia* or Ape stone also consists of ellagic acid. These concretions were held in higher esteem than those from the Goat, and were generally included, for the sake of preserving them, in a small cavity scooped out of two portions of a very light wood, which were held together by hoops wove from the twigs of the Rotang cane. There is in the Museum a specimen preserved in this manner. Kæmpfer informs us that they were found in a species of ape termed Antar by the Mongols, which he believes to be the *Babianum cynocephalum*. The composition of these concretions renders their origin no longer a matter of uncertainty, and confirms, in a very remarkable manner, the statements of Tavernier and Kæmpfer, that they are derived from the juices of the plants on which the animals fed."

MM. Merklein and Wöhler have proposed that the word ellagic should be changed for that of bezoaric acid, partly because the German word "Gall," when reversed, is not capable of being converted into ellagic, and partly from its want of euphony. If we were however, for the sake of euphony, to reject all the inharmonious appellations which the industry of modern chemists has introduced into the science, we might alter half the names at present in use; besides, as the ellagic acid was first procured from the gall-nut by Chevreul and subsequently named by Braconnot, I think it a matter of courtesy to adhere to its French derivation. The term bezoaric is also peculiarly improper, inasmuch as it would imply that the entire class of bezoars consisted of this peculiar acid.

I remain, dear Sir,

Yours most truly,

New Bridge Street, Nov. 20, 1845.

THOMAS TAYLOR.

\* *Op. cit.* p. 400.

X. *Inquiries in the Elements of Phonetics\**.By C. B. CAYLEY, *Fellow of Trinity College, Cambridge.**To the Editors of the Philosophical Magazine and Journal.*

GENTLEMEN,

**S**CHHEME of Consonants.—*Liquid Aspirates*.—I start with that scheme which has the authority of Messrs. Latham and Whewell (waiving some undecided points), to which I would make a few additions. I denote, for convenience, the aspirates (so called perhaps fitly, as the breathing is more heard in them) universally by adding to the corresponding non-aspirates a small *h*, which thus becomes a symbol from a letter, and I make *J*, suitably to its form and origin, stand for German *J*, English *Y*.

Sharp.	Flat.	Liquid.
P Ph	B Bh	M . . W
T Th	D Dh	N
K Kh	G Gh	L . . J
S Sh	Z Zh	R . . (place uncertain)

I observe that the flat consonants appear formed from the sharp by an effort (to make their sound stronger and more continuous), which also, being repeated, converts the flat into the liquid. This effort has not prevented, in the flat order, the introduction of the second force of aspiration as in the sharp: hence there is some reason to conjecture that the liquids might be susceptible of this force. In short, the symmetry of the system requires an order of liquid aspirates, which might be written *Mh*, *Nh*, *Lh*, *Rh*. May we not identify *Mh* with French *m* or *n* nasal, *Nh* with English *Ng*, *Lh* with Welsh *Ll*, *Rh* with Arabic Ghain?

Concerning the latter two, my chief difficulty is to know whether they are simple or compound sounds; but

The Welsh *Ll* is stated in Davis's Grammar to be the aspirate of *L*, pronounced by pressing the tongue against the teeth on both sides with a forcible emission of breath.

The Arabic Ghain is described by De Sacy as a sound resembling *R* and *G*, "comme l'*R* graissayé des Provençaux." We often hear imperfect pronunciations of *R* which might give the idea of this secondary liquid.

But I shall chiefly insist on *Mh* or *m* nasal, because it coincides with *F* and *V*, the aspirates as *M* with *B*, *N* with *D*, &c. Compare the pronunciation of *combattre*, *entendre*, *enfant*, *envahir*; likewise it has the same resemblance to the dental *N*, which *Ph*, *Bh* being formed by the teeth in part have to *Th* (whence the confusion made by children in those sounds, and

\* From letters addressed to Mr. Latham.

the Russian corruptions Feodor, Marpha, &c. for Theodore, Martha), so that all the labial aspirates would approximate to dentals, and on this principle might it be that the liquid aspirate *Nh* approaches the next order of gutturals.

Should this paper be admitted, I shall hereafter consider some objections that have been communicated to me, and proceed to an attempted analysis of vowel sounds.

C. B. CAYLEY.

XI. On Fresnel's Theory of Double Refraction.

By ARCHIBALD SMITH.

To the Editors of the *Philosophical Magazine and Journal*.

GENTLEMEN,

IN an article on Fresnel's Theory of Double Refraction, in the Supplement to the December Number of the *Philosophical Magazine*, Mr. Moon has quoted part of an article of mine in the first volume of the *Cambridge Mathematical Journal*, in which the following passage occurs:—"Let the particle receive a small displacement, the projections of which on the co-ordinate axes are  $\delta x$ ,  $\delta y$ ,  $\delta z$ . Then supposing the displacement to be very small, *the force of restitution may be taken as proportional to it, so that we have,*" &c. I am not surprised that Mr. Moon should remark on this passage, "what is meant by the mysterious principle 'supposing the displacement to be very small, the force of restitution may be taken as proportional to it,' I profess myself unable to understand."

The clause in italics, which was added to my manuscript when it was sent to the press, to remove, I believe, what was thought an abruptness in the reasoning, is certainly incorrect when applied to a doubly refracting medium. What was intended to be expressed, no doubt, was, that in the case supposed terms involving powers of  $\delta x$ ,  $\delta y$ ,  $\delta z$  higher than the first might be neglected. But this expression is only equivalent to the other in the case of a singly refracting medium. I may mention that in the middle of page 7 of the article in the *Cambridge Mathematical Journal*, the word "rays" was, by a mistake, substituted for "waves." These mistakes are corrected in the second edition of the first volume of the *Cambridge Mathematical Journal* which is now printing. As they have been noticed in your *Journal*, I shall feel much obliged by your inserting this explanation when you can afford space for it.

Your obedient Servant,

25 Old Square, Lincoln's Inn, Dec. 20.

ARCHIBALD SMITH.

XII. *On the Origin of the constituent and adventitious Minerals of Trap and the allied Rocks.* By JAMES D. DANA\*.

THE minerals of trap and the allied rocks may be arranged in two groups:—

1. Those essential to the constitution of the rock, or intimately disseminated through its texture.

2. Those which constitute nodules or occupy seams or cavities in these rocks.

Of the first group, are the several felspars, with augite, hornblende, epidote, chrysolite, leucite, specular, magnetic and titanite iron; and occasionally Hauyne, sodalite, sphene, mica, quartz, garnet and pyrites. Of the second group are quartz, either crystallized or chalcedonic, the zeolites or hydrous silicates, Heulandite, Laumonite, stilbite, epistilbite, natrolite, scolecite, mesole, Thomsonite, Phillipsite, Brewsterite, harmotome, analcime, chabazite, dysclasite, pectolite, apophyllite, prehnite, datholite, together with spathic iron, calc-spar and chlorite. Native copper and native silver might be added to both groups, yet they belong more properly to the latter. To the same also might be added sulphur, and the various salts that are known to proceed from decompositions about active volcanoes, including the crystallizations of alum, gypsum, strontian, &c.; but these more properly form still a *third* group, and being well understood, will not come under consideration in the remarks which follow.

We observe with regard to the minerals of the first group, that they are all anhydrous, that is, contain no water. In this respect, the essential constituents of trap and basalt are like those of granite and syenite. But in the second group, consisting of the minerals occurring in cavities or seams, all contain water except pectolite, quartz, calc-spar and spathic iron; and the last three are known to be always deposited in an *anhydrous* state from aqueous solutions.

We proceed to give a few brief hints with regard to the first group, intending only to glance at this branch of the subject, and then take up more at length the group of adventitious minerals.

*Essential constituents of modern Plutonic rocks.*—It is obvious that modern igneous rocks, although in some cases derived from the original material of the globe, have proceeded to a great extent from a simple fusion of rocks previously existing, and especially of the older igneous rocks. In accordance with this view, we may with reason infer that the tra-

\* Read before the Association of American Geologists and Naturalists, May, 1845, and communicated by the Author.

chytes and porphyries, which consist essentially of felspar, have proceeded, in many instances at least, from felspathic granites; the basalts and trap from syenites, hornblende or augitic rocks.

A theory proposed by Von Buch supposes that the felspathic rocks, as they are of less specific gravity, are from the earliest eruptions, or the more superficial fusings, while the heavier basalt has come from greater depths. Darwin thus accounts for the granites of the surface being intersected by basaltic dykes; the latter having originated from a deeper source, where their constituents took their place at some former period from their superior gravity. It virtually places hornblende rocks below felspathic granites in the interior structure of our globe. The hypothesis is ingenious and demands consideration; but it may not be time to give it our full confidence.

But supposing these more modern rocks to have been derived from the more ancient granitic—what has become of the quartz and mica which occur so abundantly in the latter, while they are so uncommon in the former? By what changes have they disappeared?

In the fusion produced by internal fires, the elements are free to move and enter into any combinations that may be favoured by their affinities. If silica, alumina, magnesia, lime, iron, the alkalies, potash and soda, were fused together—and these are the actual constituents of basalt—what result might we expect? From known facts, we should conclude that the silica would combine with the different bases, and these simple silicates would unite into more complex compounds. The silicates of alumina and the alkalies or lime, form thus one set of compounds, the felspars; the silicates of magnesia and the isomorphous bases, iron and lime, another set, to which belong augite, hornblende and chrysolite; and if much iron is present, we might have with the lime and alumina, the mineral epidote. The experiments of Berthier, Mitscherlich and Rose, and the facts observed amongst furnace slags, confirm what is here stated.

But not to go back to a resolution of the fused minerals into their elements, we may consider for a moment what changes the minerals themselves might more directly undergo in the process of fusion.

Much of the mica in granite differs from felspar in containing half the amount of silica in proportion to the bases, the bases in each being alumina and potash or soda. The change then in the conversion of the mica into felspar would require an addition of silica, which might be derived from the



free quartz of granite. Other varieties of mica contain magnesia, which would go towards the formation of some mineral of the magnesian series. It is possible that trachytes and porphyry have thus been made from granite; but trap rocks could not have been so derived, as they contain from 10 to 25 per cent. less of silica.

Again, hornblende and augite are so nearly related, that they have been considered by Rose the same mineral, the different circumstances attending the cooling giving rise to the few peculiarities presented. There can be no difficulty therefore in deriving augite by fusion from hornblende rocks. This moreover has been actually confirmed by experiment.

Augite, by giving up half of its silica, and receiving additional magnesia in place of its lime, is reduced to chrysolite\*. The Gehlenite, nepheline, anorthite and meionite of Vesuvius, contain, like scapolite, only 40 to 45 per cent. of silica and a large proportion of lime, and it is no improbable supposition, judging from the small amount of silica, and from the lime present, that scapolite rock, or rather limestones containing scapolite, may have contributed in part towards the lavas of that region. The ejections of unaltered granular limestones, and many mineral species pertaining to such beds, strongly support this view; and it is no less sustained by the fact, that in the Vesuvian basalts, Labradorite, which includes lime instead of the alkalies, replaces common felspar. The original felspar seems to have given way to leucite and Labradorite †.

An important source of new combinations is found in the sea-water which gains access to the fires of volcanoes. The decomposition which takes place eliminates muriatic acid, so often detected among volcanic vapours; but the soda and other fixed constituents remain, to enter into combination with some of the ingredients in fusion. Is not this one source of the soda forming the soda felspar, or albite, and of the muriatic acid and soda in sodalite? Phosphates have been long known to occur occasionally in volcanic rocks, and lately phosphoric acid has been proved to be generally common in small quantities. Sea-water is also a very probable source of this ingredient, as has been shown by late analyses of the same by Dr. Jackson.

\* The formula of augite is  $\bar{R}^3 \bar{Si}^2$ ; that of chrysolite,  $\bar{R}^3 \bar{Si}$ .

† Using  $\bar{R}$  for the bases and Si for silica, the formula of leucite is  $\bar{R} \bar{Si}^3$ ; that of common felspar,  $\bar{R} \bar{Si}^2$ ; that of Labradorite,  $\bar{R} \bar{Si}$ . From this, it appears that felspar may be reduced to leucite by giving up one-third of its silica, the bases being the same in the two; and with this excess and other silica combining with the lime at hand, Labradorite might be formed.

These few hints are barely sufficient to indicate something of the interest that attaches to this field of investigation, which the future developments of science will probably open fully to view. We do not attempt to explain why in these modern fusings, mica should not have remained mica, and the quartz still free uncombined quartz. The facts prove some peculiarity of condition attending the formation of the granitic rocks. Of this condition we know nothing certain, and can only suggest the common supposition of a higher heat and slower cooling, attending a greater pressure and different electrical conditions, and the same circumstances may have existed during the granites of different ages.

With these brief suggestions I pass to the second division of the subject before us.

2. *Minerals occupying cavities and seams in amygdaloidal trap or basalt.*—These minerals have been attributed to a variety of sources, and even at the present time there are various opinions respecting their origin. According to some writers, they result from the process of segregation; that is, a separation of part of the material containing rock during its cooling by the segregating powers of crystallization; and in illustration of the process we are pointed to the many segregations of felspar, quartz and mica, in granite and other rocks, the siliceous nodules in many sandstones, the pearlstones in trachytes and obsidian. Others have thought them foreign pebbles, enclosed at the time the rock was formed. Again, they are described as proceeding from the vapours which permeated the rock while still liquid, and which condensed as the rock cooled, in cavities produced by the vapours. By a few it is urged, admitting that the cavities are inflations by vapours like those of common lava, that they may have been filled either at the time the rock cooled or at some subsequent time, either by crystallization from vapours, or from infiltrating fluids, but more generally the latter.

Of these views we believe the last to accord best with the facts. Macculloch, in his *System of Geology*—a work which anticipated many of the geological principles that have since become popular—dwells at length on this subject, and supports the opinion here adopted with various facts and arguments. Lyell also admits the same principles. A review of the facts will enable us to judge of its correctness.

1. In the first place, the cavities occupied by the nodules are in every respect similar to the common inflations or air-bubbles in lava. These cavities are open and unoccupied in common lava, and may be no less frequently so in the ejections under water; and should they not be expected to fill in

some instances by infiltration? They are the very places where an infiltrating fluid would deposit its sediment, or collect and crystallize, if capable of crystallization; and such infiltrating fluids are known to permeate all rocks, even the most solid, and especially if beneath a body of water. It is evident, therefore, that we are supporting no strange or improbable hypothesis. On some volcanic shores one variety of the process may be seen in action. The cavities of a lava may be detected in the process of being filled with lime from the seawater washing over dead shells or coral sand, and at times a perfect amygdaloid is formed. But the positions and characters of the minerals themselves establish clearly the view we support.

2. The mineral in these cavities sometimes only fills their lower half, as if deposited from a solution; and again, it incrusts the upper half or roof, as if solidified on infiltrating through. In the large geodes of chalcedony, stalactites depend from above like those of lime from the roof of caverns, and, as Macculloch states, the stalactite is often found to correspond to an inferior stalagmite, the fluid silica having dripped to the bottom and there become solid; moreover, the superior pendent stalactite is sometimes found united with the stalagmite below. The same results are here observed as with lime stalactites in caverns, and often a similar laminated or banded structure, the result of deposition in successive layers. Such results can proceed only from a slow and quiet process,—a gradual infiltration of a solution from above into a ready-formed cavity; they can no more be supposed to arise from ascending vapours, or gaseous emanations from below, than the stalactite in the limestone cavern.

Another fact is often observed. A geode of quartz crystals, sometimes amethystine, in which every crystal is neatly and regularly formed, is found with the surface coated over with an incrustation of chalcedony, the part above hanging in small stalactites; and this chalcedonic coat sometimes scarcely adheres to the crystals it covers; or is even loose, and may be easily separated. There can scarcely be a doubt of a subsequent infiltration in a case of this nature.

We might rest our argument here, since the fact being ascertained with regard to quartz, it is necessarily established as a general principle with reference to the zeolites and other amygdaloidal minerals; for quartz or chalcedony, when present in these cavities, is, with rare exceptions, the *lower* or *outer mineral*. We find zeolites implanted on quartz, but very seldom quartz on zeolites. I have met with no instance of the latter, while the former is the usual mode of occurrence.

Any deduction, therefore, respecting quartz, holds equally for the associated minerals.

How a cavity coated with a deposit of chalcedony can still be afterwards filled up with other minerals, has been deemed a mystery in science, but the possibility of it is now not doubted. Even flint and agate, as Macculloch states, are known to give passage to oil and sulphuric acid; and much more will this take place in the moist rocks before the agate has been hardened by exposure to the air. Silica remains in a gelatinous state for a long period after deposition, and in this condition is readily permeable by solutions. It is not necessary that the fluid which has acted the part of a solvent and filled the cavity, should yield place to another portion of fluid; for the process of crystallization having commenced, a new portion of the material is constantly drawn in to the same fluid, and the necessary chemical changes are also promoted by the inductive influence of the changes in progress—the catalytic action as it is called—one of the most efficient, and at the same time one of the most universal agencies in nature.

Other evidence with reference to amygdaloidal minerals is presented by the zeolites themselves.

3. The zeolites occupy veins or seams as well as cavities. Often the seams were opened by the contraction of the cooling rock, and at other times they were of more recent origin. In either case the minerals filling these seams must be subsequent in formation to the origin of the rock itself, and could not have proceeded from vapours attending the eruption. These seams sometimes open upward and can be seen to have no connexion with the parts below, the rock in this portion being solid. Origin from above or from either side, is the only supposition in such cases.

Messrs. Jackson and Alger, in their valuable memoir on the geology of Nova Scotia, mention the occurrence of crystals of analcime attached to the extremity of a filament of copper, the copper having been the nucleus about which the solution crystallized, and state that their formation must have been subsequent to the formation of the rock.

4. Zeolites, moreover, have been found forming stalactites in basaltic caverns, as was observed by the writer in some of the Pacific islands: and Dr. Thomson has described and analysed one (Antrimolite) from Antrim in Ireland near the Giant's Causeway.

These facts favour throughout the view we urge, that the amygdaloidal minerals have in general resulted from infiltration, and were not necessarily formed simultaneously with the erupted rock.

5. We remark further, that no lavas have ever been shown to contain at the time of ejection any of the zeolitic minerals. The zeolites of Vesuvius are known to occur only in the older lavas, and afford no evidence against our position. The cavities in lavas, as far as observed, are empty as they come from the volcanic fires, with the exception of those containing sparingly some metallic ores which are condensed within them. Considering the fusibility of the zeolites and their easy destruction by heat and by volcanic gases, sulphureous and muriatic, we should *à priori* say that they could not be formed under such circumstances.

6. Besides, as we have stated, none of the proper constituents of trap or basalt—or the minerals disseminated through these rocks,—contain water. They are all anhydrous. The minerals formed accidentally in furnaces are anhydrous. The constituents of granite, syenite and porphyry, are all anhydrous. It is only those minerals which are found in geodes or seams that contain water. Of equal importance is the fact, that none of the essential constituents of these rocks have ever been found in these geodes or cavities along with the zeolites, as might have been the case had they been formed together, by segregation or otherwise. Neither felspar, although so abundant, nor augite, nor chrysolite, have been found filling, like zeolites, or with them, the cavities of amygdaloid. There is then a wide distinction between the anhydrous constituents of these rocks, and the hydrous zeolitic minerals.

A few zeolites have been found in granite or gneiss, but they are so disseminated that they can be shown to be of more modern origin than the rock, and to have resulted from some decompositions of true granite minerals. They differ entirely in their mode of distribution from the felspar, garnet, &c. of granite. Along with the decomposing felspar it is not unusual to find stilbite in the cavities formed by the decomposition.

Zeolites also have been found disseminated through the texture of basalt, clinkstone, &c., like the felspar, augite, &c. But the proportion varies widely, and in some parts of the same bed they are found to be wanting: so that we have sufficient reason for classing these disseminated zeolites with those in the cavities, as formed or introduced by infiltration.

7. Bearing upon this subject, it should be observed, that the constituents of amygdaloidal minerals are, in general, those of the containing rock. Silica, potash, soda, alumina, are found in the felspars; lime, magnesia and iron, in augite or hornblende; iron and magnesia in chrysolite. These are all

the constituents needed, except a little baryta for one species. The felspar decomposes readily and gives up its ingredients, its potash or soda, silica and alumina; the same is true of augite and chrysolite, which afford magnesia, lime, silica and iron. With water to infiltrate, we should therefore have all the necessary ingredients at hand for the required compounds. The fact already stated, that zeolites have been found as stalactites in caverns, seems to prove that they *do* actually result from decompositions and recompositions, such as have been supposed. Thus we have all the conditions at hand necessary for producing, by infiltration, the zeolites and the chlorite nodules of these rocks; the alumina, alkalies and lime, contribute, along with a portion of the silica, to the zeolites, and the magnesia, iron, and another portion of the silica, to the chlorite\*, often as abundant as the former. The amygdaloidal nodules frequently have a green coating, which further indicates the probable truth of these views; for it appears evidently to be a precipitate from the solution before a crystallization of the zeolites took place—a settling, perhaps, of the insoluble impurities taken up by the filtrating fluid in its passage through the rock, or of the formed chlorite, less soluble than the zeolites. Occasionally, when the rock contains copper, these nodules have an earthy coating of green carbonate of copper—the carbonate having proceeded, apparently, from the native copper of the rock, by the same process as explained.

The hypothesis of filtration seems, then, to be at least the principal source of these minerals. In some instances the filtrating fluid may have derived its ingredients from distant sources. The salts of sea-water may act an important part in these changes. Silica is dissolved on a grand scale during submarine eruptions, as we have elsewhere urged, and is thence distributed to the rocks around. Lime, also, is taken up in a similar manner. But the rock itself has often afforded the ingredients for the forming minerals, during the passage of the filtrating fluid through it. By the same means, the adjoining walls of a seam or dyke—which received the drainings from the rock of the dyke—are often penetrated by zeolitic minerals.

It may be thought that I am giving undue influence to a favourite theory, and in the minds of some, these conclusions may be set down among mere speculations in science. But

\* Chlorite consists of the same elements as augite or hornblende, except that the lime is excluded and water added. They are silica, alumina, magnesia, oxide of iron, with 12 per cent. of water.

the circumstances attending submarine igneous action, I am persuaded, are not generally apprehended. What is the condition of the deep bed of an ocean? Even at a depth of three miles, the waters press upon the bottom with a force equivalent to a million of pounds to the square foot; and with such a forcing power above, can we set limits to the depth to which these sea-waters—magnesia and soda solutions—will penetrate? Will not every cavern, every pore, far down, be filled under such an enormous pressure? Let a fissure open by an earthquake effort, and can we conceive of the tremendous violence with which the ocean will rush into the opened fissure? Let lava ascend, can we have an adequate idea of the effect of this conflict of fire and water? The rock rises, blown up with cavities like amygdaloid, and will a long interval elapse before every air-cell will be occupied from the incumbent water? Suppose an Hawaii to be situated beneath the waves, pouring forth its torrents of liquid rock;—this island contains about five thousand square miles, which is less than the probable extent of many a region of submarine eruption;—suppose, I say, the fires were opened and active over an area of some thousands of square miles—are there no effects to be discovered of this action? There is no geologist that pretends to deny the premises—the fact of such submarine eruptions, the ocean's pressure, the effect of fire in heating water, and in giving it increased solvent power; and why should they not reason upon the admitted facts, and study out the necessary consequences? Surely, if there have been effects, we might expect to see some of them manifested in the cavities of the ejected rocks, which were opened at the time to receive the waters and any depositions they might be fitted under the circumstances to make.

We are led by these considerations to another point in connexion with this subject,—the probable condition under which the different amygdaloidal minerals have been formed. Have they all proceeded from heated solutions, or all from cold solutions? or can we distinguish some which are indubitably of one or the other mode of formation?

Bearing on these questions, we notice such facts as are afforded by the condition and relative positions of the minerals in geodes. And I would here acknowledge my obligations to the valuable memoir, before alluded to, by Messrs. Jackson and Alger. The paucity of information on this subject to be found in the various accounts of similar rocks by other writers is surprising. Even where special pains have been taken to describe the mineral species, the relative positions of the minerals are very seldom noted. It has been altogether too com-

mon among geologists to treat mineral information with a degree of neglect almost amounting to contempt, although, as facts will probably hereafter show, they lie at the basis of an important branch of geological science.

But to proceed with the subject before us. We find that

*Quartz* or *chalcedony*, and *datholite*, very seldom overlie other mineral species in geodes or amygdaloidal cavities, while the latter often overlie them\*.

*Prehnite* is usually lowermost with reference to all the species except the two just mentioned. Occasionally it is found upon analcime, as at the Kilpatrick hills.

*Analcime* is commonly situated below all, except quartz, datholite and prehnite.

Of the remaining species, chabazite, stilbite, harmotome, Heulandite, scolecite, mesole, Laumonite and apophyllite, it is more difficult to distinguish an order of arrangement. My investigations only enable me to state that chabazite is usually covered by the rest (when associated with them), yet it is sometimes superimposed on stilbite; and apophyllite is almost uniformly above all with which it may be associated; calc-spar is at different times above and below. We thus arrive at the following as the usual order of superposition:—

1. Quartz.
2. Datholite.
3. Prehnite.
4. Analcime.
5. Chabazite, harmotome.

6. Stilbite, Heulandite, scolecite, natrolite, mesole, Laumonite, apophyllite.

It is a reasonable inference that the species which covers the bottom of a cavity was first deposited, and, as a general rule, that the others above were formed, either simultaneously, or in succession upon the lowermost, as their order may indicate. Each is usually perfect in its most delicate crystallizations, so that we cannot suppose that the minerals first deposited often underwent change after their deposition, though instances of this may no doubt be detected.

It is also evident, that if there were any species formed previous to the complete cooling of the rock, or if any require for their formation an elevated temperature, they are those first deposited—the first in the above series. A few considerations will place this, if possible, in a clearer light.

Quartz, as we have stated in a preceding page, and fully remarked upon elsewhere, enters largely into solution during

\* The writer has observed stilbite, apophyllite, calc-spar and prehnite overlying datholite, and various species over prehnite.



submarine eruptions. This solution has been shown, by actual experiment, to be a necessary consequence of such action. This fact corresponds most completely with the above deductions. Quartz usually forms the first lining of the geode or amygdaloidal cavity, when it is found at all, and, moreover, it is the most abundant of all amygdaloidal minerals.

Quartz may also proceed from decompositions of the rock in the cold, and incrustations of this kind are known to occur; but such an explanation does not account for its generally preceding all other species in filling cavities and seams in trap rocks, and is insufficient to produce the large deposits of silica, sometimes amounting to many tons in a single geode.

It should not be understood that the quartz is supposed to be derived always from the same heated waters that attended the formation of the containing rock; for later eruptions in the same region might, at a subsequent period, produce a like result: yet, as its place in the series proves it to be the earliest in formation, it has probably been generally deposited from the water heated during the eruption of the rock. Leaving quartz, we pass to the other minerals.

It is a striking fact that the minerals next to quartz in the table given—*datholite*, *prehnite* and *analcime*—contain less water than either of the following species. While the others include from 10 to 20 per cent., the first, *datholite*, has but 5 per cent., *prehnite* about  $4\frac{1}{4}$  per cent., and *analcime* 8 per cent.\* This fact certainly leans towards the view of their having originated at a somewhat more elevated temperature than the other species—the same conclusion that is drawn from their lower position in geodes.

The fact, also, that *prehnite* has been found forming pseudo-morphs, bears the same way; for heat would be necessary, in all probability, to aid in removing the original mineral. The vast extent of some *prehnite* veins—occasionally, as Dr. Jackson has observed, three or four feet wide—refers to an origin like that of the quartz in similar rocks. Indeed, there seems little doubt that *prehnite* is often derived from that

\* The following table shows the per-centage of water, and gives at the same time a general view of the composition of the zeolites:—

*Silica, boracic acid, lime.*—*Datholite* (5 Aq.).

*Silica, alumina, lime.*—*Prehnite* ( $4\frac{1}{4}$  Aq.). *Heulandite* (14 Aq.). *Scolecite* ( $13\frac{1}{2}$  Aq.). *Epistilbite* (14 Aq.). *Stilbite* (17 Aq.). *Laumonite* (17 Aq.).

*Silica, alumina, lime and potash, or soda.*—*Mesole* (12 Aq.). *Thomsonite* (13 Aq.). *Phillipsite* (17 Aq.). *Chabazite* (21 Aq.).

*Silica, alumina, and either soda, baryta or strontia.*—*Analcime* (8 Aq.). *Natrolite* ( $9\frac{1}{2}$  Aq.). *Harmotome* (15 Aq.). *Brewsterite* (13 Aq.).

*Silica, lime and potash.*—*Apophyllite* (16 Aq.).

*Silica, lime.*—*Dysclasite* ( $16\frac{1}{2}$  Aq.).

portion of the silica in solution which entered into combinations at the time with the alumina and lime which the siliceous waters contained; and probably the lime as well as silica was derived in part from an external source. The pseudomorphs prove that prehnite may have been the result also of subsequent eruptions, at the same time that they show the probable necessity of heat for its formation.

Datholite is a compound of silica, lime and boracic acid, with about 5 per cent. of water. Besides the small percentage of water, and its being, next to quartz, the lowermost mineral in geodes, we find an additional fact, alone almost decisive with regard to its origin, in its containing boracic acid. Boracic acid is often evolved about volcanoes or in volcanic regions. The hot lagoons of Tuscany, and the volcano of Lipari, are the most noted examples.

Although boracic acid has never been detected in sea-water, there can be little doubt of its occurring in it. The usual modes of analysis by evaporation would dissipate it, and of course it could not thus be detected except with special care and by operating on a large quantity of water. Borate of soda (boracite) is found only in beds of salt and gypsum,—both sea-water products. Moreover, borate of lime has been lately found on the dry plains in the northern part of Chili, along with common salt, iodine salts, gypsum and other marine salts; and all are so distributed over the arid country, that the region has been lately described as having been beyond doubt once the bed of the sea. These facts render it altogether probable that sea-water which gains access to volcanic fires is the source of the boracic acid in volcanic regions\*.

If this be its origin, the necessity of heat and pressure must be admitted, in order to produce the chemical combinations in datholite. Its elements are not those of the felspar or other trap minerals, like the zeolites superimposed on it; but they have come from an extraneous source, and none is more probable than the sea waters, which were heated at the submarine eruption, and permeated the bed of molten rock shortly after ejection. Thus placed in circumstances of pressure and confinement, along with silica in solution, the volatile boracic acid might enter into the combination presented in datholite.

An interesting fact bearing upon the history of datholite

\* The only other known source is the mineral tourmaline, quite an improbable one in the case before us. It is possible that tourmaline may have received its boracic acid from the sea during granitic eruptions, and the occurrence of this mineral in the vicinity of trap dykes is explained in the same manner.

was observed by Dr. Jackson at Keweena Point, Lake Superior. The datholite is often found there in veins with native copper, and is associated in some places with a curious slag of boro-silicate of iron and copper. Sometimes the crystals of datholite, as well as the prehnite and calc-spar, contain scales or filaments of native copper. These very important observations seem to establish the same origin for the three minerals, for Dr. Jackson states that they appear to be contemporaneous; and if calc-spar has been deposited from a solution, the same holds true of the others. They have all been formed subsequent to the copper filaments of the cavities, for they were deposited around them; yet may have been the next to form during the cooling of the rock. The boro-silicate of iron and copper has resulted from the same causes.

Analcime approaches the zeolites in composition, but like the prehnite and datholite it contains less water, and is very different in its crystallization. We have less evidence as to the heat necessary for its formation; yet it was probably formed at a somewhat elevated temperature.

With regard to the other amygdaloidal minerals, we are in still greater doubt as to the necessity of heat. We cannot at present fully appreciate the efficiency of chemical agents in a nascent state acting slowly without heat through long periods. Many of them may require heat, and some may be the last depositions from the filtrating waters after they have nearly or quite attained their reduced temperature. But the formation of zeolitic stalactites in caverns favours the view that some at least may form at the ordinary temperature by the slow decomposition of the containing rock after it had emerged from the waves\*. Kersten has lately described a modern stellated zeolite forming incrustations on the pump-wells of the Himmelsfahrt mine near Freyberg. It consisted of silica, oxides of iron and manganese and water. Further examination will probably bring more of these modern products to light †.

The formation of particular minerals in certain regions depends of course upon the supply of the necessary ingredients. Where the supply of lime has been large, we should expect to find some of the minerals, prehnite, Heulandite, Laumonite, stilbite, scolecite, dysclasite, chabazite, for carbonate of lime decomposes the silicates of potash and soda. Instances of

\* *Annales des Mines*, ii. (4th Ser.) 465, 1842.

† Carbonate of iron seems never to form from water at the surface, its solutions depositing a hydrated peroxide of iron instead of the carbonate; it may therefore require a submerged condition of the rock, although not necessarily a raised temperature.

this association of the lime zeolites with a large supply of lime in the vicinity are common. When there is little or no lime, or only the results proceeding from the decomposing rock, the other zeolites are formed—the hydrous silicates of alumina and potash or soda, occasionally with some lime. But if a salt of baryta or strontia is present, the decomposition of the silicates of the alkalies takes place as by the lime, and the mineral harmotome or Brewsterite is produced.

In the above explanations we have scarcely appealed to one source of amygdaloidal minerals admitted in the outset—their proceeding from vapours rising with the erupted rock; for it seems to be of but limited influence. Besides the arguments already brought forward, we state that the vapours which rise at the moment of eruption are insufficient. They inflate the rock or blow up the cavities; but the little vapour required to open the cavities most assuredly could not afford by condensation the mineral matter necessary to fill them,—to produce stalactites, stalagmite and successive layers of minerals. The vapours then, if the source, must have continued to rise for some time afterward. But is it possible that vapours should rise up through the solid rock? Such does not happen in the case of recent volcanoes; for fissures are first opened and then the vapours escape. And could it happen with the water above pressing down into the rock with the force of an ocean even a mile deep?

There may be instances of this mode of formation; but that it should be the usual mode is irreconcilable with the many facts stated. The form and condition of quartz or chalcedony in geodes, as well as the vast amount of this mineral in some cases,—the relative positions of the zeolites, and their occurrence as incrustations on rocks, or as fillings of cavities or seams, and never in disseminated crystals through the texture of the rock,—the green coating of the nodules, which is sometimes a carbonate of copper when there is native copper in the rock to undergo alteration,—the correspondence between the elements of the minerals and the composition of the including rock, and at the same time their contrast in being hydrous while the constituents of the latter are anhydrous,—and the known formation of zeolites in caverns,—these various facts appear to establish infiltration as the principal means by which amygdaloidal minerals have been produced.

XIII. *Reflections on the Resolution of Algebraic Equations of the Fifth Degree.* By G. B. JERRARD.

[Continued from vol. xxvi. p. 574.]

47. **T**HE remarks in No. 44. related to a difficulty which must arise if we can, as seems to have been proved, succeed in tracing the equation for  $W$  to a class of equations of the sixth degree, the solution of which can be effected. I have lately reconsidered the subject of that number, and the exact nature of the difficulty in question will, I think, appear from what follows.

Since every symmetric function of the quantities  $V_I, V_K, V_L$  will be such as to remain unchanged whilst one of the roots,  $x_1$ , continues fixed, and  $x_2, x_3, x_4, x_5$  are permuted in every possible way among themselves, it might easily be shown that the equation for  $V$ ,

$$V^{15} + C_1 V^{14} + \dots + C_{15} = 0,$$

will admit of being resolved into five factors of the form

$$V^3 + \frac{1}{5} C_1 V^2 + r_2(x_\alpha) V + r_3(x_\alpha) = 0,$$

obtained by writing 1, 2, 3, 4, 5 successively for  $\alpha$ :  $r_2$  and  $r_3$  being expressive of rational functions, and such that  $r_n(x_\alpha)$  shall essentially involve  $x_\alpha$ . In this equation, therefore, we cannot generally write  $r_n(0)$  instead of  $r_n(x_\alpha)$ .

But the equation for  $V$  will evidently lead to twenty-five expressions for the five roots  $x_1, x_2, \dots, x_5$ , obtainable from a system of functions of  $x_\alpha$ ;

$$\Psi_1(x_\alpha), \Psi_2(x_\alpha), \dots, \Psi_5(x_\alpha)^*.$$

The question therefore suggests itself: Is it permitted, since the number of distinct values of  $x$  cannot exceed 5, to suppose that

$$\Psi_m(x_\alpha) = \Psi_n(0),$$

or that the five roots,  $x_1, x_2, \dots, x_5$ , without considering in what order they will arise, may be expressed by

$$\Psi_1(0), \Psi_2(0), \dots, \Psi_5(0);$$

and thus to avoid the conclusion that the equation of the third degree, at which we shall arrive, will, in the ordinary meaning of the term, be simultaneous with  $V^{15} + C_1 V^{14} + \dots + C_{15} = 0$ ?

In fine, if we can effect the resolution of algebraic equations of the fifth degree, it must be possible to withdraw the terms involving  $x_\alpha$  from  $\Psi(x_\alpha)$  considered throughout its extent, although we retain those which involve  $x_\alpha$  in  $r(x_\alpha)$ .

London, December 13, 1845.

\* See (31.).

XIV. *Proceedings of Learned Societies.*

## ROYAL SOCIETY.

Nov. 27, "EXPERIMENTAL Researches in Electricity." By 1845. Michael Faraday Esq., D.C.L., F.R.S., &c. Nineteenth Series. Section 25: On the Magnetization of Light, and the Illumination of Magnetic Lines of Force.

For a long time past the author had felt a strong persuasion, derived from philosophical considerations, that among the several powers of nature which in their various forms of operation on matter produce different classes of effects, there exists an intimate relation; that they are connected by a common origin, have a reciprocal dependence on one another, and are capable, under certain conditions, of being converted the one into the other. Already have electricity and magnetism afforded evidence of this mutual convertibility; and in extending his views to a wider sphere, the author became convinced that these powers must have relations with light also. Until lately his endeavours to detect these relations were unsuccessful; but at length, on instituting a more searching interrogation of nature, he arrived at the discovery recorded in the present paper, namely, that a ray of light may be electrified and magnetized; and that lines of magnetic force may be rendered luminous.

The fundamental experiment revealing this new and important fact, which establishes a link of connexion between two great departments of nature, is the following. A ray of light issuing from an Argand lamp is first polarized in the horizontal plane by reflexion from a glass mirror, and then made to pass, for a certain space, through glass composed of silicated borate of lead, on its emergence from which it is viewed through a Nichol's eye-piece, capable of revolving on a horizontal axis, so as to intercept the ray, or allow it to be transmitted, alternately, in the different phases of its revolution. The glass through which the ray passes, and which the author terms the *dimagnetic*, is placed between the two poles of a powerful electro-magnet, arranged in such a position as that the line of magnetic forces resulting from their combined action shall coincide with, or differ but little from the course of the ray in its passage through the glass. It was then found that if the eye-piece had been so turned as to render the ray invisible to the observer looking through the eye-piece before the electric current had been established, it becomes visible whenever, by the completion of the circuit, the magnetic force is in operation; but instantly becomes again invisible on the cessation of that force by the interruption of the circuit. Further investigation showed that the magnetic action causes the plane of polarization of the polarized ray to rotate, for the ray is again rendered visible by turning the eye-piece to a certain extent; and that the direction of the rotation impressed upon the ray, when the magnetic influence is issuing from the south pole, and proceeding in the same direction as the polarized ray, is right-handed, or similar to that of the motion of the hands of a watch, as estimated by an observer at the eye-piece. The direction in which the rotation

takes place will, of course, be reversed by reversing either the course of the ray or the poles of the magnet. Hence it follows that the polarized ray is made to rotate in the same direction as the currents of positive electricity are circulating, both in the helices composing the electro-magnet, and also in the same direction as the hypothetical currents, which, according to Ampère's theory, circulate in the substance of a steel magnet. The rotatory action was found to be always directly proportional to the intensity of the magnetic force, but not to that of the electric current; and also to be proportional to the length of that portion of the ray which receives the influence. The interposition of substances which occasion no disturbance of the magnetic forces, produce no change in these effects. Magnets consisting only of electric helices act with less power than when armed with iron, and in which magnetic action is consequently more strongly developed.

The author pursues the inquiry by varying in a great number of ways the circumstances in which this newly-discovered influence is exerted; and finds that the modifications thus introduced in the results are all explicable by reference to the general law above stated. Thus the effect is produced, though in a less degree, when the polarized ray is subjected to the action of an ordinary magnet, instead of one that derives its power from a voltaic current; and it is also weaker when a single pole only is employed. It is, on the other hand, increased by the addition of a hollow cylinder of iron, placed within the helix, the polarized ray traversing its axis being then acted upon with great energy. Helices act with equal power in any part of the cylindric space which they enclose. The heavy glass used in these experiments was found to possess in itself, no specific magneto-inductive action.

Different media differ extremely in the degree in which they are capable of exerting the rotatory power over a polarized ray of light. It is a power which has no apparent relation to the other physical properties, whether chemical or mechanical, of these bodies. Yet, however it may differ in its degree, it is always the same in kind; the rotation it effects is invariably in one direction, dependent, however, on the directions of the ray and of the magnetic force. In this respect it differs essentially from the rotatory power naturally possessed by many bodies, such as quartz, sugar, oil of turpentine, &c., which exhibit the phenomena of circular polarization; for in some of these the rotation takes place to the right, and in others to the left. When, therefore, such substances are employed as dimagnetics, the natural and the superinduced powers tend to produce either the same or opposite rotations; and the resulting effects are modified according as they are cumulative in the former case, and differential in the latter.

In the concluding section of the paper, the author enters into general considerations on the nature of the newly-discovered influence of electricity and magnetism over light, and remarks that all these powers possess in common a duality of character which constitutes them a peculiar class, and affords an opening which before was

wanting for the appliance of these powers to the investigation of this and other radiant agencies. The phenomena thus brought to light confirm the views entertained by the author relative to the constitution of matter as being spheres of power, for the operation of which the conception of a solid nucleus is not necessary; and leads to the presumption that the influence of magnetism on bodies which exhibit no magnetic properties consists in producing in them a state of electric tension tending to a current; while on iron, nickel, and other bodies susceptible of magnetism, currents are actually established by the same influence.

The author states that he is still engaged in the prosecution of these inquiries.

“On the Action of the Rays of the Spectrum on Vegetable Juices:” being an Extract from a Letter by Mrs. M. Somerville to Sir John F. W. Herschel, Bart., dated Rome, September 20, 1845. Communicated by Sir John F. W. Herschel, Bart., F.R.S.

In the experiments of which the results are here recorded, the solar spectrum was condensed by a lens of flint glass of seven inches and a half focus, maintained in the same part of the screen by keeping a pin-hole or pencil-mark constantly at the corner of the red rays, which were sharply defined by being viewed through blue spectacles; and the apparatus was covered with black cloth in order to exclude extraneous light. Thick white letter-paper, moistened with the liquid to be examined, was exposed wet to the spectrum, as it was found that the action of the coloured light was thus rendered more immediate and more intense, than when the surface of the paper was dry.

The action of the spectrum at the junction of the lavender with the violet rays was found in some cases to be different from what it is with either of these colours separately, indicating a break in the continuity of action, and suggesting the idea of a secondary spectrum. In many instances the yellow and green rays exert a powerful influence on vegetable substances, an influence apparently unconnected with heat; for the darkening is generally least under the red rays and immediately below them, where the calorific rays are most abundant. The action, in a great number of cases, produces insulated spots in different parts of the spectrum, but more especially in the region of the rays of mean refrangibility, in which neither the calorific nor the chemical powers are the greatest. The point of maximum intensity is sometimes altered by the addition of acids, alkalies, or diluted alcohol. But altogether, as the author states, the action of the different parts of the spectrum seems to be very capricious, the changes of colour produced being exceedingly irregular and unaccountable.

## XV. *Intelligence and Miscellaneous Articles.*

### ACTION OF NITRIC ACID ON WAX.

**W**HEN wax is boiled in nitric acid, the same phænomena, according to M. Gerhardt, result as when the acid is made to act



upon stearic acid or other fatty bodies; much nitrous vapour is disengaged; but the action is not so vivid, as when olive oil, for example, is treated by the acid.

About 4300 grains of wax were boiled with rather less than two pints of common nitric acid for about two hours, and the mixture, allowed to cool, became a solid mass; this was perfectly dissolved by carbonate of soda, with the production of slight effervescence. On cooling the whole became one mass; the wax was unctuous and of an apricot colour. After twenty-four hours' ebullition, the greater part of the wax was dissolved in the nitric acid; an oily substance, having the smell of rancid butter, floated on the solution; this was entirely dissolved by potash: this oil was acid, and could not be distilled without decomposing, and possessed all the properties attributed by M. Laurent to azoleic or œnanthylic acid. The formation of this acid has been observed to occur, as is well known, during the oxidizement of stearic and oleic acids, and other fatty bodies.

Wax was afterwards boiled with twice its weight of nitric acid, during several days, until all the oily matter disappeared; the first crystalline grains which deposited by the cooling of the solution, were pimelic acid, as shown by analysis, which gave carbon 52, hydrogen 7·8, indicating as its composite, carbon 52·5, hydrogen 7·5, oxygen 40 in 100 parts, or  $C^7 H^6 O^4$ .

The mother-water yielded a considerable quantity of adipic acid, but which appeared to be mixed with lipic acid. The last portions of the mother-water yielded no crystals, but were rendered turbid by the addition of water, and deposited fresh portions of oily azoleic acid. Lastly, when the wax was treated with nitric acid, till red vapours ceased to be produced, fine crystals of succinic acid were obtained. The formation of this body has been already shown by Mr. Ronalds.—*Ann. de Ch. et de Phys.*, Oct. 1845.

#### DRY DISTILLATION OF WAX.

M. C. Gerhardt states, that when wax is submitted to dry distillation, there condenses in the receiver a solid, white granular matter, floating in an oily liquid, and during the whole time of the operation a mixture of carbonic acid and bicarburetted hydrogen gases is evolved. The condensed portions consist of a fatty acid, a solid carburetted hydrogen, and several liquid carburetted hydrogens; the products become more and more impure as the operation approaches its termination, and sometimes, when the last remains of the wax are carbonized, a small quantity of a reddish solid matter is obtained. If the products be separately received at different times, it is found that the fatty acid passes first, and afterwards the solid carburetted hydrogen; the liquid carburetted hydrogens are among the last products. When the distillation is rapidly performed, there remains little else than a coaly residue.

The first portions of the distillation saponify almost entirely, except a few particles of solid carburetted hydrogen. The soap yields, by the action of hydrochloric acid, a perfectly white fatty acid; when

crystallized once or twice from æther slightly alcoholized, this acid melts at 140° Fahr., and becomes a radiated mass on cooling; this acid, after being fused, yielded by analysis,—

Carbon .....	75·1
Hydrogen .....	12·8
Oxygen .....	12·1
	100·

which gives as the formula  $C^{17} H^{17} O^2$ , and the substance produced was therefore margaric acid. The solid carburetted hydrogen which accompanies the above substance is paraffin, as shown by the experiments of M. Etting.—*Ann. de Ch. et de Phys.*, Nov. 1845.

---

ANALYSIS OF PHOSPHATE OF ALUMINA. BY M. A. DELESSE.

M. Danhauser discovered at Bernay, near Epernay, a white substance, considerably resembling alumina dried on a filter; it invested a gangue coloured by the oxides of iron and manganese, and appeared to belong to the plastic clay formation. Several collections in Paris contain specimens of it, but that examined by M. Delesse contained phosphoric acid.

In the closed tube this substance blackens and yields much water, containing bituminous matter; it is acid, reddens litmus paper, and appears also to corrode glass slightly, which may indicate the presence of a little hydrofluoric acid. In the outer flame of the blow-pipe, the black colour produced by the carbon of the organic matter disappears and the substance becomes white; it is infusible. With the salt of phosphorus it readily dissolves, and a very transparent bead is formed; with carbonate of soda this substance swells, but does not dissolve; with nitrate of cobalt it yields a fine blue colour.

When not calcined this substance dissolves entirely and with the greatest facility in acids; it also dissolves, but with difficulty, in potash. After calcination, it is scarcely and with difficulty acted upon by acids.

It will be observed that the substance possesses all the properties of pure alumina, and, as already observed, it has the appearance of it; the presence of phosphoric acid was, however, ascertained by the process of Vauquelin and Thenard; it also contains a little lime, which is undoubtedly in the state of carbonate, for when acted upon by acids there is a disengagement of gas.

After several hours' drying, so as to expel the hygrometric moisture, the loss amounted to about 10 per cent; and by analysis the substance yielded,—

Phosphate of alumina .....	46
Water and organic matter.....	49
Carbonate of lime and loss ....	5
	100

M. Delesse states that he did not possess a sufficient quantity of the mineral to determine the quantity of phosphoric acid; but it is

evident that it must form a distinct species from wavellite, which contains 20 to 30 per cent. of water, while the phosphate of Bernon contains 49 per cent. and an organic substance. Vauquelin has also described, in the 21st vol. of the *Annales de Chimie et de Physique*, an hydrated phosphate of alumina, from the Isle of Bourbon, the composition of which is also different from that of wavellite, and likewise contains ammonia.—*Annales des Mines*, 1844.

#### A NEW PLANET.

The Astronomer Royal has forwarded to the *Times* newspaper the following letter from Prof. Encke of Berlin, relating the discovery of a new planet. Mr. Hind had previously communicated an extract of a letter from Prof. Schumacher, announcing the fact of Mr. Hencke's new planet, accompanied with a statement on the part of Mr. Hind, that he could not find any star answering the description of the supposed new one.

“Berlin, Dec. 15th.

“On the 13th of December, Mr. Hencke, of Driessen, gave notice that he had found a star of the ninth magnitude, in a place where before there was none. He gave its position by reference to the star-map of the Berlin Academy, 4th hour (which particular map was very carefully drawn by Prof. Knorr), from which its place appears to have been: Dec. 8.—At 8 hours; right ascension in arc,  $65^{\circ} 25'$ ; declination north,  $12^{\circ} 41'$ .

“Yesterday, Dec. 14, we sought for it with our refractor, and found, by comparison with the star-map of the Berlin Academy (which alone, on account of the fulness of its details, could have enabled us to discover it), a star of the ninth magnitude, not marked in the map, whose place was: Dec. 14.—At 6 hours 28 min. mean time, right ascension in arc,  $64^{\circ} 4' 53''\cdot 2$ . At 12 hours 43 min. mean time, right ascension in arc,  $64^{\circ} 1' 10''\cdot 3$ .

“We then determined the following places with the wire micrometer, each place being the mean of five observations. At 13 hours 34 min. 55·6 sec. mean time, right ascension in time, 4 hours 16 min. 2·44 sec.; declination north,  $12^{\circ} 39' 54''\cdot 2$ . At 13 hours 42 min. 36·5 sec., right ascension in time, 4 hours 16 min. 2·08 sec.; declination north,  $12^{\circ} 39' 53''\cdot 1$ . At 14 hours 33 min. 27·1 sec., right ascension in time, 4 hours 16 min. 0·2 sec.; declination north,  $12^{\circ} 39' 52''\cdot 1$ . Or, taking the mean, at 13 hours 56 min. 59·7 sec. mean time; right ascension in arc,  $64^{\circ} 0' 23''\cdot 6$ ; declination north,  $12^{\circ} 39' 53''\cdot 1$ .

“The motion is retrograde, and its daily amount, as determined from the observations, eight hours apart, is—in right ascension,  $14' 21''\cdot 2$  of arc; in declination it is quite insignificant.

“Mr. Hencke's place of December 8th agrees very nearly with this.

“The star is probably a new planet near its opposition. Vesta is pretty near it, and is also in opposition.

“On account of the difficulty of following it, I have thought it

best to send you the news directly; and I beg you to make it known in England, that a sufficient number of observations may soon be collected. Excuse the shortness of this letter, which is written in great haste. Yours, &c. "ENCKE."

Professor Airy says, there appears to be no reasonable doubt that the object to which the foregoing relates is a new planet.

Mr. Hind has since observed the new star: At 0 h. 20 min. 15 sec., sidereal time, on Wednesday evening, the right ascension of the new planet was 4 hs. 8 min. 17.58 sec., and the declination  $12^{\circ} 45' 32'' \cdot 6$ , north. He was enabled to establish its motion in R.A. from the observations made at Mr. Bishop's Observatory, Regent's Park, on that evening. The planet has the appearance of a star of the ninth or tenth magnitude.—*Literary Gazette*, Dec. 27.

The following letter from the Astronomer Royal has since been published in the *Times* newspaper of the 29th inst. :—

Royal Observatory, Greenwich, Dec. 27.

Sir,—I have this day received from Prof. Schumacher a letter relating to the new planet, of which I request you to publish the following extract.

I am, Sir, your obedient Servant,  
G. B. AIRY.

(Extract of a letter from Professor Schumacher.)

"Mr. Encke obtained an observation on the 20th of December, and this has enabled him to give an approximate sketch of the orbit of the new planet. I send you the elements :—

"Epoch of mean longitude, 1846, Jan. 0, at 0 hour,  $89^{\circ} 32' 12'' \cdot 1$ ; longitude of perihelion,  $214^{\circ} 53' 7'' \cdot 0$ ; longitude of ascending node,  $119^{\circ} 44' 37'' \cdot 5$ ; inclination,  $7^{\circ} 42' 8'' \cdot 4$ ; eccentricity, 0.207993; logarithm of semiaxis major, 0.42144; daily mean motion in longitude,  $827'' \cdot 65$ ; periodic time, 1565 days.

"The discoverer has left the determination of the name to Mr. Encke, and Mr. Encke calls it 'Astræa.'

"Yours, &c., H. C. SCHUMACHER.

"Altona, Dec. 23."

---

NOTICE OF AN AURORA BOREALIS SEEN AT MANCHESTER.

*To the Editors of the Philosophical Magazine and Journal.*

GENTLEMEN,

I will thank you to record in your *Philosophical Magazine*, &c. an aurora borealis which was seen at this place on Wednesday evening, the 3rd instant. It was first seen as a luminous arch about six o'clock; at half-past six the arch was complete throughout, from the eastern to the western horizon, with a span of upwards of  $100^{\circ}$ . The arch of light was perfectly steady and of an unusual breadth, much broader indeed than any I have before noticed. The altitude of the arch also was unusually great.  $\alpha$  *Ursæ Majoris*, then near the meridian beneath the pole, was within the lower margin of the

luminous band, whose highest point was about the meridian of *a Draconis*. From the upper edge of the western limb sprang several extensive streamers; but the light of the moon (then about four days old) prevented their being very brilliant. Shortly after seven, heavy clouds completely covered the aurora, showing their distance to be less than that of the luminous meteor which they obscured, and which, by their long continuance, they finally closed.

We had some very heavy hail showers during the day, and a perfect gale of wind the previous night. The same day a terrific thunderstorm visited a great part of Wales, killing several head of cattle and doing other serious damage to property.

I am, Gentlemen,

Your obedient Servant,

Manchester, December 19, 1845.

W. STURGEON.

METEOROLOGICAL OBSERVATIONS FOR NOV. 1845.

*Chiswick*.—November 1. Slight haze: very fine. 2. Slight fog: overcast. 3. Frosty: fine: clear and frosty. 4. Frosty, with dense fog: clear and frosty at night. 5. Frosty and foggy: very fine: overcast. 6. Very fine: rain. 7. Clear and fine: cloudy: rain. 8. Cloudy. 9. Very fine: slight rain. 10. Very fine: heavy clouds. 11. Hazy: rain. 12. Very fine. 13. Hazy: very fine. 14. Foggy throughout. 15. Foggy: fine. 16. Densely clouded: rain. 17. Fine: rain. 18. Cloudy: clear. 19. Boisterous, with rain: showery: very clear at night. 20. Fine. 21. Overcast: heavy rain. 22. Fine: clear and cold. 23. Sharp frost: fine. 24. Very fine: foggy at night. 25. Uniformly overcast: slight rain: foggy. 26. Densely overcast. 28. Cloudy. 29. Heavy rain. 30. Cloudless: overcast at night.—Mean temperature of the month  $1^{\circ}45$  above the average.

*Boston*.—Nov. 1. Fine. 2, 3. Cloudy. 4—7. Fine. 8. Cloudy: rain early A.M. 9. Fine. 10. Foggy. 11. Fine: rain P.M. 12. Cloudy. 13. Fine. 14, 15. Cloudy. 16. Cloudy: rain early A.M. 17. Cloudy: rain early A.M.: rain P.M. 18. Cloudy: rain early A.M. 19. Stormy: rain A.M. 20—23. Fine. 24. Fine: snow and rain early A.M. 25—28. Cloudy. 29. Cloudy: rain P.M. 30. Fine.

*Sandwich Manse, Orkney*.—Nov. 1. Bright: cloudy. 2. Fine: cloudy. 3. Fine: frost: cloudy. 4. Bright: clear. 5. Clear. 6. Damp: cloudy. 7. Damp: hazy. 8. Drizzle: cloudy. 9. Cloudy: damp. 10. Damp. 11. Cloudy: fog in valleys. 12. Frosty: fog: clear. 13. Fine. 14. Fine: frost: fine. 15. Fine: cloudy. 16. Fine: rain. 17. Fine: showers. 18. Cloudy. 19. Rain: cloudy. 20. Showers. 21. Showers: sleet. 22. Cloudy: showers. 23. Cloudy: snow-showers. 24. Cloudy: snow: rain. 25. Showers: rain. 26. Showers: thunder and showers. 27. Showers: hail: showers. 28. Cloudy: showers. 29. Cloudy: showers: sleet. 30. Sleet-showers: snow on hills.

*Applegarth Manse, Dumfries-shire*.—Nov. 1. Fair and fine. 2. Fair and chilly. 3. Fair, but dull: frost A.M. 4. Frost, hoar: clear and cold. 5. Frost: dull. 6. Fair and fine: fresh. 7—10. Rain early A.M. 11. Fair and fine. 12—13. Hoar-frost: fine. 14. Raw and cloudy. 15. Rain P.M. 16. Heavy rain P.M. 17. Fine: dry. 18, 19. Heavy showers. 20. Fine A.M.: rain P.M. 21. Showers. 22. Frost. 23. Frost: a few drops of rain. 24. Frost: cloudy P.M. 25. Wet. 26—28. Very heavy rain. 29. Showers. 30. Heavy rain P.M.

Mean temperature of the month .....  $42^{\circ}7$   
 Mean temperature of Nov. 1844 .....  $43^{\circ}6$   
 Mean temperature of Nov. for twenty-three years.  $40^{\circ}2$



THE  
LONDON, EDINBURGH AND DUBLIN  
**PHILOSOPHICAL MAGAZINE**  
AND  
**JOURNAL OF SCIENCE.**

---

[THIRD SERIES.]

---

FEBRUARY 1846.

XVI. *On the Application of the Photographic Camera to Meteorological Registration.* By HENRY COLLEN, Esq.\*

[With a Plate.]

IN April 1844, Mr. Ronalds applied to me for the purpose of obtaining some photographic representations of figures, forming "a sort of pictorial register of atmospheric electricity" upon glass plates coated with Canada balsam, which figures had been executed at the Kew Observatory by means of his electrograph, described in the Fourteenth Report of the British Association. The desired result was quickly obtained by the usual photogenic process, and also by the camera; the latter being found however, as was to be expected, the greatly superior mode. Several other impressions were afterwards made from figures on coated metallic plates, some of which were shown attached to Mr. Ronalds's report to the meeting at York. The sharpness and delicacy of the positive impressions thus obtained gave rise to some experiments, made by us conjointly, for the purpose of applying the photographic camera to the registration of Volta's electrometer, the thermometer, and the siphon barometer. The projection of shadows on photographic paper, which, by the way, had been already proposed and tried by several persons, was at once objected to by Mr. Ronalds, whose knowledge of the delicacy required in observing and registering the various instruments at the Observatory, made him fully aware of the necessity of obtaining as perfect definition as the best optical arrangement would produce; an excellent compound lens, made and kindly lent to us by Mr. Ross, was therefore used, and has been employed on each of the instruments, *i. e.* the electro-

\* Communicated by the Author.

meter, the barometer, and the thermometer, and a series of experimental observations permanently registered at Kew.

The accompanying figure (Plate III. fig. 1) is part of a day's registration of the effect of atmospheric electricity on Volta's electrometer; the gradual decline of daylight is shown, and also the continuation of the registration, by artificial light; without the use of the latter, it is obvious that the application of photography to these purposes would be very incomplete, if not wholly useless; and it may perhaps, in some cases, be advisable to make its use constant.

The various intensities of light from a *clouded* sky frequently give rise (of course) to variations in depth of tint on the paper, which thus becomes an approximation to Sir John Herschel's actinograph; and it may be here worth while to remark, that sometimes, when with *such a sky* these intensities of action on the paper are *augmented*, the electricity of serene weather manifests a tendency to *increase* also; this fact may be compared with the almost invariable tendency of the sun's light and heat, in a *clear* sky, to *diminish* the tension of those electrometers which receive their charges by absorption.

The calotype process is that which is used, being, of all those upon paper, the most sensitive, which quality is highly essential during the use of artificial light; it is very advantageously employed for these purposes, instead of the Daguerreotype, on account of its cheapness, and also on account of the facility with which representations can be obtained of any required length.

In the apparatus at present constructed, the paper is moved by a clock at the rate of one inch per hour, and is cut into pieces nine inches long; but for constant use they should be twelve inches long, so that by the introduction of two pieces during twenty-four hours, a continuous register of the effects would be preserved without further attention than the application of the artificial light (if not used constantly) at the decline of daylight; at present an Argand lamp is used, which, of course, requires some attention, but where available, a common gas-light would be greatly preferable; this however is not the case at the Kew Observatory, and for this reason only, the experiments have not been continued during the night.

The construction of the apparatus is very simple, although many tedious experiments have been made to produce the result; it consists essentially of the following arrangement:—The instrument to be registered is placed so as to be between the light and a lens of considerable aperture, with very short focus, and flat field of sufficient extent for the purpose; and



the paper is placed so as to be in the exact focus for obtaining an image of the same size as, or larger than, the object. When the electrometer is the instrument to be registered, the figures of the extreme ends only of the straws are allowed to fall upon the paper, an opaque diaphragm pierced with a slit, the curve of which is part of a circle of which the length of the straws is the radius, being placed very near the paper.

In the registration of the thermometer or barometer, the difficulty arising from the refraction of light by the glass tube was proposed to be met in two ways, the first of which, the one adopted, consists in the use of a diaphragm with a straight slit, which can be opened from, or contracted towards, its exact centre by a very simple arrangement, and is placed in front of the mercury, *i. e.* on the side next to the light, so as to regulate the quantity admitted; this regulation has also the effect of preserving the necessary sharpness of figure, which too much light tends to injure.

The second method proposed, which has not yet been tried, consists in the employment of a piece of glass tube, the bore of which is a trifle larger than the outside of the tube of the instrument; this, having two opposite surfaces ground flat and polished, and being long enough to include the range of variation, is cemented on to the tube of the instrument with Canada balsam, and would render it easy (by making all but a central slit opaque) to get rid of the partial illumination of the column of mercury on the side which is required, for a good impression on the paper, to be quite dark.

The surface of the mercury in the barometer sustains a blackened pith-ball of the same diameter as the bore of the tube, but freely sliding therein; it is proposed however to make a float of platinum foil with a sharp edge, which will probably be found to be more advantageous.

The thermometer used is mercurial, with a *broad* flat bore.

The wet-bulb, hair hygrometer, &c., as well as every other instrument which by its action affords a distinct sign, may obviously be registered in the same manner.

Several minor points of difficulty remain still to be overcome, but it is hoped that enough has been done to justify the expectation that the photographic camera may become a really useful and convenient instrument in the hands of the exact meteorologist.

The electrical experiments were made by means of a small conductor, insulated for the occasion; Mr. Ronalds not feeling either authorised, or disposed, to interrupt the course of observations carried on by means of the ordinary high conductor, until the proposed mode of registration is quite matured.

XVII. *On Fresnel's Theory of the Aberration of Light.* By G. G. STOKES, M.A., *Fellow of Pembroke College, Cambridge\**.

THE theory of the aberration of light, and of the absence of any influence of the motion of the earth on the laws of refraction, &c., given by Fresnel in the ninth volume of the *Annales de Chimie*, p. 57, is really very remarkable. If we suppose the diminished velocity of propagation of light within refracting media to arise solely from the greater density of the æther within them, the elastic force being the same as without, the density which it is necessary to suppose the æther within a medium of refractive index  $\mu$  to have is  $\mu^2$ , the density in vacuum being taken for unity. Fresnel supposes that the earth passes through the æther without disturbing it, the æther penetrating the earth quite freely. He supposes that a refracting medium moving with the earth carries with it a quantity of æther, of density  $\mu^2 - 1$ , which constitutes the excess of density of the æther within it over the density of the æther in vacuum. He supposes that light is propagated through this æther, of which part is moving with the earth, and part is at rest in space, as it would be if the whole were moving with the velocity of the centre of gravity of any portion of it, that is, with a velocity  $\left(1 - \frac{1}{\mu^2}\right)v$ ,  $v$  being the velocity of the earth. It may be observed however that the result would be the same if we supposed the whole of the æther within the earth to move together, the æther entering the earth in front, and being immediately condensed, and issuing from it behind, where it is immediately rarefied, undergoing likewise sudden condensation or rarefaction in passing from one refracting medium to another. On this supposition, the evident condition that a mass  $v$  of the æther must pass in a unit of time across a plane of area unity, drawn anywhere within the earth in a direction perpendicular to that of the earth's motion, gives  $\left(1 - \frac{1}{\mu^2}\right)v$  for the velocity of the æther within a refracting medium. As this idea is rather simpler than Fresnel's, I shall adopt it in considering his theory. Also, instead of considering the earth as in motion and the æther outside it as at rest, it will be simpler to conceive a velocity equal and opposite to that of the earth impressed both on the earth and on the æther. On this supposition the earth will be at rest; the æther outside it will be moving with a velocity  $v$ , and the æther in a refracting medium with a velocity

\* Communicated by the Author.

$\frac{v}{\mu^2}$ , in a direction contrary to that of the earth's real motion.

On account of the smallness of the coefficient of aberration, we may also neglect the square of the ratio of the earth's velocity to that of light; and if we resolve the earth's velocity in different directions, we may consider the effect of each resolved part separately.

In the ninth volume of the *Comptes Rendus* of the Academy of Sciences, p. 774, there is a short notice of a memoir by M. Babinet, giving an account of an experiment which seemed to present a difficulty in its explanation. M. Babinet found that when two pieces of glass of equal thickness were placed across two streams of light which interfered and exhibited fringes, in such a manner that one piece was traversed by the light in the direction of the earth's motion, and the other in the contrary direction, the fringes were not in the least displaced. This result, as M. Babinet asserts, is contrary to the theory of aberration contained in a memoir read by him before the Academy in 1829, as well as to the other received theories on the subject. I have not been able to meet with this memoir, but it is easy to show that the result of M. Babinet's experiment is in perfect accordance with Fresnel's theory.

Let  $T$  be the thickness of one of the glass plates,  $V$  the velocity of propagation of light in vacuum, supposing the æther at rest. Then  $\frac{V}{\mu}$  would be the velocity with which light would traverse the glass if the æther were at rest; but the æther moving with a velocity  $\frac{v}{\mu^2}$ , the light traverses the glass with a velocity  $\frac{V}{\mu} \pm \frac{v}{\mu^2}$ , and therefore in a time

$$T \div \left( \frac{V}{\mu} \pm \frac{v}{\mu^2} \right) = \frac{\mu T}{V} \left( 1 \mp \frac{v}{\mu V} \right).$$

But if the glass were away, the light, travelling with a velocity  $V \pm v$ , would pass over the space  $T$  in the time

$$T \div (V \pm v) = \frac{T}{V} \left( 1 \mp \frac{v}{V} \right).$$

Hence the retardation, expressed in time,  $= (\mu - 1) \frac{T}{V}$ , the same as if the earth were at rest. But in this case no effect would be produced on the fringes, and therefore none will be produced in the actual case.

I shall now show that, according to Fresnel's theory, the laws of reflexion and refraction in singly refracting media are

uninfluenced by the motion of the earth. The method which I employ will, I hope, be found simpler than Fresnel's; besides it applies easily to the most general case. Fresnel has not given the calculation for reflexion, but has merely stated the result; and with respect to refraction, he has only considered the case in which the course of the light within the refracting medium is in the direction of the earth's motion. This might still leave some doubt on the mind, as to whether the result would be the same in the most general case.

If the æther were at rest, the direction of light would be that of a normal to the surfaces of the waves. When the motion of the æther is considered, it is most convenient to define the direction of light to be that of the line along which the *same portion* of a wave moves relatively to the earth. For this is in all cases the direction which is ultimately observed with a telescope furnished with cross wires. Hence, if A is any point in a wave of light, and if we draw AB normal to the wave, and proportional to  $V$  or  $\frac{V}{\mu}$ , according as the light is passing through vacuum or through a refracting medium, and if we draw BC in the direction of the motion of the æther, and proportional to  $v$  or  $\frac{v}{\mu^2}$ , and join AC, this line will give the direction of the ray. Of course, we might equally have drawn AD equal and parallel to BC and in the opposite direction, when DB would have given the direction of the ray.

Let a plane P be drawn perpendicular to the reflecting or refracting surface and to the waves of incident light, which in this investigation may be supposed plane. Let the velocity  $v$  of the æther in vacuum be resolved into  $p$  perpendicular to the plane P, and  $q$  in that plane; then the resolved parts of the velocity  $\frac{v}{\mu^2}$  of the æther within a refracting medium will be  $\frac{p}{\mu^2}$ ,  $\frac{q}{\mu^2}$ . Let us first consider the effect of the velocity  $p$ .


It is easy to see that, as far as regards this resolved part of the velocity of the æther, the directions of the refracted and reflected waves will be the same as if the æther were at rest. Let BAC (fig. 1) be the intersection of the refracting surface and the plane P; DAE a normal to the refracting surface; AF, AG, AH normals to the incident, reflected and refracted waves. Hence AF, AG, AH will be in the plane P, and

$$\angle GAD = \text{FAD}, \quad \mu \sin \text{HAE} = \sin \text{FAD}.$$

Take  $AG = AF, \quad AH = \frac{1}{\mu} AF.$

Draw  $Gg$ ,  $Hh$  perpendicular to the plane  $P$ , and in the direction of the resolved part  $p$  of the velocity of the æther, and

Fig. 1.



$Ff$  in the opposite direction; and take

$$Ff : Hh : FA :: p : \frac{p}{\mu^2} : V, \text{ and } Gg = Ff,$$

and join  $A$  with  $f$ ,  $g$  and  $h$ . Then  $fA$ ,  $Ag$ ,  $Ah$  will be the directions of the incident, reflected and refracted rays. Draw  $FD$ ,  $HE$  perpendicular to  $DE$ , and join  $fD$ ,  $hE$ . Then  $fDF$ ,  $hEH$  will be the inclinations of the planes  $fAD$ ,  $hAE$  to the plane  $P$ . Now

$$\tan F D f = \frac{p}{V \sin F A D}, \quad \tan H E h = \frac{\mu^{-2} p}{\mu^{-1} V \sin H A E},$$


and  $\sin F A D = \mu \sin H A E$ ; therefore  $\tan F D f = \tan H E h$ , and therefore the refracted ray  $Ah$  lies in the plane of incidence  $fAD$ . It is easy to see that the same is true of the reflected ray  $Ag$ . Also  $\angle gAD = fAD$ ; and the angles  $fAD$ ,  $hAE$  are sensibly equal to  $FAD$ ,  $HA E$  respectively, and we therefore have without sensible error,  $\sin fAD = \mu \sin hAE$ . Hence the laws of reflexion and refraction are not sensibly affected by the velocity  $p$ .

Let us now consider the effect of the velocity  $q$ . As far as depends on this velocity, the incident, reflected and refracted rays will all be in the plane  $P$ . Let  $AH$ ,  $AK$ ,  $AL$  be the intersections of the plane  $P$  with the incident, reflected and refracted waves. Let  $\psi$ ,  $\psi_r$ ,  $\psi'$  be the inclinations of these waves to the refracting surface; let  $NA$  be the direction of the resolved part  $q$  of the velocity of the æther, and let the angle  $NAC = \alpha$ .

The resolved part of  $q$  in a direction perpendicular to  $AH$  is  $q \sin(\psi + \alpha)$ . Hence the wave  $AH$  travels with the velocity  $V + q \sin(\psi + \alpha)$ ; and consequently the line of its inter-

section with the refracting surface travels along A B with the

Fig. 2.



velocity  $\text{cosec } \psi \{V + q \sin (\psi + \alpha)\}$ . Observing that  $\frac{q}{\mu^2}$  is the velocity of the æther within the refracting medium, and  $\frac{V}{\mu}$  the velocity of propagation of light, we shall find in a similar manner that the lines of intersection of the refracting surface with the reflected and refracted waves travel along A B with velocities

$$\text{cosec } \psi_i \{V + q \sin (\psi_i - \alpha)\}, \quad \text{cosec } \psi' \left\{ \frac{V}{\mu} + \frac{q}{\mu^2} \sin (\psi' + \alpha) \right\}$$

But since the incident, reflected and refracted waves intersect the refracting surface in the same line, we must have

$$\left. \begin{aligned} \sin \psi_i \{V + q \sin (\psi_i - \alpha)\} &= \sin \psi \{V + q \sin (\psi_i - \alpha)\}, \\ \mu \sin \psi' \{V + q \sin (\psi_i - \alpha)\} &= \sin \psi \left\{ V + \frac{q}{\mu} \sin (\psi' + \alpha) \right\}. \end{aligned} \right\} \quad (\text{A})$$

Draw H S perpendicular to A H, S T parallel to N A, take S T : H S :: q : V, and join H T. Then H T is the direction of the incident ray; and denoting the angles of incidence, reflexion and refraction by  $\phi$ ,  $\phi_i$ ,  $\phi'$ , we have

$$\phi - \psi = \text{S H T} = \frac{\text{S T} \sin \text{S}}{\text{S H}} = \frac{1}{V} \times \text{resolved part of } q \text{ along A H}$$

$$= \frac{q}{V} \cos (\psi + \alpha). \quad \text{Similarly,}$$

$$\phi_i - \psi_i = \frac{q}{V} \cos (\psi_i - \alpha), \quad \phi' - \psi' = \frac{q}{\mu V} \cos (\psi' + \alpha):$$

whence  $\sin \psi = \sin \phi - \frac{q}{V} \cos \phi \cos (\phi + \alpha),$

$$\sin \psi_i = \sin \phi_i - \frac{q}{V} \cos \phi_i \cos (\phi_i - \alpha),$$

$$\sin \psi' = \sin \phi' - \frac{q}{\mu V} \cos \phi' \cos (\phi' + \alpha).$$

On substituting these values in equations (A), and observing that in the terms multiplied by  $q$  we may put  $\phi_1 = \phi$ ,  $\mu \sin \phi' = \sin \phi$ , the small terms destroy each other, and we have  $\sin \phi_1 = \sin \phi$ ,  $\mu \sin \phi' = \sin \phi$ . Hence the laws of reflexion and refraction at the surface of a refracting medium will not be affected by the motion of the æther.

In the preceding investigation it has been supposed that the refraction is out of vacuum into a refracting medium. But the result is the same in the general case of refraction out of one medium into another, and reflexion at the common surface. For all the preceding reasoning applies to this case if we merely substitute  $\frac{p}{\mu'^2}$ ,  $\frac{q}{\mu'^2}$  for  $p$ ,  $q$ ,  $\frac{V}{\mu'}$  for  $V$ , and  $\frac{\mu}{\mu'}$  for  $\mu$ ,  $\mu'$  being the refractive index of the first medium. Of course refraction out of a medium into vacuum is included as a particular case.

It follows from the theory just explained, that the light coming from any star will behave in all cases of reflexion and ordinary refraction precisely as it would if the star were situated in the place which it appears to occupy in consequence of aberration, and the earth were at rest. It is, of course, immaterial whether the star is observed with an ordinary telescope, or with a telescope having its tube filled with fluid. It follows also that terrestrial objects are referred to their true places. All these results would follow immediately from the theory of aberration which I proposed in the July number of this Magazine; nor have I been able to obtain any result, admitting of being compared with experiment, which would be different according to which theory we adopted. This affords a curious instance of two totally different theories running parallel to each other in the explanation of phænomena. I do not suppose that many would be disposed to maintain Fresnel's theory, when it is shown that it may be dispensed with, inasmuch as we would not be disposed to believe, without good evidence, that the æther moved quite freely through the solid mass of the earth. Still it would have been satisfactory, if it had been possible, to have put the two theories to the test of some decisive experiment.

XVIII. *Observations on the Development and Growth of the Epidermis.* By ERASMUS WILSON, F.R.S., Lecturer on Anatomy and Physiology in the Middlesex Hospital\*.

IT is the commonly received doctrine at the present day, that the cells of the epidermis and of epithelium in general, originate out of materials furnished by the liquor sanguinis or plasma of the blood. In order that this purpose may be effected, the liquor sanguinis is conveyed by endosmosis through the walls of the capillary vessels and through the peripheral boundary of the surface, the "basement membrane" of Bowman. Having reached the exterior plane of the latter, the changes commence which result in the development of granules in the previously fluid liquor sanguinis, or rather, perhaps, in the aggregation of the molecules of the organisable material or blastema, which was previously held in intimate suspension or solution by the liquor sanguinis. Out of the body an action of this kind would be termed coagulation, and where inorganic matter is concerned, crystallization. The process to which I am now referring, though taking place within the body, is analogous to these phænomena, with the difference of being controlled and directed by the power of life, of being, in point of fact, a vital coagulation or crystallization. Indeed, coagulation, although occurring out of the body, and sometimes after the lapse of a considerable period, may be regarded as the last act of vital existence, or as a vestige of the atmosphere of life with which the coagulating fluid was previously charged in abundance.

As regards the tissue under consideration, there is every ground for belief, that the organisable material or blastema of the liquor sanguinis is appropriated by the epidermis the very instant it reaches the exterior plane of the "basement membrane;" some portion of it, and the greater part of the serum of the liquor sanguinis, being taken up by the newly-formed cells to be transmitted in succession to more superficial ranges of cells, and the remaining portion being converted on the spot into the primitive granules of the tissue. This belief is supported by the fact of the absence of any fluid stratum between the epidermis and the dermis, and by the close connexion known to subsist between those two membranes. It is well known that to separate the epidermis from the dermis, until the former is so thoroughly saturated with fluid by maceration as to have acquired a considerable addition to its dimensions in all directions, or until decomposition has com-

\* Read before the Royal Society, June 19, 1845, and communicated by the Author.



menced, is next to impossible; and in the living state of the body, separation never takes place until the mutual connexion between the layers has been destroyed by the effusion of fluid. The microscope gives additional weight to this evidence. I have observed that the cells of the deep surface of the epidermis are in immediate contact with the boundary limit of the dermis, and that moreover it is frequently difficult to determine the exact line between them. I have also made the following experiment:—I cut very thin vertical slices of the skin, at daily periods, from the moment of death until decomposition had become established, and submitted them to the action of the compressor on the field of the microscope, but in every instance, while fresh, the two tissues yielded to the pressure in equal proportion without any separation occurring. As soon, however, as decomposition had commenced, separation was produced, and in the early stages took place with difficulty. This experiment proves that the firm adhesion subsisting between the epidermis and dermis is not alone due to the numerous inflexions of the former into the latter, which take place at the sudoriferous tubes, hair tubes, and sebaceous ducts, although these inflexions must co-operate powerfully in the result.

Being desirous of examining the under surface of the epidermis with the higher powers of the microscope, and failing in all my attempts to effect this object by taking the entire thickness of the epidermis or scraping, I awaited the first indication of its separation from the dermis, and then removing it carefully made a thin slice parallel with the surface which I wished to examine. This plan succeeded beyond my expectations; for not only did I obtain parts so diaphanous as to enable me to see the surface distinctly, but the septa between the depressions for the papillæ of the dermis afforded natural laminae of such transparency as permitted their structure to be well examined.

When the under surface of the epidermis was exposed to view, I found it to be composed of four kinds of elements, arranged in such a manner as to constitute an irregular mosaic plane. These elements are,—1, *granules*, measuring about  $\frac{1}{20000}$ th of an inch in diameter; 2, *aggregated granules*, measuring about  $\frac{1}{10000}$ th; 3, *nucleated granules*, measuring  $\frac{1}{6000}$ th to  $\frac{1}{4000}$ th; 4, *cells*, measuring  $\frac{1}{3000}$ th to  $\frac{1}{2500}$ th of an inch.

1. The granules, which I may distinguish by the name of *primitive granules*, are globular in form, homogeneous, solid, brightly illumined by transmitted light when the centre is under the focus of the microscope, but dark when viewed upon the surface, the darkness being increased whenever they are

congregated in clusters. These granules I conceive to be the first organic shape of the blastema of the liquor sanguinis.

2. The *aggregated granules*, measuring about  $\frac{1}{10000}$ th of an inch in diameter, are minute masses, composed of four, five or six of the preceding, or as many as can be aggregated without leaving an unoccupied space in the centre of the mass. With an imperfect focus, these granules have the appearance of possessing a transparent globular nucleus, but this appearance ceases when the focus is perfect, and then the component granules are quite obvious, and the centre becomes a dark point, namely, the shadow caused by the meeting of the primitive granules.

3. The *nucleated granules*, measuring between  $\frac{1}{8000}$ th and  $\frac{1}{4000}$ th of an inch in diameter, are in point of construction an "aggregated granule" with a single tier of "aggregated granules" arranged around it, so as to give the entire mass a circular or oval form. The central aggregated granule has now become a nucleus, and at the same time has undergone other changes which indicate its longer existence. For example, the primitive granules composing it are denser than they were originally, and they are separated from each other by a very distinct interstitial space filled with a transparent and homogeneous matter. Sometimes this interstitial substance presses the granules asunder equally on all sides, constituting a circular nucleus; but more frequently two opposite granules are more widely separated than the rest, and the nucleus receives an elongated form. The interstitial substance is most conspicuous at the line of junction of the nucleus with the secondary tier of "aggregated granules," and in this situation gives a defined character to the nucleus. Close observation and a perfect focus render it quite obvious that the peripheral tier of granules are in reality aggregated, they are lighter than the shaded granules of the nucleus, and apparently softer in texture.

The nucleated granules are more or less flattened in form, and present a flat surface of contact with the dermis. It is the latter circumstance that gives the facility of determining their mode of construction.

4. The cells of the deep stratum of the epidermis, measuring  $\frac{1}{3000}$ th to  $\frac{1}{2500}$ th of an inch in their long diameter, are the most striking feature of this layer, and may be said to be its chief constituent. They originate, as is evident from their structure, in the nucleated granules previously described, and consist of a transparent layer added to the exterior of the former; or, if I might be permitted to describe them as they appear in their tessellated position, they are constituted by the

addition of a transparent border to the last-described nucleated granule. The periphery of this transparent border is bounded by a dark interstitial substance, which gives the border a defined outline; and in the latter situation I imagine a cell-membrane to exist. I am not satisfied, however, that this is the case; and the difficulty of isolating these cells, and their roughness of outline when separated, serve to prove that if a membrane be really present, it must be exceedingly thin and easily torn. Assuming therefore, from analogy rather than from demonstrative evidence, that there exists a boundary membrane to the bodies I am now describing, I have termed them "cells;" the cavity of the cell I apprehend to be the "transparent border," the "nucleated granule" is the *nucleus* of the cell, the "aggregated granule" of the latter the *nucleolus*, and the entire body a "nucleolo-nucleated cell."

Before quitting the structure of the "nucleolo-nucleated cell," or primitive cell of the epidermis, there is a point of much interest to be mentioned with regard to it, which is, that the "transparent border" just described is itself a tier of "aggregated granules." The nucleolus therefore is an "aggregated granule," the nucleus a tier (taking its flat surface) of "aggregated granules" surrounding the former, and the cell a tier of "aggregated granules" enclosing the whole.

To return to the mosaic-like plane of the under surface of the epidermis, the largest of the pieces composing this plane are the nucleolo-nucleated cells. These are placed without order, some being closely pressed together, others being separated by moderate intervals, and here and there some separated by interspaces equal to the breadth of the cells. The interspaces, or intercellular spaces, are occupied by the "nucleated granules," "aggregated granules," and "primitive granules," irregularly set in a homogeneous interstitial substance, which fills up all vacuities. The granules and interstitial substance modify the light transmitted through them variously at different foci of the microscope; sometimes the granules look dark while the interstitial substance is light, and sometimes the reverse is the case.

Such is the structure of the mosaic-like plane of the under surface of the epidermis, and so far, my observations, having reference to facts, are demonstrable and admit of being spoken to positively. The interpretation of the facts I would willingly leave to others, but feel that I am called upon to state any opinion, founded on the above observations, that I may have formed of the signification of these appearances. In the first place, then, I must acknowledge myself wholly divided between

a belief in the formation of the "aggregated granule" by the *aggregation of primitive granules*, the idea which prompted me to give them that name, and the formation of the aggregated granule by the cleavage of a primitive granule. If this question related merely to the formation of the "primary aggregated granule," it would be unimportant, but it has a more extended application. The outermost layer of the nucleus is composed, as I have shown, of "aggregated granules," and so also is that layer which alone forms the space in the nucleolo-nucleated cells. To them the hypothesis of cleavage of a simple granule would be most suitable, and this theory would explain better than any other, changes which remain to be described in the further growth of the epidermic cell. In the second place, the relation of cell and nucleus is a question on which I feel considerable doubt. The process of development appears to consist in the successive production of granules, one layer of granules succeeding another, so that if the organisable principle exist in each separate granule, the organisable force may be supposed to be more and more weakened in successive formations, until the moment arrives when it ceases entirely. Is that which I have described as a nucleolo-nucleated cell really a cell or still a nucleus? The only solution of the question that occurs to me is, determining the presence of a cell-membrane, in which I have not satisfactorily succeeded.

Admitting the nucleolo-nucleated bodies now described to be cells in their earliest state of formation, their size is  $\frac{1}{3000}$ th to  $\frac{1}{2300}$ th of an inch in the long diameter, and that of their nucleus from  $\frac{1}{6000}$ th to  $\frac{1}{4300}$ th of an inch. In the stratum immediately above the deepest layer, I find cells measuring  $\frac{1}{2000}$ th of an inch with nuclei of  $\frac{1}{4300}$ th. Above these, cells measuring  $\frac{1}{1800}$ th, with nuclei varying from  $\frac{1}{4000}$ th to  $\frac{1}{3000}$ th, and above the latter cells measuring  $\frac{1}{1300}$ th, with nuclei of  $\frac{1}{2300}$ th. In following the layers of the epidermis upwards to the surface, cells may be observed possessing every intermediate degree of size between the last-mentioned cell, namely  $\frac{1}{1300}$ th and  $\frac{1}{600}$ th, which is the measurement of the scales which constitute the uppermost stratum of the epidermis. It must not be supposed, however, that the growth of the epidermic cells reaches its maximum only at the surface: I have found cells of that magnitude in the deeper strata; and there is every indication of the growth of these cells being completed in the stratum immediately above the mosaic-like layer.

Young cells are remarkable for the large size of the nucleus as compared with the entire bulk of the cell; and it is quite

evident also that the nuclei, up to a certain point, grow with the cells, their mode of growth appearing to be, the separation of the original granules by the deposition between them of interstitial matter; and in addition, as I believe, by cleavage of the latter and the consequent multiplication of the granules. In cells measuring  $\frac{1}{2000}$ th and  $\frac{1}{1800}$ th of an inch, I found the granular character of the nucleus to be very manifest. Besides growth, it is apparent that other changes are taking place in the nucleus; imbibition and assimilation of organisable material must necessarily be in action in order to accomplish the formation of interstitial matter; but in addition to this the central granules undergo another change, by which they are altered in character and become distinguished from the rest when submitted to chemical experiment. For example, when dilute acetic acid is added to the cells measuring  $\frac{1}{2000}$ th of an inch and less, the entire nucleus is rendered transparent and less discernible than before; but when cells of a somewhat larger size, and consequently longer growth, are submitted to the same process, the nucleus is rendered much more distinct than it was previously. But the body which is made so conspicuous in this latter experiment is not the entire nucleus, but simply the central and older granules of the nucleus; the younger granules retain the character of those of the young cells, they are made more transparent than they were before, and have faded from sight. I may mention also, that the nucleus brought into view by the acetic acid is more or less irregular in form, and has the appearance of being constituted by the fusion of the original granules. How much of this appearance may be real and how much the effect of the acid, I do not pretend to say, and I set no value on the experiment beyond the demonstration of the mere fact which it is made to illustrate.

I now turn to the growth of the cells. I have remarked, in an earlier paragraph, that the formation of the young cell appears to be due to the development of a stratum of "aggregated granules" externally to the nucleated mass, which I have regarded as the cell-nucleus. Now nothing is more certain than that the growth of the cell is due to a successive repetition of this process, the growth of the cell-membrane being consentaneous with the development and growth of "aggregated granules" within it. In cells of  $\frac{1}{1800}$ th to  $\frac{1}{1500}$ th of an inch, the "aggregated granules" of the periphery are not easily discernible; but in cells measuring  $\frac{1}{1000}$ th, and thence upwards to the complete size of the epidermic cell, the fact is quite evident, and is apparent even in the cell-scale. Indeed a cell, at the full period of growth, is a kind of cell-microcosm,

containing in its interior secondary cells, tertiary cells, nucleolo-nucleated cells, nucleated granules, aggregated granules, and primitive granules.

It will be observed that this hypothesis of cell-growth differs from that of Schwann. The theory of Schwann always appeared to me to be incompetent to the explanation of the growth of the large scale of epidermis and epithelium in a tissue manifestly subjected to considerable pressure. I sought in vain for the watch-glass cells, elliptical cells, and globular cells in the epidermis; but my search has been rewarded by the discovery of the above-described beautiful process of formation and growth. It will be seen that, according to this view of the growth of the epidermic cells, they never possess anything approaching to a globular shape, that the scales are not flattened spheres, but on the contrary always possessed a flattened form, and have increased by a peripheral growth. This mode of growth again is made manifest by the observation of a vertical section of the epidermis. The most careful examination can distinguish no difference between the size of the deeper and the superficial strata of cells; they have all the same average thickness, all the same average length,—an appearance easily explained when we regard them as parent-cells containing secondary and tertiary cells of the same average size as the cells of earlier formation. It is true that the complete size of a cell is very quickly attained, and that its growth, taking place in the deepest stratum of the epidermis, could not be expected to produce any difference of character in the middle and superficial strata; but this is not mentioned, as far as I know, by Schwann.

The process of growth here described explains also the fact of the disappearance of the nucleus in the scales of epidermis. The outermost granules of the nucleus have become the nuclei or nucleoli of secondary cells, and have consequently been moved away from their original position in the performance of their office of centres of growth to secondary cells. The original nucleus, therefore, is not lost, but merely robbed of some of its component granules, which may be discovered in many parts of the epidermic scale, instead of being concentrated in a single mass. In these scales, and particularly in epithelial scales, the central and dense part of the original nucleus is generally perceptible; in the latter it constitutes the scale nucleus; and in the epidermic scale there is always some one little mass larger than the rest, particularly if the scale have been for some time immersed in fluid, as when it is examined in the serum of a blister. In an epidermic cell, measuring  $\frac{1}{800}$ th of an inch in long diameter, I found several

secondary cells measuring  $\frac{1}{1500}$ th, others measuring  $\frac{1}{3000}$ th; and in the interstices, primitive granules, aggregated granules, and nucleated cells.

My observations, it will be seen, have been chiefly directed to the epidermis, and I am prevented at present from carrying them further, but I have no doubt that the epithelium will be found to be identical in the phenomena of development and growth with the epidermis. I have observed the same structure in the epithelium of the mouth and fauces, and also in that of the bladder and vagina. Incomplete epithelial cells, measuring  $\frac{1}{750}$ th and  $\frac{1}{700}$ th of an inch from the fauces, presented a very remarkable appearance; they had a rounded lobulated border, evidently composed of a row of secondary cells and a depressed centre, as though the action were subsiding in the latter, while it was progressing in the circumference.

Another illustration of the structure now described, I found in the cells of melanosis, and in the pigmentary cells of the choroid membrane of the eyeball. I am induced to believe that the same structure will be discovered more extensively than at present can be anticipated. The corpuscles of melanosis, according to my observations, are parent-cells, having an average admeasurement of  $\frac{1}{1000}$ th of an inch, containing secondary cells and nucleated and aggregated granules, as well as separate primitive granules. The "aggregated granules" measured from  $\frac{1}{11000}$ th to  $\frac{1}{7000}$ th of an inch, and the primitive granules about  $\frac{1}{20000}$ th.

There is another feature in the history of development of the epidermic cell, which I regard as peculiarly interesting. This relates to an organic change taking place in the assimilative powers of the primitive granules, by which the latter are altered in their colour, in short, are converted into "pigment granules." Pigment granules appear to differ in no respect from the "primitive granules," excepting in that of colour, and perhaps also in chemical composition. They have the same globular form, the same size, and occupy the same position in the cell, being always accumulated around the nucleus, and dispersed less numerously through the rest of the cell. The nucleus of the cell in the epidermis of the Negro appears to consist wholly of pigment granules, while in the European there is a greater or less admixture of coloured and uncoloured granules. The central granules are generally lighter in tint than the rest, and give the idea of a colourless nucleolus, while those around the circumference are deeper coloured. Besides a difference in the depth of colour of the separate granules entering into the composition of a single

cell, there is also much difference in the aggregate of the granules composing particular cells. For example, intermingled with cells of a dark hue, there are others less deeply tinted, which give the tissue in which they are found a mottled appearance. This fact is well-illustrated in the hair, and also in the nails, in which latter it is no uncommon thing to find an isolated streak produced by the accumulation of a number of cells containing coloured granules, in the midst of colourless cells.

When pigment granules are examined separately, they offer very little indication of the depth of colour which is produced by their accumulation. I have observed some to have the hue of amber, while others scarcely exceeded the most delicate fawn. The depth of colour of the deep stratum of the epidermis in the Negro, is evidently due to the composition of that layer of these granules, while the grayness of the superficial layers of the same tissue results not merely from the desiccation of these granules, but also from the fact of those subsequently produced being less strongly coloured, and also from the addition of a colourless cell-membrane.

The epidermic scale of the Negro has a mottled appearance, from the numerous secondary nuclei and their attendant coloured granules which are scattered through its texture.

---

P.S. Since my communication of the above paper to the Royal Society, I have confirmed its truth by further observations, and have ascertained that the same principle of growth is applicable to the formation of mucus and pus-corpuscles.

December 1845.

---

XIX. *On the Aberration of Light, in Reply to Mr. Stokes.*  
By the Rev. J. CHALLIS, M.A., Plumian Professor of Astronomy in the University of Cambridge\*.

**T**HE remarks Mr. Stokes has made on my Explanation of the Aberration of Light, since they have little reference to the more important parts of the communication, require from me but brief notice.

I agree with all Mr. Stokes has said about the direction of vision through a telescope, but cannot perceive what it has to do with aberration. In selecting the wire of an astronomical telescope for the terrestrial object to which the direction of the celestial object is referred, I had not the least reference to vision through a telescope. It would have answered my pur-

\* Communicated by the Author.





just before it enters the eye, is the *true* direction of the object, atmospheric refraction not being considered.

It seems probable also that this is the direction in which the object is *seen*; if so, the point  $p$  is seen out of its true place. This, however, is not an essential consideration.

I think it important to remark, that the foregoing explanation of aberration rests on no hypothesis whatever, being a strict deduction from ascertained facts, without reference to any theory of light. The cause assigned for aberration is, therefore, a *vera causa*, which consequently excludes every explanation of a hypothetical kind, such, for instance, as that which Mr. Stokes proposed in the July number of this Magazine.

Aberration being explained in this manner, it is interesting to inquire whether a proposed theory of light be consistent with this explanation. The object of such an inquiry would be to test the truth of the proposed theory. The only condition the theory is required to fulfil, in addition to that of temporary transmission, is, that the light from an object traverse, just before it enters the eye, a straight line directed to the true position of the object.

The above condition is satisfied on the theory of emission, because according to that theory light passes from the object to the eye in a straight line. In the undulatory theory, the direction of transmission of light is the direction of transmission through space of a given point of a wave in a given phase of vibration. Where the æther is undisturbed, this direction is normal to the front of the wave. Where the æther is in motion, it is the direction resulting from the composition of the motion of propagation of the wave with the motion of translation of the æther. It is easy, therefore, to determine, for a given motion of translation of the æther, the angle which the normal to the front of the wave makes with the direction of transmission of light. In the figure, let  $p'n$  (not necessarily equal nor parallel to  $e'e$ ) represent the motion of the æther,  $p'e$  representing the velocity of light; then  $en$  is the direction of the normal to the wave that enters the eye at  $e$ . If the normal underwent no angular deviation the whole distance from the object to the eye,  $en$  would also be the direction of the object, and consequently aberration on this theory would not be accounted for. I gave Mr. Stokes the credit of having first shown that the normal is shifted through a certain angle as the wave is propagated through the æther set in motion by the earth, and by reasoning as he has done, and supposing certain analytical conditions, which I shall speak of presently, to be satisfied, the deviation is found to be from  $p'$  towards  $n$ , exactly through the angle  $p'en$ . Consequently  $ep'$  is the di-

rection of the object, and the required condition is satisfied by the undulatory theory of light.

I admit the correctness of Mr. Stokes's strictures on that part of my communication to which he principally objects. Mr. Stokes's own reasoning in the July number, or the following, may be substituted for the part objected to. The point  $a$  is carried with the velocity  $V-w$ , and the point  $b$  with the velocity  $V-w'$ , in the direction of the axis of  $z$ . As  $w$  is less than  $w'$ ,  $a$  is carried further than  $b$  in the small time  $\delta t$ , by  $(V-w)\delta t - (V-w')\delta t$ , that is, by  $(w'-w)\delta t$ , or  $\frac{dw}{dx}\delta x\delta t$ .

Dividing by  $\delta x$ , the interval between  $a$  and  $b$ , the angular displacement of the front of the wave in the plane of  $zx$  is  $\frac{dw}{dx}\delta t$ ,

which is equal to  $\frac{dw}{dx} \cdot \frac{\delta z}{V}$ , since  $V = \frac{\delta z}{\delta t}$  very nearly. To in-

tegrate this expression, it is necessary to assume that  $\frac{dw}{dx} = \frac{du}{dz}$ .

So considering the motion in the plane  $zy$ , the integration requires that  $\frac{dw}{dy} = \frac{dv}{dz}$ . These conditions, which are alluded to

above, I agreed with Mr. Stokes that it was necessary the motion of the æther should satisfy. I went a step further, and endeavoured to show that they do not restrict the motion. The reasoning for this purpose was based on hydrodynamical equations, in which the squares of the velocities were neglected. This may generally be done when the motion is small. But obviously all cases of motion for which  $\frac{du}{dt}, \frac{dv}{dt}$ ,

and  $\frac{dw}{dt}$  vanish are to be excepted, and the instance before us

may be one of this class; for the motion must be nearly symmetrical about the line in which the earth's centre moves, and if the earth's centre be taken for origin of co-ordinates, the velocity must be very approximately a function of co-ordinates independent of the time. On this account I doubt the applicability of those equations, and in the present state of our knowledge of the subject, it seems the best course simply to suppose the motion of the æther to be such as to satisfy the

two conditions  $\frac{dw}{dx} = \frac{du}{dz}$  and  $\frac{dw}{dy} = \frac{dv}{dz}$ . There is nothing improbable in the supposition: it saves the undulatory theory; but I must protest against its being considered necessary for the explanation of the aberration of light.

Cambridge Observatory, January 8, 1846.

XX. *Observations on certain Molecular Actions of Crystalline Particles, &c; and on the Cause of the Fixation of Mercurial Vapours in the Daguerreotype Process.* By AUGUSTUS WALLER, M.D.\*

[With a Plate.]

WHEN a piece of glass is covered with a solution containing the double phosphate of ammonia and magnesia, and traces are made upon it by any hard body, it is known that they become visible shortly afterwards by the salt being precipitated upon them. Berzelius, who mentions this test in his *Elements of Chemistry*, states that Wollaston proposed to make use of this fact as a test of the presence of magnesia in solution, which has since been frequently adopted. According to Berzelius, "the cause of this property is of a mechanical nature, probably from the glass being covered with microscopic crystals, the facets of which take a different position on the traces, for some reason which is not easily explained." More recently, Prof. Liebig has alluded to this subject in his *Vegetable Physiology*, § 157. These effects are referred by him to a state of unstable equilibrium of the various particles which compose the liquid, which is destroyed whenever a dynamical action is created sufficiently powerful to overcome the feeble attractions, or the inertia of the molecules in solution. He ascribes to the same cause the sudden solidification of water, which had remained liquid when below the freezing-point, upon being agitated; the precipitation of a mixture of potash and tartaric acid; also the detonation of fulminating powder from the contact of any solid body. Neither of these eminent observers mentions having submitted these traces to microscopic examination, although that is the only manner to test the hypothesis advanced by Berzelius.

On the present occasion it is my intention to describe some observations I have made, in order to elucidate the influence of molecular action on the precipitation of saline bodies, similar to that observed in the double phosphate, and to show that a similar influence is exerted over bodies in a gaseous state and in a state of vapour, and afterwards to point out some phænomena hitherto unexplained, such as the fixation of the mercurial vapours in the Daguerreotype for instance, which evidently depend upon a like cause.

In order to obtain the double phosphate, I have generally used a solution containing about ten grains of phosphate of soda with about three of carbonate of ammonia in an ounce and a half of water. I have preferred this mixture, because the ingredients are more easily procured, and are less acted

\* Communicated by the Author.

upon by the atmosphere than the phosphate of ammonia. The magnesian solution was generally a few grains of sulphate of magnesia to the same quantity of water as above.

A small quantity of the first mixture is poured on a piece of glass, and to this are added a few drops of the magnesia in solution; if it be allowed to remain undisturbed, in a few minutes the surface of the liquid becomes covered with a thin film, and on the glass appear minute shining crystals; but if before these crystals have time to form, any solid substance, as a glass rod or an empty pen, for instance, is passed over the glass through the liquid, the course it follows becomes visible shortly after. The images which are thus formed are double, and may be termed the upper and lower images.

I will first describe the upper images:—They appear on the surface of the liquid itself, when the film would otherwise have been formed. They are seen immediately after the passage of the pen through the liquid, whereas the lower ones only become apparent a few moments after. Being formed on a moveable surface, they are not perfect representations of the traces that have been made, and are changed and distorted by any movement of the liquid. When the solution of the salt is weak, they frequently disappear a few moments after their formation and are redissolved in the liquid; when the liquid is more concentrated, they likewise disappear, owing to the formation of the film on the surface. The production of these images appears to be independent of the chemical nature of the body used for tracing. They may be obtained independently of the lower ones, by drawing a thread gently over the surface of the liquid, without its coming in contact with the surface of the glass.

The lower images are formed on the surface of the glass, under the upper ones. A few seconds after the tracing has been made upon the glass, they begin to appear, and gradually become more distinct. The space of time which elapses before their appearance depends upon the strength of the solution. When it is strong they appear quickly, and when weak they take several minutes before they are visible.

To cause the formation of any images, the tracing must always be made after the mixture of the two solutions; under no other circumstances have I been able to create them. Thus, when the tracing is made on a perfectly dry glass, or on one slightly wet, and then immediately covered with the solution, no images will be created. This is likewise the case when we make traces in either the magnesian or the phosphate solution before their mixture together.

The passage of any solid substance in the proper solution

on glass will cause the formation of a deposit. Wood, glass, slate, and other similar substances, all have equal power in this respect, but metallic substances are less active. Other polished surfaces may be used instead of the glass plate, and I have formed these images on quartz and agate with the same effect.

The difference of crystalline texture exerts no influence, but the images seem to be with more difficulty produced on polished silver and copper than on a vitreous surface.

A very slight degree of friction will excite the formation of an image, although a moderate degree of pressure is more favourable.

Electricity exerts no influence in the formation of these images. In one experiment, in order to diminish the friction, I adapted two fine wires of a spiral form to a battery sufficiently strong to decompose water freely. These wires were moved through the solution in various directions, and the marks of the passage of the two poles became equally apparent without any difference on either side; and when afterwards disconnected from the battery and used in a similar manner, they produced the same effects.

It is remarkable with what fidelity the traces of lines become visible in this manner. Letters thus formed by a pen, are much more faithfully rendered than when written on paper with ink, and lines may be formed which are scarcely visible to the naked eye. Microscopic inspection shows this extreme exactness to a much greater degree than could have been anticipated; for we see a simple line become as it were decomposed into a number of parallel lines, which represent the point of contact between the two solids (see Plate III. fig. 2). These lines are composed of very minute and confused crystals, of an irregular appearance and joined together. Their diameter varies from 0.02 of a millimetre to about double that size. Between these parallel lines are frequently seen others still more minute. The other crystals which become deposited by the common crystalline powers over the untouched parts of the glass, are much larger than either of these. When the point of intersection of two lines is examined under the microscope, we perceive the appearance represented. While crystalline masses are in process of formation, it is impossible to prevent the deposition of crystals on other parts of the glass; but if while these are fresh they are subjected to a sharp current of water, the irregular crystals are mostly carried away, while the images are left almost intact. It is therefore evident that the same power which causes this deposit, renders them more adherent to the surface of the glass than the other cry-

stals. Another method of demonstrating the difference of their adherence, is by allowing the solution to dry on the glass, when by brushing it slightly with the feather of a pen, most of the irregular crystals are taken off and the images remain.

*Other substances capable of forming a like deposit.*—Chloride of platinum and nitrate of potash, mixed together, form a double chloride, with which images can be obtained with as much ease as with the double phosphate. The only difference is, that the double chloride precipitates in the shape of octahedrons, &c. Solutions of tartaric acid and nitrate of potash deposit crystals of bitartrate of potash, which are capable of forming upper and lower images with nearly as much facility as the double phosphate. The lower images formed by the bitartrate differ in one respect from those by the phosphate, for shortly after their formation they appear to lose their adhesion to the glass, and the slightest agitation of the liquid causes them to be detached; and if a sentence has been written, the curious appearance is presented of fragments of words and letters floating about in confusion. Under the microscope also they differ, fewer parallel lines are perceived, and the crystals are larger and unequal in size. Liquor potassæ added to a solution of tartaric acid will form images exactly similar to those just mentioned. Caustic soda and tartaric acid produce the same result, but the solution must be much more concentrated.

*Images formed by gaseous bodies.*—These traces are formed in the same manner as those which are crystalline, by passing a solid body over a piece of glass covered with a liquid containing a gas in solution, when they are immediately perceived by the bubbles which are deposited. On account of the specific gravity of the gas, these images are not very durable, for after a short time the gas which composes them rises to the surface. As a general rule, the ingredients, whose combination causes the formation of the gas, should be added together gently, and so diluted that whatever gas is formed they remain dissolved in the liquid. I have been surprised to find how much gas may be in this way made to remain in solution; and as most of them appear capable of being dissolved in this unstable manner, traces may be obtained from them all; and I have ascertained by experiment, that such is the case with carbonic, acetic and hydrochloric acids.

To obtain carbonic acid, I have generally used the subcarbonate of soda and tartaric acid. Acetate of ammonia was employed to liberate acetic acid, and hydrochloric acid was obtained from common salt and sulphuric acid. A mixture

capable of forming traces has the property of disengaging its gas in bubbles, whenever it is brought in contact with any dry surface; as for instance, when a mixture of this sort formed on a slip of glass is caused to spread over a part of the surface which has not previously been wetted, bubbles of gas are immediately evolved on that spot, although none are perceived elsewhere. This effect is also produced with champagne, seltzer and other effervescing waters, which however have not the property of forming gaseous traces. Any surface, whether metallic or non-metallic, will be found to effect the separation of the gas from the liquid; and I have not perceived that there was any difference from the surface being perfectly polished or rough.

The immersion of a piece of bread in champagne to renew the effervescence, is merely an example of the contact of a fresh surface with the gas; in a short time it ceases to have this effect, but if a fresh piece is used, the effervescence is renewed as before. The difference of effect between this and a piece of metal arises solely from the superior extent of surface presented by the cavities of the bread. The disengagement of steam from boiling water by platinum foil or any other solid substance, is likewise of the same nature. After a very short time this effect ceases, unless renewed by a fresh surface. The most natural explanation of these phenomena, is to refer them to some molecular action of the solid on the gas, probably of a mechanical nature, which lasts a very short time, when the solid acquires a "droit de domicile" in the liquid, and becomes perfectly inert. M. Legrand, who has made most correct experiments on the point of ebullition of saline solutions, remarks, that platinum possesses no power in equalizing ebullition after a few moments, when, according to him, all the air has been expelled from its surface; but on the contrary, zinc and iron will act as long as they are present in the liquid, which he attributes to their power of decomposing water.

Previously to showing the existence of the same action in bodies in a state of vapour or of fume, I will make a short digression with respect to the constitution of vapours in general.

The term vapour is commonly applied to bodies in three different conditions,—1st, that of temporary gas diffused in the atmosphere; 2nd, that of liquid particles mechanically suspended there; 3rd, that of solid particles suspended in like manner. To the two latter, to speak more correctly, may be applied the term of fumes. The first correspond to solution in a liquid, and the other two to that of suspension in the same. As examples of the first, we have the vapours of water while in an invisible state, and those of bromine, &c. Of the



second, water as in mists, fogs, &c. ; and of the third, the vapours of arsenic and of corrosive sublimate. Bodies in either of these conditions possess the faculty of assuming a definite crystalline form on becoming solid. The properties of the gaseous vapours are so well known, that it is unnecessary to dwell upon them here.

The second class, or the liquid globular vapours or fumes, which, as we have said, cause those accumulations known under the name of fogs, clouds, or mists, are those which I intend at present to examine, as they comprehend the theory of the fixation of the mercurial vapours in the Daguerreotype. It was formerly believed that vapour or mist was composed of minute spherules or globules of liquid water, and in Newton's works we find evidence that such was his opinion. According to another view, first advanced I believe by De Saussure, these vapours were composed of vesicles or very minute hubbles, exactly resembling, on a small scale, the common soap-bubble. This opinion has received the assent of Fresnel and Berzelius, and at present obtains general credence. The proofs on which it is considered to be founded, are principally the observations of De Saussure, who asserts that on high mountains, or in the clouds, he has been able to detect these air-vesicles with the naked eye, and has seen them burst as they came in contact with each other. Berzelius recommends the examination of the vapour of water over a dark surface, such as that of ink, with a lens of a short focus. He says, that vesicles may be detected in this manner, varying in size from  $\frac{1}{4500}$  to  $\frac{1}{2780}$ th of an inch, which occasionally burst as they touch each other. The suspension of clouds is also used as an argument in favour of the vesicular theory, as it is contended that liquid spherules would descend to the ground by their specific gravity in such situations. Fresnel indeed compares the globules to small balloons, which dilate or contract, according to the temperature of the air they contain.

A few days' stay at the convent of St. Bernard gave me an opportunity of repeating the observations on the clouds, as mentioned by De Saussure, which may be also made in this season on our London fogs. Globules of various sizes in these circumstances are frequently discerned by the naked eye floating in all directions. I have endeavoured to ascertain their vesicular structure, but have been unable to do so from direct observations. It is frequently a most difficult point, in microscopic investigation, to decide upon the existence of a thin transparent membrane. It is still more so to pronounce upon the vesicular or spherular structure of globules in constant agitation ; and I believe that if minute spherules and vesicles

could be mixed together, we do not possess any means at present of distinguishing them.

I have never been able to detect that appearance of bursting of the globules mentioned by De Saussure, but sometimes, when the agitation of the air is slight, two of the larger globules may be seen floating towards each other, and afterwards disappear suddenly, which may be explained, if we admit that it is caused by the union of the two spherules into one, which is too heavy to remain any longer in suspension, and whose rapid deposition conceals it from the sight.

There may be urged as objections to the vesicular theory, that if the pellicle become extremely thin, the vesicle would no longer be perceived any more than the apex of an air-bubble before bursting, or the central black spot of a system of Newton's coloured rings. It will be seen below that the globules of vapour possess the power of depositing themselves in a crystalline form, which requires a tranquil deposition of particles, such as could scarcely be deemed possible, if the air contained in each had to escape at the moment of its crystallization.

I have endeavoured to fix the globules of water on glass and other substances, so as to be enabled to submit them to microscopic inspection, but from their volatile nature and other causes have not succeeded. However, it is easy to do so with almost any other volatile substance; and I have examined several in this way without detecting the slightest appearance of a vesicular structure. Mercury is deposited under the form of globular particles, with a metallic lustre whose diameter is  $\frac{1}{300}$ th of a millimetre, in which I have never detected any internal cavity by the most careful examination\*. Flour

\* In order that others who may wish to verify these results may operate in the same conditions as myself, it is proper to state that the mercurial vapours were disengaged in a box, such as is used in the Daguerreotype process; and after the mercury had been raised to a temperature of about 90° centigrade, it was allowed to cool. Three experiments were made in this manner: in the two first the glass plate was placed four inches above the mercury, in the other it was eight inches distant. The appearance of the globules was the same in each case; if any difference existed in their size, those of the last experiment were rather larger. In another experiment, where a common Daguerreotype plate was substituted for one of glass, the appearance of the globules was in all respects the same. From the manner in which they are deposited, they appear to exert an influence over each other, as they are frequently found in groups of three or four, or more. Mr. Ross has stated on the part of Mr. Solly (Microscopical Society, December 1843), that these globules are deposited in hexagonal groups; but with preconceived ideas no doubt it would be very easy to form such shapes, as it would be to form triangles or any other simple geometrical figure, particularly when the illusions inseparable from catoptric microscopy

of sulphur is found to consist of solid globules, several of which adhere together; when acted upon by a gentle solvent, their external portion is dissolved, and there remains a regular octahedron. An interesting experiment may be made on the fumes of sal-ammoniac, which appear whenever muriatic acid and ammonia are brought together. Two small phials, each containing one of these substances, are covered by an inverted tumbler: above the surface of the acid are seen at a short distance the fumes of the salt, which at the end of a few hours are found to have condensed into a thin snowy pellicle, completely obturating the mouth of the bottle. This partition is so delicate, that the slightest agitation will cause it to fall into the liquid.

In all these cases it is found that the fumes possess the power of remaining suspended a much greater length of time than would be expected from the difference of their specific gravity with that of air, which is also the case with the fumes of other substances, and smoke in particular. This can only be accounted for by the continual state of agitation of the air, even within an enclosed space, and by the elasticity of the solid and liquid particles. In the case of solid particles this can be readily admitted, but with regard to liquid globules, there is probably some action similar to that which takes place on the impinging of solid elastic balls, which after becoming flattened rebound in virtue of their tendency to recover their original shape.

The causes which act in fixing different vapours and fumes are the same as those which determine the precipitation of solid particles in solution, such as for instance, sharp points of any kind, minute filaments, and more especially the existence of a crystalline particle to act as a nucleus. Non-conducting substances, as woollen cloth, the nap of a hat, the web of the spider, &c., are covered with aqueous globules when no rain has fallen, and when polished surfaces near present no such deposition.

Having now shown the existence of a crystalline power in vapours, we shall proceed to prove the influence of a force which disturbs this equilibrium in the same manner as in the saline solutions above mentioned. The friction of a solid body on glass will leave traces which are invisible until breathed upon.

are added to those of physiology. This tendency of the mind, of which a good account has been given by Müller in his *Elements of Physiology*, is so strong, that where groups of globules are concerned, I would always advise their being mapped down under the microscopic camera lucida, and put by for some time for future inspection. I shall have occasion to advert to this subject more fully hereafter.

Many bodies possess this property, but the mineral steatite, or soap-stone, produces the effect better than any other I know. A considerable degree of friction may be used over the traces thus produced by steatite, without affecting the appearance of the traces when breathed upon repeatedly. The glass may even be heated considerably without affecting them. By examining with the microscope the parts that have been traced upon by steatite, we are unable, any more than with the naked eye, to detect any material cause for the deposition of vapours in these places, as it probably depends upon the transparency of the mineral, which being so attenuated is unable to affect the rays of light. When the traces have been brought out by breathing upon them, they must be covered with another piece of glass, which impedes the evaporation of the water and allows them to be submitted to the microscope. The parts untouched by the steatite present the appearances that have been already mentioned. On the lines created by the mineral, the drops of water are differently disposed, their long diameters being parallel to the direction of the lines. These minute drops very much resemble the globules of gas deposited from a liquid, the only difference between the two consisting in the deviation from the globular form in the liquid traces, which evidently arises from the power which the water possesses of wetting glass.

It is evident, therefore, that the secondary cause of these images is a difference in the position of the minute drops of water, reflecting the light differently from the other drops, which are irregularly disposed on the other parts of the glass.

There exists another method of fixing vapours, which has been long known, and to which I believe attention was first directed by Prof. Draper. It consists in merely placing a body on a plain surface, such as that of a metallic speculum, or even of glass; after a short time it is found that simple contact, such as this, has caused some molecular action, as the spot occupied by the object will become apparent by breathing on it in the same way as with the images of steatite. This observation is the more interesting, as it serves as a connecting link between the effects of mechanical power and those caused by other agents.

The experiments of Mr. Hunt have shown the influence of heat in causing the fixation of vapours.

An image of this sort formed on glass by the breath, when examined under the microscope, presents exactly the same appearance as those formed by steatite. The same difficulty is experienced in bringing out, by mercurial vapours, the thermographic images on glass, as is found with the traces of

steatite, which possess but in a very slight degree the power of fixing mercurial vapours. It appears therefore that the power which water has of wetting glass, causes it to have a greater tendency to deposit than mercury, which does not wet glass. The cause of the production of thermographic images is evidently similar to that which causes the deposition of a solid body from a solution.

The fixation of the mercurial vapours in the Daguerreotype process, which has excited so much interest, and for which so many theories have been advanced, is but another example of the force which causes the deposition of solid and gaseous particles from a liquid, and which produces so many other effects. In this case the chemical rays of light act in the same manner as mechanical action and caloric in causing a certain molecular disturbance. By the discoveries of Moser, it is shown that these rays possess the power of acting upon almost any body, in such a manner as to render it capable of fixing the particles of various vapours. Thus simple minerals, glass, &c. may be made to fix the mercurial vapour.

It appears however that silver, gold, copper, &c., which form amalgams, or in other words, are capable of being wetted by mercury, possess this property in a greater degree than any other bodies which are incapable of being wetted by it; in the same way as we have seen that glass has the greatest power to fix the vapour of water. Admitting the truth of this theory of the Daguerreotype process, we are naturally led to inquire whether the same agent may not likewise cause the fixation of particles in a state of solution or of vapour, in the same manner as by simple mechanical action. After several unsatisfactory attempts, I finally succeeded in clearly proving this fact. The solution which shows the influence of light the most evidently, is that of the neutral chloride of gold. A few grains of this salt dissolved in an ounce of water, when exposed to the light, deposits minute crystals of a metallic appearance on that side of the glass nearest the light.

The action of light in causing the deposition of gaseous vapours may be shown by placing some iodine in a bottle closed with a glass stopper. After being exposed to the sunshine for several hours, minute black crystals will appear on the side nearest the light, which will change their position according to the side of the glass exposed. Another substance which shows this action still better, is camphor, a piece of which, merely covered with a glass shade, will give rise to a crystalline deposit, after an hour or two of exposure to light, and which presents the same phænomena as that of iodine. By a prolonged exposure these crystals become very abundant, and

are very beautiful\*. I have applied this property to the construction of an instrument for measuring the chemical rays of light. As the details respecting this would be foreign to our present subject, I will defer them to another occasion, and confine myself now to prove that these phænomena are independent of the deposits caused by radiation.

1st. The crystals are formed on the side exposed to the action of direct or diffused light.

2nd. They are not formed during the night, when the radiation from the earth is sufficient to cause the deposition of water.

3rd. Green glass, which retards photographic action, likewise impedes this deposit.

In an experiment which is now going on, a bottle of pale green common glass is exposed to the north, while another of white glass is placed in a southern aspect. The first became covered with minute crystals, in size averaging about a millimetre, which have remained stationary for a week; the second is covered with arborescent ramifications, which are daily increasing.

Several familiar, but hitherto unexplained phænomena, may in my opinion be easily accounted for by these molecular actions.

The formation of hail I consider to be an instance of an action precisely similar to that which causes the deposition of the solids of gaseous and liquid particles. If we admit the influence of this force on the globular vapours of water, it is not at all improbable that certain conditions may arise in nature when these vapours may be much more liable to this influence than we find them in our imperfect experiments. We have seen that a solution of sulphate of soda or water in a pure state may be brought by the abstraction of caloric to such a condition of unstable equilibrium, that the slightest perturbing cause will immediately reduce them to a solid form.

If we admit that the globules which form the clouds are capable of being placed in a similar condition, we have sufficient data to explain all the phænomena that occur in the production of hail. Any nucleus formed within a cloud in this state, would create around it a deposition of all the neighbouring particles; and the size of the hail-stones would be dependent upon the thickness of the cloud it had to traverse. In the storm at Ordenburg, in 1825, mentioned by Dr. Eversman, pyrites was found in the centre, and had acted like a nucleus

\* I am informed by a friend, that this action of camphor was mentioned twenty years since by Dr. Hope in his lectures, but I am not aware of anything having been published upon the subject.

round which the crystallization had taken place. Where the centre is not formed by a foreign body of this sort, it has frequently been mentioned that it consisted of an opaque nucleus of a spongy nature, like congealed snow, which may be easily accounted for. The succession of concentric layers would be caused by the passage of the particles through strata of liquid globules not all at the same temperature; and the radiated structure indicates a gradual increase of crystalline action proceeding from the centre. The temperature of the hail-stones, which has generally been found below the freezing-point, is a further corroboration of this view.

The formation of butter is likewise in all probability another instance of molecular action of the same nature. It is well known that after the cream has been agitated for a certain length of time, the globules suddenly coalesce, and by their union butter is produced. The sudden appearance of this product is the more remarkable, as it takes place at different temperatures, although more quickly at some than others, and not gradually, as might have been expected, which precludes the idea of its being owing to any caloric developed by friction. The most minute observations have been unable to show any material alteration in the appearance of the fatty globules at the moment before the butter is formed. Little doubt can be entertained of its being caused by some molecular action, or engendered in the globules by the continued agitation they have undergone.

Some of the most permanent gases likewise exhibit phenomena closely allied to the above, by their action on platinum and other metals. According to Dulong and Thenard, platinum foil newly beaten has the property of acting at the common temperature, on a mixture of hydrogen and oxygen; but after a few minutes' exposure to the air, it entirely loses that power, which may however be restored to it in a stronger degree than before by heating it in a covered crucible. If it be kept in a covered vessel, so as to exclude the air, it will retain the power without decrease for four-and-twenty hours.

Platinum filings, made with an ordinary sized file, have the same property immediately after their formation, and which they retain for above an hour. It has also been observed, that a hollow ball of platinum has the power of condensing and absorbing different gases, which are generally disengaged at a temperature below the boiling-point (Pouillet, *Elémens de Physique*, § 131). The action of the gases on platinum in all the above cases greatly resembles that of carbonic acid on glass, except that not merely simple lines, but the whole surface of the metal exerts its influence, and that the gases themselves are invisible.

XXI. *Note to Mr. Hennessy's Paper on the Connexion between the Rotation of the Earth and the Geological Changes of its Surface\**.

THE values of  $I_1$  and  $K$  must be altered, as some incorrect assumptions were made in obtaining them. This alteration will produce no changes in the general conclusions which have been arrived at.

The method for obtaining the moment of inertia of a solid of revolution contained in equation (3.), appears to have been inapplicable to the case of the internal spheroid of the earth from the nature of the expression for  $\rho$ . The expression for the earth's moment of inertia, which is used for obtaining the theoretical coefficients of precession and nutation, is, however, adapted to our purpose. In this case we have †,

$$I_1 = \frac{8A}{3\pi^3} \left(\frac{6p}{5}\right)^4 \left\{ \left(5\pi - \left(\frac{5}{6}\right)^3 \pi^3\right) \cos \frac{5}{6}\pi + \left(\frac{25}{12}\pi^2 - 6\right) \sin \frac{5}{6}\pi \right\},$$

$$K = \frac{10368}{1875} \left(\frac{\alpha - D}{\beta - D}\right)^4 \left\{ \left(5\pi - \frac{125}{216}\pi^3\right) \cot \frac{5}{6}\pi + \frac{25}{12}\pi^2 - 6 \right\}.$$

When this value of  $K$  is substituted in (19.), that resulting for  $P$  will evidently be less than what has been already found, and it will give an amount of denudation of the earth's surface still more within the limits of geological observations than that which has been previously obtained.

H. HENNESSY.

XXII. *Letter to Henry Lord Brougham, F.R.S., &c., containing Remarks on certain Statements in his Lives of Black, Watt and Cavendish. By the Rev. WILLIAM VERNON HARCOURT, F.R.S. &c.*

MY DEAR LORD,

IN a volume of biography which you have lately published, I perceive that you have reprinted your contribution to M. Arago's historical notice of Watt, in which the distinguished author attempted to transfer to the subject of his *éloge* the credit of a celebrated chemical discovery, hitherto by the common consent of chemists attributed to Cavendish.

Your personal challenge to myself would not have moved me to enter again on a question which I scarcely think open to dispute since the publication of the fac-similes of Cavendish's original notes of that discovery, in the Transactions of the British Association for the Advancement of Science, had I not

\* Phil. Mag. vol. xxvii. p. 376, November 1845.

† Airy's Tracts, Precession and Nutation, Art. 43.



observed, as it seems to me, other mistakes in this volume, on points of scientific history, which, venial as they are in one who cannot be supposed to have devoted much of his valuable time to these *umbratile studies*, are yet such as ought not to pass without some notice.

I must begin, however, my criticisms on your historical chemistry, by repeating the grounds on which I deemed it needful to controvert the statements of M. Arago respecting the discovery of the composition of water. "The *éloge* of Watt, delivered before the French Academy by one of its secretaries, and subjoined to the *Annuaire* for 1839, had just been published. It was blemished by statements which reflect unjustly on the character of one whose memory is cherished among us as a bright example of the union of modesty with science, of the purest love of truth with the highest faculties for its discovery, and the most eminent success in its attainment. Perceiving these statements to be founded in error, I took the earliest opportunity of rectifying them, at the meeting of the British Association which followed within two or three weeks after I became acquainted with them, rejoiced that I had it in my power, from the position in which, as President of that body, I had then the honour to be placed, to make the correction of the error as formal and public as its promulgation had been; and persuaded that M. Arago, as soon as he should be fully possessed of the facts, would consider it a duty which he owes both to the Academy and himself, to retract the suspicions which he had expressed\*."

Those who feel that a sense of justice is a material part of the character of illustrious men and illustrious bodies, are still "waiting," not "till your fellow champion," as you express it, "shall seal your adversary's doom," but till he makes the *amende honorable* by withdrawing, in explicit terms, imputations which since the lithographing of the Cavendish MSS. he must know to be unfounded. I am not content, my dear Lord, that you should either for your "colleague" or yourself, half retract and half retain those doubts which I perceive that you have *republished* in one part of your volume, whilst you *disclaim* them in another.

"I cannot easily suppose," you say, "that M. Arago ever intended, and I know that I never myself intended, to insinuate in the slightest degree a suspicion of Mr. Cavendish having borrowed from Mr. Watt." Certainly, as regards yourself at least, no declaration can be more explicit than this. But what then, give me leave to ask, is the significance of the following words in your *now republished* appendix to M. Arago's

\* Report of the British Association for 1839, p. 22.

éloge? “Whether or not Mr. Cavendish had heard of Mr. Watt’s theory previous to drawing his conclusions, appears more doubtful: the supposition that he had so heard rests on the improbability of Sir C. Blagden and many others knowing what Mr. Watt had done and not communicating it to Mr. Cavendish, and on the omission of any assertion in Mr. Cavendish’s paper, even in the part written by Sir C. Blagden with the view of claiming priority as against M. Lavoisier, that Mr. Cavendish had drawn his conclusion before April 1783. Mr. Watt’s theory was well known among the members of the Society some months before Mr. Cavendish’s statement appears to have been reduced into writing, and eight months before it was presented to the Society. That the first letter of April 1783 was for some time,—two months as appears from the papers of Mr. Watt,—in the hands of Sir Joseph Banks and other members of the Society during the preceding spring, is certain from the statements in the note to p. 330; and that Sir C. Blagden, the Secretary, should not have seen it seems impossible, for Sir Joseph Banks must have delivered it to him at the time when it was intended to be read at one of the Society’s meetings (Phil. Trans. p. 330, note); and as the letter itself remains among the Society’s records in the same volume with the paper into which the greater part of it was introduced, it must have been in the custody of Sir C. Blagden. It is equally difficult to suppose that the person who wrote the remarkable passage already referred to respecting Mr. Cavendish’s conclusions having been communicated to M. Lavoisier, should not have mentioned to Mr. Cavendish that Mr. Watt had drawn the same conclusion in the spring of 1783, that is, in April at the latest; for the conclusions are identical, with the single difference that Mr. Cavendish calls dephlogisticated air water deprived of its phlogiston, and Mr. Watt says that water is composed of dephlogisticated air and phlogiston.”—(Life of Watt, pp. 396–398.)

To what does all this argument tend?—Would it lead any one to guess that you mean to acquit Cavendish of plagiarism, or that “you have yourself,” as you elsewhere affirm, “always been convinced that Mr. Watt had, *unknown to Cavendish*, anticipated his great discovery?” Allowing a certain interval of time and place, I should not wonder at your having forgotten or laid aside your doubts whether Cavendish, with the connivance of Blagden, had not purloined the *conclusions* of Watt; but I have never before known an instance of so deliberate a disavowal of a suspicion contemporaneous and in juxtaposition with its no less deliberate reiteration.

Your *reprinting now* these old doubts is the more unaccountable, not only because they consist so ill with your profession of belief in the good faith of Cavendish, and are indeed a mere trifling after that point has been satisfactorily established, but because I have corrected the *particular* error out of which this tissue of suspicions was spun; and you are now apprised that the Secretary of the Royal Society at that time was *Mr. Maty*, and not, as you persist in taking for granted, Cavendish's friend *Dr. Blagden*, who did not enter on the office till *May 1784*. "So that," as I told you in the Appendix to my address to the British Association, "he is not liable to the suspicion intimated by Lord Brougham, of having shown Watt's letter to Cavendish, nor to the reproach which M. Arago casts upon him, of not speaking the whole truth respecting the precise date at which Watt's opinions were made known in London."

The confidence which you place, with so much simplicity, in the innocence of M. Arago's "*intentions*," contrasts strangely with the disposition you have shown to suspect Cavendish and Blagden: for M. Arago does not, like yourself, "just hint a fault," but retorts in good set terms on the English philosophers the imputation which Blagden had cast on Lavoisier, "That he had told the truth, but not the whole truth." "This is a heavy charge," says your illustrious colleague; "let us see whether *all who took part in this affair* are not liable to the same reproach;"—and then in a style of pointed irony, into the spirit of which I should have thought you apt enough to enter, he proceeds to fix the charge on all the parties concerned. I believe I have given no more than the plain meaning of these clever sarcasms when I said, "The Secretary of the Academy has not confined himself to taking from Cavendish the honour of this discovery, but has in fact imputed to him the claiming a discovery which he borrowed from another; of inducing the Secretary of the Royal Society to aid in the fraud, and even causing the very Printers of the Transactions to antedate the presentation copies of his paper."

The real truth is, that M. Arago having, when in England, heard but one side of the story, was persuaded of the *insincerity* of Cavendish. If he is now disabused of this persuasion, I hope he will choose another method of withdrawing what he wrote under such an impression than that which you have framed for him in the following protest. "As a strange notion seems to pervade this paper that every thing depends on the character of Cavendish, it may be as well to repeat the following disclaimer, already very distinctly made, of all intention to cast the slightest doubt upon that great man's per-

fect good faith in the whole affair, I never having supposed that he borrowed from Mr. Watt, though M. Arago, Professor Robison and Sir H. Davy, as well as myself, have always thought that Mr. Watt had, unknown to him, anticipated his great discovery."

Of the deceased philosophers, whose names are here pressed into this service, I shall presently have occasion to speak; but let me first venture to answer for M. Arago, that if *he* has "read the fac-similes" of Cavendish's notes, you will not find him at the same loss as yourself to discover the *inferences* of the experimental philosopher in *the steps of his investigation*; he will not join you in propounding, "that in all Cavendish's diaries and notes of his experiments, not an intimation occurs of the composition of water having been *inferred* by him earlier than Mr. Watt's paper of spring 1783."

Those celebrated experiments of 1781, which pass with chemists for a model of a well-combined train of analytical and synthetical research, you imagine to have been without object or *inference*, till an imperfect attempt to repeat them had the good luck to be reasoned upon by Watt in 1783. You appear to think that the manner in which the great facts of experimental philosophy are ascertained is by one man's stumbling on the *proofs*, and another some time after hitting on the *conclusion*. If it be so, I believe that you would have been as capable of interpreting such experiments, once made, as James Watt himself; and could you have been at hand when Cavendish, in July 1781, completed the discovery of those facts which prove the composition of water, he need not have waited so long to learn what to *infer* from them: I doubt not but that you would at once have drawn the *inference* for him, established the *theory*, and become for ever memorable as the true discoverer; you would, in your own amended phrase, "*unknown to him, have anticipated his great discovery.*"

But I own I do not suspect your "colleague" of these peculiar views. Once satisfied that Cavendish spoke truth when he said that all the experiments on this subject published in his paper were made by him in the summer of 1781, he will no longer doubt to whom the discovery of this important fact is due; once convinced that the experiments were communicated to Priestley, and that the attempt to repeat them was made in consequence of that communication; once aware that the repetition was abortive because made with a *wrong* gas, that neither the phlogiston nor the inflammable air of Priestley and Watt were convertible terms for *hydrogen*, and that their notions of the change of water into air and air into water had no reference to that particular gas, but first to

nitrogen, and afterwards to a mixture of gases, the chief of which was *carbonic oxide*--M. Arago will keep you "waiting" long before he rejoins you in the advocacy of any part of the supposed claims of your client, or thanks you for classing him with yourself as still cherishing the conviction that "Mr. Watt had, unknown to Cavendish, anticipated his great discovery."

That which renders the self-devotion of this knight-errantry complete, is the singular fact that you are fighting for Watt *against himself*. I had formerly come to the conclusion that he never thought of claiming the discovery in the sense which you suppose, nor in any other respect than as regards the theory of the extrication of heat and light from the combining gases; and a circumstance has lately been pointed out to me by a friend, which establishes this conclusion.

The edition of Robison's *Mechanical Philosophy*, published by Sir D. Brewster, was revised by Watt himself. In that revision we find him by no means indifferent to his own just fame. Writing to the Editor he says, "I have carefully perused my late excellent friend Dr. Robison's articles, 'Steam and Steam-Engines,' in the *Encyclopædia Britannica*, and have made remarks upon them in such places, where either from the want of proper information, or from too great a reliance on the powers of his extraordinary memory at a period when it probably had been weakened by a long state of acute pain, and by the remedies to which he was obliged to have recourse, he had been led into mistakes in regard to facts, and also in some places where his deductions have appeared to me to be erroneous. Dr. R. qualifies me as 'the pupil and intimate friend of Dr. Black:' he afterwards, in his dedication to me of Dr. Black's *Lectures upon Chemistry*, goes the length of supposing me to have professed to owe my improvements upon the steam-engine to the instructions and information I had received from that gentleman, which was certainly a misapprehension; as though I always felt and acknowledged my obligations to him for the information I had received from his conversation, and particularly for the knowledge of the doctrine of latent heat, I never did nor could consider my improvements as originating in those communications. He is also mistaken in his assertion, p. 8 of the preface to the above work, that I had attended two courses of the Doctor's lectures; for unfortunately for me, the necessary avocations of my business prevented me from attending his or any other lectures at College." Mr. Watt then quotes from these lectures a passage in which Black is made to say, "My own fortunate observation of what happens in the formation and condensation of steam, had

suggested to my friend Mr. Watt his improvements in the steam-engine," and remarks, "it is very painful to me to controvert any assertion or opinion of my revered friend; yet in the present case I find it necessary to say that he appears to have fallen into an error\*."

But in revising the article on Steam, and making remarks on those places in which Dr. Robison had been led into mistakes, Watt makes no remark on the following very decisive passage:—"We know that in vital or atmospheric air there is not only a prodigious quantity of fire which is not in the vapour of water, but that it also contains light, or the cause of light, in a combined state. This is fully evinced by the great discovery of Mr. Cavendish of the composition of water: there we are taught that water, and consequently its vapour, consists of air from which the light and greatest part of the fire have been separated; and the subsequent discoveries of the celebrated Lavoisier show that almost all the condensable gases with which we are acquainted, consist either of airs which have lost much of their fire, and perhaps light too, or of matters in which we have no evidence of light and fire being combined in this manner."

Thus you see, that jealous as Watt appears of any undue share in his own discoveries being attributed even to his "revered friend" Dr. Black, he allows "the great discovery of the composition of water" to be assigned to Cavendish without reclaiming the least participation in it for himself.

These extracts entirely relieve his memory from any suspicion of his having been a party to the erroneous statements contained either in the article 'Water' in the first edition of the *Encyclopædia Britannica*, to which you have referred, or in the posthumous lectures of Black. Nor do I hold Black responsible for the fabulous history of this discovery given in the latter work. It is well known that that unambitious man left behind him no MSS. of any account, and that the Lectures published under his name were chiefly composed out of the reminiscences of the able but incorrect Editor. Robison, on historical points, was a very inaccurate writer; and to his inaccuracy I attribute the extraordinary string of errors on this subject which I have formerly pointed out.

It is from the latter work that you seem to have taken your

\* I conceive Watt to mean that the *facts* known to him respecting the condensation of steam, independent of Black's *theoretical explanation* of them, were the foundation of his improvements; and I am bound therefore, on his own showing, to allow that M. Arago has done right in not placing the merit of Watt in the study and application of *abstract philosophical principles*, so much as in ingenuity of mechanical contrivance and the happy adaptation of well-observed facts.

supposed facts; and you have in consequence entirely misstated the nature of Cavendish's experiments. Where, allow me to ask, do you find in his paper, or his notes, any such matter as this? "He then weighed accurately the air of both kinds, which he exposed to the stream of electricity; and he afterwards weighed the liquid formed by the combustion: he found that the two weights corresponded with great accuracy" (Life of Cavendish, p. 433): and again, "Water equal to the weight of the two gases taken together remained as the produce of the combustion." *Cavendish made no such experiments*; as you will find whenever you take the trouble to read either the documents themselves\*, or my account of them†. I have already stated that *this* method of determining the composition of water, which is attended with great practical difficulties, was tried indeed at a later time by the French philosophers with such accuracy as it admits of, but that Cavendish, with his usual sagacity, had taken an easier and more certain road: having mastered beyond any of his cotemporaries the analysis of gases, and possessed himself of their specific properties, he was enabled to substitute the method of volume for that of weight; he found that about two volumes of hydrogen and one of oxygen, when burnt together, entirely disappeared without loss of weight, and that *pure water* was the result. To draw from these premises the obvious conclusion, there was no need to weigh or compare the weight of the airs, and the water that lined the glass after combustion; and he did *not* compare it. Lavoisier followed in his steps: and should you ever read his papers, you will find that *he* too in the first instance contented himself with *deducing* the equality of the weights as a *corollary* from experiments of the same kind as those of Cavendish.

Had you happened to consult the *second* edition of the *Encyclopædia Britannica* as well as the *first*, you would have found it purged both of these, and some other of Robison's historical mistakes. You would have found all that you have referred to *omitted*; and in the article 'Chemistry,' compiled under the revision of friends and connections of Watt, the following account substituted in its place. "In the year 1781, Mr. Cavendish proved that water is not a simple element, but that it is composed of pure or vital air, and inflammable air." "In the mean time the French chemists were not idle; the celebrated Lavoisier, in conjunction with some of his philosophical friends, confirmed by the most decisive experiments the truth of *Mr. Cavendish's discovery of the composition of water*, which

\* Phil. Trans. vol. lxxiv. Experiments on Air, by H. Cavendish, Esq. Report of the British Association for 1839, autograph notes of experiments.

† Report of the British Association for 1839, pp. 35, 36.

was now received and adopted by almost every chemist." A detailed account is then given of Cavendish's experiments; and it is added, "*these experiments were made in 1781, and they are undoubtedly conclusive of the composition of water.*" It would appear that Mr. Watt entertained the same ideas on this subject. When he was informed by Dr. Priestley of the result of *these* experiments, he observed, Let us consider what obviously happens in the deflagration of oxygen and hydrogen gases," &c. "Thus it appears that Mr. Watt had a just view of the composition of water, and of the nature of the process by which its component parts pass to a liquid state from that of an elastic fluid."

In this account the ideas entertained by Mr. Watt obtain more notice perhaps than would have been accorded to them by an indifferent historian; but the statement of the discovery is correct, as is also that of the view which Watt took of the subject, if we confine the assertion of the justness of his ideas to his apprehension of the relation of Cavendish's discovery to certain theories of light and heat; for of the material base of water he had certainly no just conception when he wrote the letter which is quoted above. I have shown that his views in 1783 and 1784 were founded on several suppositions:—1st, that Priestley had converted water into *atmospheric air*; 2nd, that he had obtained a weight of water equal to the weight of a mixture of oxygen with *the gases extricated by heat from moist charcoal*; 3rd, that he had shown good reason to believe that carbon, combined in a certain proportion with oxygen, constitutes water. All these suppositions agreed perfectly with the opinions which Watt really expressed, that water was formed of *dephlogisticated air* and *phlogiston*; but no one of them is consistent with the opinions attributed to him by an erroneous translation of his words, that water is formed by the combination of oxygen with *hydrogen gas*\*.

From your mention of Sir H. Davy's sentiments without a quotation, I suppose that he, like Dr. Henry, has been among the number of those on whose attention this untenable claim has been privately pressed; all I know of Davy's opinion on the subject is from his *published works*, in which he has spoken, like other chemists, of the composition of water and of nitric acid as "*the two grand discoveries of Mr. Cavendish.*" But in referring to the name of this much-honoured and regretted friend, I must take the opportunity of noticing what I think a serious error in your impressions respecting one point in his personal character. You begin your sketch of his life with these words: "Sir H. Davy being now removed beyond the reach of

\* See Report of the British Association for 1839, pp. 24, 25.



such feelings, as he ought always to have been above their influence, that may be said without offence of which he so disliked the mention; he had the honour of raising himself to the highest place among the chemical philosophers of the age, emerging by his merit alone from an obscure condition." A simple anecdote may suffice to set his feelings on this subject in a more favourable light. When Davy was exhibiting to myself and three others the discoveries which he had then recently made relative to his safety-lamp, and when those present, among whom were the Hanoverian minister and the late Lord Lonsdale, were highly admiring the beauty of his experiments, with still higher admiration I heard him reply, "Yes, I have some reason to be proud of them, for my experiments on flame were first made *with a tallow candle in an apothecary's shop.*"

In these slight sketches which you have given us of the history of men eminent in science, there is one other scientific subject besides the discovery of the composition of water, on which you appear to have bestowed some consideration, namely,—the first discoveries of the gases. Here Cavendish is still out of favour with you. You pluck another feather from his wing; and having made a present of the discovery of water to Mr. Watt, dispense that of hydrogen gas to Dr. Black.

"The nature of hydrogen," you say, "was perfectly known to him, and both its qualities of being inflammable, and of being so much lighter than atmospheric air; for as early as 1766 he invented the air-balloon, showing a party of his friends the ascent of a bladder filled with inflammable air: Mr. Cavendish only more precisely ascertained its specific gravity, and showed, what Black could not have been ignorant of, that it is the same from whatever substance it is obtained\*."

You ought to have recollected, when again contravening the received opinion of chemists †, your own remarks on the supposed omission of Cavendish to state exactly the time when he had communicated to Priestley his experiments on the composition of water. "Dans une addition de Blagden faite avec le consentement de Cavendish, on donne aux expériences de ce dernier le date de l'été de 1781. On cite une communication de [à] Priestley, *sans en préciser l'époque, sans parler de conclusions, sans même dire quand ces conclusions se pré-*

\* Life of Black, p. 383.

† The *received* account of the discovery of hydrogen is this:—"Its combustible quality is described in the works of Boyle and Hales, of Boerhaave and Stahl; but it was not till the year 1766 that its properties were particularly ascertained, and the difference between it and atmospheric air pointed out by Mr. Cavendish."—Encycl. Brit., Art. Chemistry, 1810.

sentaient à l'esprit de Cavendish. Ceci doit être regardé comme une très grosse omission (*a most material omission*\*)” Nothing indeed can be more unfounded than this animadversion. In the passage to which you refer, the words of Cavendish are these:—“All the foregoing experiments on the explosion of inflammable air with common and dephlogisticated airs, except those which relate to the cause of the acid found in the water, were made by me in the summer of 1781, and mentioned by me to Dr. Priestley, *who in consequence of it made some experiments of the same kind, as he relates in a paper printed in a preceding volume of the Transactions.*” Now you need only have referred to the volume of the Transactions which Cavendish quotes, to have found the “*epoch*” which you wanted. Priestley’s paper was printed in March 1783; and therefore Cavendish’s communication of his “conclusive” experiments was *anterior* to Watt’s speculations in April, as well as to Lavoisier’s experiments in June of the same year.

But though this “*most material*,” or in M. Arago’s translation, this “*grosse*” omission turns out to be *none*, you ought, I repeat it, to have remembered your own demand for preciseness of dates, when you ascribed to Black a prior knowledge of the distinguishing properties of hydrogen gas. In proof that Black knew before Cavendish that this gas is “*so much lighter than atmospheric air*,” you allege, that “as early as 1766 he invented the air-balloon, showing a party of his friends the ascent of a bladder filled with inflammable air: Mr. Cavendish only more precisely ascertained its specific gravity.”

As early as 1766?—Are you not aware that Cavendish’s paper on factitious airs was published in this year? Is it not a “*most material omission*” that you have forgotten to “*préciser l’époque*” of Black’s experiment with the balloon, so as to show whether it was before or after the publication of Cavendish’s paper? Professor Leslie tells the story of the balloon somewhat differently from you. “The late most ingenious and accurate Mr. Cavendish, in 1766, found, by a most nice observation, this fluid to be at least seven times lighter than atmospheric air. It *therefore* occurred to Dr. Black of Edinbro’, that a very thin bag filled with hydrogen gas would rise to the ceiling of a room. He provided accordingly the allantois of a calf, with a view of showing at a public lecture such a curious experiment before his numerous auditors; but owing to some unforeseen accident or imperfection it chanced to fail, and that celebrated Professor, whose infirm state of health and indolent temper more than once allowed the finest dis-

\* Historical note, Life of Watt, p. 383.

coveries when almost within his reach to escape his penetration, did not attempt to repeat the exhibition, or seek to pursue the project any further."

If you are dissatisfied with Leslie's version of your anecdote, let me refer you to other authorities. In one of those articles of the *Encyclopædia Britannica* which are stated to have been composed or revised by Professor Miller, Dr. Muirhead, and Sir David Brewster, the circumstance is thus narrated:—"In the year 1766 Mr. Henry Cavendish ascertained the weight and other properties of this gas, determining it to be at least seven times lighter than atmospheric air. *Soon after which* it occurred to Dr. Black that perhaps a thin bag filled with hydrogen gas might be buoyed up by the common atmosphere."

I hope I have now illustrated sufficiently the value of the canon of criticism which you have laid down for these delicate inquiries,—that nothing is so necessary as to "*préciser l'époque.*"

"Cavendish," you say, "only more precisely ascertained the specific gravity of inflammable air; and showed, *what Black could not be ignorant of, that it is the same from whatever substance it is obtained.*" Now, in the first place, *inflammable air* is *not* the same from whatever substance it is obtained. This was the error into which Priestley fell when he attempted to repeat the experiments by which Cavendish had discovered the composition of water; this was the error under which Watt laboured till after the publication of Cavendish's paper in 1784, and which nullified the researches of the one and the speculations of the other. But supposing you to mean "that inflammable air is the same, *whether obtained from zinc or iron,*" why do you say that Black could not be ignorant of *that*? How do you think he was to know it? How did Cavendish know it? He tells you that he learnt it by having ascertained by experiment that the specific gravity of the gas from either material was the same. Had Black ascertained this? Had he any test whatever by which he could know that these gases were the same?

But Cavendish "*only more precisely ascertained the specific gravity of inflammable air.*" If any person conversant in the history of pneumatic discoveries were to be asked to enumerate the most important of the early advances in that branch of science, he would certainly name—1st, the discovery of the weight of the air by Galileo; 2nd, the discoveries of its law of compression, and of the factitious gases, by Boyle; 3rd, the theory of the fixation of gases by chemical attraction, propounded by Newton; 4th, the discovery of specific and elective affinities in one of those gases, by Black; 5th, the

discovery of the difference of specific gravity in several gases, by Cavendish.

You do not distribute their honours to any of these great discoverers with a severe attention to matter of fact; but I must do you the justice to own that you preserve a principle of equity in your adjudications. You omit, it is true, to dwell upon, or even to mention, the main point of novelty in the researches of Black; but then you give to Black the discoveries of Boyle and Cavendish, and make it up to Cavendish by allowing him a slice of the merit which belongs to Galileo.

For Cavendish you say, "He carried his mathematical habits into the laboratory; and not satisfied with showing the other qualities which make it clear that these two æriform substances are each *sui generis*, and the same from whatever substances, by whatever processes they are obtained,—not satisfied with the mere fact that one of them is heavier, and the other much lighter than atmospheric air," (a previous acquaintance with all which facts you have taken care to ascribe to Dr. Black) "he inquired into the precise numerical relation of their specific gravities with one another and with common air, and *first showed an example of weighing permanently elastic fluids*: unless indeed Torricelli may be said before him to have shown the relative weight of a column of air and a column of mercury, or the common pump to have long ago compared in this respect air with water. It is however sufficiently clear that neither of these experiments gave the relative measure of one air with another; nor indeed could they be said to compare common air with either mercury or water, although they certainly showed the relative specific gravity of the two bodies, taking air for the middle term or common measure of their weights."

What a strange qualification of a still stranger assertion! If instead of this confusion of specific gravities with equiponderant columns, ending with the grave suggestion, that "the relative specific gravities of water and mercury" might have been taken by the intermediation of "air," you had said that philosophers have attempted, from the relative heights of the barometer at different elevations, to calculate the mean specific gravity of the atmosphere\*, there would have been mean-

\* The following quotation will show the nature of these calculations (Dan. Bernoulli Joh. Fil. Hydrodynamica, *Argentorati*, 1738. Sect. 10. 16. p. 209):—"Patet exinde quid censendum sit de illa methodo qua in Anglia aliquando usos esse recenset D. Du Hamel, in Hist. Acad. Sc. Paris. ad indagandam rationem inter gravitates specificas aëris et mercurii: observata nimirum altitudine mercurii in loco humiliori, tum etiam in altiori, gravitates specificas in aëre et mercurio statuerunt, ut erat differentia

ing at least in the qualification; but then what an assertion to hazard! considering the great number of experiments extant for the *direct* determination of the weight of air compared with that of water, first instituted by Galileo, and then repeated successively by Descartes, Mersenne, Boyle, Hook, Newton, Cotes, and lastly by Hawksbee, whose determination was assumed and quoted by Cavendish himself for the purpose of comparing the specific gravity of common air with those of the factitious gases,—it is a strong instance of the kind of equity for which I have given you credit, that you should have allotted to Cavendish the merit of having “*first showed an example of weighing permanently elastic fluids.*” Even Descartes allowed that Galileo’s “method of weighing

altitudinum mercurii in barometro ad altitudinem inter locos observationum interceptam; etiamsi aër ejusdem densitatis ponatur ab imo observationis loco ad alterum usque, non licet tamen inde judicare de ejus gravitate specifica ratione mercurii. Quicquid ab experimento colligere licet hoc solum est:—

“Consideremus scilicet integram crustam aëream terram ambientem atque inter ambo observationis loca interceptam, et erit pondus istius crustæ ad superficiem terræ ut pondus columnæ mercurialis qualis in barometro descendit ad basin ejus; manifesta hæc sunt ex eo quod summa basium A et B sustinent quidem summam ponderum quæ habent columnæ aëreæ A C et B D, neque tamen quævis basis premitur suæ columnæ pondere seorsim, et quod idem resectis columnis A g et B h intelligi debet de columnis g C et h D, diaphragmatis in g et h positis, incumbentibus. Igitur experimentum non tam gravitatem specificam aëris in quo factum est indicat, quam omnis aëris terræ proximi gravitatem specificam mediam determinat; prior admodum variabilis est, altera procul dubio constanter eadem fere permanet.

“Faciamus computum *gravitatis specificæ istius mediæ aëris* omnis qui terram ambit. Multis vero experimentis, quæ in diversis locis parum supra mare elevatis sumpta fuerunt, id constat, elevationi 66 pedum proxime descensum respondere unius lineæ in barometro. Sequitur inde quod aëris gravitas specifica media ratione mercurii sit, ut altitudo unius lineæ ad altitudinem 66 pedum, *i. e.* ut ut 1 ad 9504, ergo posita gravitate specifica mercurii = 1, erit gravitas specifica *media* aëris = 0.00105. Notabile est profecto tantam esse hanc gravitatem mediam aëris: certus enim sum vel maxime sævienti hic locorum frigore aëris gravitatem specificam vixdum tantam esse quantam nunc exhibuimus pro statu medio omnis aëris terram ambientis: at sub æquatore multo erit minor, et omnibus recte pensis non crediderim *gravitatem mediam* aëris qui inter utramque latitudinem 60 gr. continetur, ultra 0.000090 excurrere; quo posito erit *gravitas media* aëris ab utroque polo ad 30 gradus terram cingentis, quod spatium paullo plus quam octavam totius terræ superficiæ efficit partem, = 0.000210, quæ dupla est aëris hic locorum densissimi: sub ipso autem polo, præsertim antarctico, admodum gravior erit aër, et fortasse aqua vix decies levior, cum est frigidissimus atque densissimus.

“32. Et quia aëris mediocriter densi gravitas specifica est ad gravitatem specificam merc. ut 1 ad 11000, ipsaque altitudo media merc. in barometro pro locis parum a superficie maris elevatis sit  $2\frac{1}{3}$  ped. Paris. erit altitudo aëris homogenii mediocriter densi 25666 pedum.”

the air was not amiss\*;" and the experiments of the great Italian philosopher, which laid the original foundation of all our knowledge of elastic fluids, ought not to have been entirely forgotten by any one who appreciates duly those capital discoveries by which the ideas of men are fixed and a new order of facts is ascertained.

To Black, on the other hand, with like even-handed justice, you ascribe a knowledge of the lightness of hydrogen and the heaviness of carbonic gas, which you have no ground for suspecting him to have possessed. Experiments, indeed, had been made with a view of ascertaining such points, and your assertion, that "Cavendish first set the example of weighing permanently elastic gases," is so far from the truth, that the factitious gases themselves had been weighed both by Hawksbee and Hales. Hales weighed the "air of tartar," which consists of a mixture of carburetted hydrogen and carbonic gases in a bladder, and then filling it with common air compared the weights †; Hawksbee ascertained accurately the specific gravity of air that had passed through tubes filled with iron wires, and heated red in the fire, which consisted partly of carbonic acid and partly of nitrogen ‡. But these mixed

\* "Sa façon de peser l'air n'est pas mauvaise, si tant est que la pesanteur en soit si notable qu'on la puisse apercevoir par ce moyen; mais j'en doute." (*Œuvres de Descartes*, tom. vii. p. 440.) Thus Descartes wrote to Mersenne in 1638. In 1642 he repeated the experiment himself by a method far less susceptible of accuracy, and obtained a result much further from the truth, which satisfied him however, "que la poids de l'air est sensible en cette façon." (*Œuvres*, tom. viii. p. 567.) Dr. Whewell has taken notice (*History of Mechanics*, p. 66) that "in a letter of the date of 1631 he (Descartes) explains the suspension of mercury in a tube closed at the top by the pressure of the column of air reaching to the clouds." In this letter the atmosphere is compared to a pack of wool, the filaments of which are all heavy, and press on each other from the clouds to the earth, being only kept apart by the æther which plays between them, "ce qui fait un grand pesanteur"—expressions which at first sight might lead to the idea that he had anticipated the theory of the elevation of the barometric column; but it is evident from many subsequent letters of Descartes, that he had no correct conception of the statical pressure of fluids, and was therefore incapable of reasoning justly on this subject. The tube in which the mercury was suspended in the case in question, was a *straight tube without a bason*: he tried to account for the phænomenon of its suspension on his principle of *circular movement in a plenum*, by supposing that the mercury, before it could quit the tube, must effect the circle of motions required to bring down from the sky a current of æther to supply the vacuum left at the top of the tube by the descent of the quicksilver; and presuming the column of air which it had to lift to be as heavy as itself, he concluded that no such circular motion in the chain of matter could take place. It is possible however that this representation of the atmosphere as a heavy column may have conduced to suggest the more correct views of the subject afterwards adopted.

† Veg. Status, p. 185.

‡ Phil. Trans., No. 328, p. 199.

gases approached too nearly to common air in that respect to enable the experimenters to establish a distinction. An attempt too had been made by Greenwood, a Professor of Mathematics at Cambridge in New England, to ascertain the specific gravity of the deleterious air in a well, which was doubtless chiefly carbonic gas; but the method employed by him was not sufficiently delicate to show a difference of density. Such was the state of knowledge, or rather ignorance, on this subject previous to the experiments of Cavendish. We have not the least reason to believe that any one had observed the different weights of the different kinds of air. Dr. Mayow\* indeed about a century before had supposed his "*nitro-igneous aura*," to the combinations of which he ascribed the phænomena of acidification, combustion and vitality, to be *heavier* than the residual air from which it is separated in those processes; and this opinion, which proved to be correct, he entertained so distinctly, as to represent the specific lightness of the vitiated air, after it had served its purpose of sustaining life, as a provision of nature for freeing us from a noxious atmosphere. But he had no better ground for entertaining such an opinion than his observation of the movements of animals which he had confined in a close vessel, and which appeared in his experiments to seek for a less suffocating air in the lower part of the receiver, whilst they avoided the upper.

Such loose surmises as these detract nothing from the great experimental discovery of Cavendish, the importance of which cannot be better expressed than in the words of an eminent chemist and chemical historian †: "It can scarcely be said that pneumatic chemistry was properly begun till Mr. Cavendish published his valuable paper on Carbonic Acid and Hydrogen Gas, in the year 1766." On the fruits of this discovery, in the hands of its author and of all succeeding chemists, and its consequences to the study of gaseous substances and their combinations, I need not dwell. It is enough to remark, that the ascertainment of *this physical* difference in the gases was the first *conclusive* proof of a *plurality* of elastic fluids.

Another point of no small consequence to pneumatic chemistry was first made out in this paper. From the earliest discovery of factitious airs, it had been observed that a considerable portion of several of these disappeared after they had been generated, though there had been no change of temperature or pressure. The usual statement of this phænomenon

\* *De parte aëria igneaque Spir. Nitri.*

† Dr. J. Thomson's Biographical Account of Priestley, Ann. Phil., vol. i. p. 91.

was, that the elasticity of the air had been *destroyed*. Dr. Hales, dissatisfied with so loose an explanation, accounted for the loss, which in the case of nitrous acid *he* first observed, after the following manner:—"When fresh air is let into the receiver, whose included air is impregnated with the fumes arising from the mixture of compound *aquafortis*, or spirit of nitre, and Whitstable pyrites, mentioned in the following experiment, then the air in the receiver turns very red and turbid, and much air is absorbed after several repeated admissions. When fresh air is thus admitted into the glasses full of sulphureous, though clear, air, a good many particles of the fresh air must needs be reduced by the sulphureous ones from an elastic to a fixed state, as in the effervescences of other liquors. Therefore the rising of the water in the glass vessel does not seem to be wholly owing to the rebating of the air's elasticity in some degree, but rather to the reduction of it from an elastic to a fixed state, which is further probable from hence, viz. that the whole quantity of air admitted at several times is equal, or nearly equal, to the quantity of sulphureous air A. Z., so that *both airs are at the same time contained within the space A. Z.*"

In this important observation, which was subsequently turned to such good account by Priestley and Cavendish, Hales gave the true theory of the loss of volume which occurs by the admission of common air to nitrous gas; but the variable, and apparently capricious, loss of elasticity which he remarked in other gases he could not explain. "Though a good part of the air," he says, "which rises from *fluids* seems to have existed in an elastic state in those fluids, yet the air which arises from *solid* bodies, either by the force of fire or effervescence, does not seem to arise only from the interstices of those bodies, but principally from the most fixed parts of them. For since the airs which are raised by the same acid spirit from a vast variety of substances have very different degrees of permanency, as was shown in Exp. 10, No. 3, 4, 5, 6, and in Exp. 11, No. 6, 7, 8, 9, 10 of experiments on stones, hence it is probable that these airs do not arise from latent interstices of the dissolved stones, &c., but from the solid fixed particles of them; and since the whole of some of these newly-generated airs does in a few days lose its elasticity, it should seem hence probable, that whatever air arises from the spirit in the effervescence is not permanently elastic, or else that in the rotation of some stones it is thrown off into a more permanently elastic state than from others."

The cause of this loss of volume was first explained in Cavendish's paper: he proved by experiment, that carbonic acid



is condensed over water, but not over mercury. You indeed tell us that Black "*found this gas incondensable,*" but he has nowhere told us as much himself; and you might with more safety have presumed the contrary; the true statement being, that he and his predecessors had found it *condensable*, and that Cavendish found the conditions under which it is *not condensed*.

In the same spirit of liberality you take "*the capital discovery,*" that the air of the atmosphere is not the only air permanently elastic, from its ancient owners, to appropriate it to Black, and expend much learned pains in setting forth the originality and importance of the "*doctrine*" which you ascribe to him. "The great step," you say, "was now made, that the air of the atmosphere is not the only permanently elastic body, but that others exist, having perfectly different qualities from atmospheric air, and capable of losing their elasticity by entering into chemical union with solid and with liquid substances, from which, being afterwards separated, they regain the elastic or aëriform state." . . . . "In order to estimate the importance of this discovery, and at the same time to show how entirely it attends the whole face of chemical science, and how completely the doctrine was original, we must now examine the state of science which philosophers had previously attained to. It has often been remarked, that no great discovery was ever made at once, except perhaps that of logarithms: all have been preceded by steps which conducted the discoverer's predecessors nearly, though not quite, to the same point. Some may perhaps think that Black's discovery of fixed air affords no second exception to this rule; for it is said that Van Helmont, who flourished at the end of the sixteenth and beginning of the seventeenth century, had observed its evolution during fermentation, and gave it the name of *gas sylvestre*, spirit from wood, remarking that it caused the phenomena of the Grotto del Cane near Naples; but though he, as well as others, had observed an aëriform substance to be evolved in fermentation and in effervescence, there is no reason for affirming that they considered it as differing from atmospheric air, except by having absorbed or become mixed with various exhalations or impurities. Accordingly a century later than Van Helmont, Hales, who made more experiments upon air than any of the old chemists, adopts the commonly received opinion, that all elastic fluids were only different combinations of the atmospheric air with various exhalations or impurities: and this was the universal opinion upon the subject, both of philosophers and the vulgar." . . . . "It is now fit that we see in what manner the subject was treated by

scientific men at the period immediately preceding Black's discoveries. The article 'Air,' in the French *Encyclopédie*, was published in 1751, and written by D'Alembert himself. It is, as might be expected, able, clear and elaborate. He assumes the substance of the atmosphere to be alone entitled to the name of air, and to be the foundation of all other permanently elastic *bodies*. When D'Alembert wrote this article, he gave the doctrine then universally received, that all the other kinds of air were only impure, and that this fluid alone was permanently elastic, all other vapours being only like steam, temporarily aëriform. Once the truth was made known, that there are other gases in nature, only careful observation was required to find them out\*."

After all this, should I venture to affirm that you have post-dated our knowledge of permanently elastic gases, other than the atmosphere, by about a hundred years, were I to suggest that in this case also the old story is the true one, and that Priestley has correctly recorded the real historical fact when he said, "Mr. Boyle, I believe, was the first who discovered that what we call fixed air, and also inflammable air, are really elastic fluids capable of being exhibited in a state unmixed with common air," were I to add that the existence of various elastic fluids was generally recognised by the philosophers of Europe, and particularly by those whom you have quoted as instances to the contrary, during the century which preceded Black's essay, as distinctly, and more distinctly, than by Black himself,—I know not what you would think of me. Nevertheless, since this is a passage in the history of science which deserves to be told with a strict regard to dry matter of fact, I must beg you to listen with patience to an account of it certainly very different from your own.

It was in December 1659 that Boyle published his "New Physico-mechanical Experiments," among which is to be found a description of two of those gases separable from fixed bodies, which he subsequently denominated *factitious airs*. The high interest which may be justly attached to all the circumstances of discoveries so important as this, induces me to give the details of it in the words of the author.

"Contenting myself," he says, "to have mentioned our author's (Kircher's) experiment as a plausible, though not demonstrative, proof that water may be transmuted into air, we will pass on to mention, in the third place, another experiment which we tried in order to the same inquiry. We took a clear glass bubble, capable of containing by guess about three ounces of water, with a neck somewhat long and wide of

\* Life of Black, pp. 331-36.

a cylindrical form: this we filled with oil of vitriol and fair water, of each almost a like quantity, and casting in half a dozen small nails we stopped the mouth of the glass, which was top-full of liquor, with a flat piece of *dia palma* provided for the purpose, that, accommodating itself to the surface of the water, the air might be exquisitely excluded; and speedily inverting the phial, we put the neck of it into a small wide-mouthed glass that stood ready with more of the same liquor to receive it. As soon as the neck had reached the bottom of the liquor it was dipped into, there appeared at the upper part, which was before the bottom of the phial, a bubble of about the bigness of a pea, which seemed rather to consist of small and recent bubbles produced by the action of the dissolving liquor upon the iron, than any parcel of the external air that might be suspected to have got in upon the inversion of the glass, especially since we gave time to those little particles of air which were carried down with the nails into the liquor to fly up again. But whence the first bubble was produced is not so material to our experiment, in regard it was so small; for soon after we perceived the bubbles produced by the action of the menstruum upon the metal, ascending copiously to the bubble named, and breaking, did soon exceedingly increase it, and by degrees depress the water lower and lower, till at length the substance contained in these bubbles possessed the whole cavity of the glass phial, and almost of its neck too, reaching much lower in the neck than the surface of the ambient liquor wherewith the open-mouthed glass was by this means almost replenished. And because it might be suspected that the depression of the liquor might proceed from the agitation whereinto the exhaling and imprisoned steams were put by that heat which is wont to result from the action of corrosive salt upon metals, we suffered both the phial and the open-mouthed glass to remain as they were in a window for three or four days and nights together; but looking upon them several times during that while, as well as at the expiration of it, the whole cavity of the glass bubble and most of its neck seemed to be possessed by air, since by its spring it was able for so long to hinder the expelled and ambient liquor from regaining its former place. And it was remarkable that just before we took the glass bubble out of the other glass, upon the application of a warm hand to the convex part of the bubble, the imprisoned substance readily dilated itself like air, and broke through the liquor in divers bubbles succeeding one another.

“Having also another time tried the like experiment with a small phial and with nails dissolved in *aquafortis*, we found

nothing incongruous to what we have now delivered. And this circumstance was observed, that the newly-generated steams did not only possess almost all the whole cavity of the glass, but divers times without the assistance of heat of my hand did break away in large bubbles through the ambient liquor into the open air: so that the experiments with corrosive liquors seemed manifestly to prove, though not that air may be generated out of water, yet that in general air may be generated anew.

“Lastly, to the foregoing arguments from experience, we might easily subjoin the authority of Aristotle and of his followers the schools, who are known to have taught that air and water, being symbolizing elements in the quality of moisture, are *easily transmutable into each other*\*; but we shall rather to the foregoing argument add *this*, drawn from reason—that if, as Leucippus, Democritus, Epicurus and others, followed by divers modern naturalists, have taught, the difference of bodies proceeds but from the various magnitudes, figures, motions, and textures of the small parts they consist of (all the qualities that make them differ being deducible from thence), there appears no reason why the minute parts of water, and other bodies, may not be so agitated and connected as to deserve the name of air; for if we allow the Cartesian hypothesis, according to which the air may consist of any terrene or aqueous corpuscles, provided they be kept swimming in the interfluent celestial matter, it is obvious that air may be as often generated, as terrestrial particles, minute enough to be carried up and down by the celestial matter, ascend into the atmosphere. And if we will have the air to be a congeries of little slender springs, it seems not impossible, though it be difficult, that the small parts of divers bodies may, by a lucky concurrence of causes, be so connected as to constitute such little springs, since water in the plants it nourisheth is usually contrived into springy bodies, and even the bare altered position and connexion of the parts of a body may suffice to give it a spring that it had not before, as may be seen in a thin and flexible plate of silver, into which, by some strokes of a hammer, you may give a spring; and by only heating it red-hot, you may make it again as flexible as before.

“These, my Lord, are some of the considerations at present occurring to my thoughts, by which it may be made probable that air may be generated anew.”

\* This I presume is the *hypothesis, doctrine, or theory* which Cavendish was suspected of deriving from Watt.

In a subsequent part of the same treatise, Boyle adds an account of another discovery of a similar kind. "I took," he says (exp. 42), "whole pieces of red coral, and cast them into as much spirit of vinegar as sufficed to swim about an inch over them: these substances I made use of that the ebullition upon the solution might not be too great, and that the operation might last the longer." It gave but few bubbles, till the receiver under which it was placed was exhausted; "then the menstruum appeared to boil in the glass like a seething-pot. To avoid suspicion, that these proceeded not from the action of the *menstruum* upon the coral, but from the sudden emersion of those many little parcels of air that are wont to be dispersed in liquors, we conveyed over distilled vinegar alone into the *receiver*, and kept it awhile there to free it from the bubbles, which were but very small, before ever we put the coral into it. The former experiment was another time tried in another small receiver with coral grossly powdered, and the success was much alike."

Of the two gases thus first obtained and separated, he observed some time afterwards that the one was inflammable\*,

\* "Having provided a saline spirit, which by the uncommon way of preparation was made exceeding sharp and piercing, we put into a phial, capable of containing three or four ounces of water, a convenient quantity of filings of steel, which were not such as are commonly sold in shops to chemists and apothecaries, those being usually not free enough from rust, but such as I had awhile before caused to be purposely filed off from a piece of good steel. This metalline powder being moistened in the phial with a little of the menstruum, was afterwards drenched with more, whereupon the mixture grew very hot, and belched up copious and very stinking fumes, which, whether they consisted altogether of the volatile sulphur of the Mars, or of metalline steams participating of a sulphureous nature, and joined with the saline exhalations of the menstruum, is not necessary here to be discussed. But whencesoever this stinking smoke proceeded, so inflammable it was, that upon the approach of a lighted candle to it, it would readily enough take fire, and burn with a bluish and somewhat greenish flame at the mouth of the phial for a good while together; and that though with little light, yet with more strength than one would easily suspect. This flaming phial therefore was conveyed to a *receiver*, which he who managed the pump affirmed that about six exsuctions would exhaust. And the receiver being well cemented on, upon the first suck the flame suddenly appeared four or five times as great as before, which I ascribed to this, that upon withdrawing of the air, and consequently the weakening of its pressure, great store of bubbles were produced in the menstruum, which breaking, could not but supply the neck of the phial with store of inflammable steams, which as we thought took not fire without some noise. Upon the second exsuction of the air, the flame blazed out as before, and so it likewise did upon the third exsuction; but after that it went out, nor could we rekindle any fire by hastily removing the receiver: only we found that there remained such a disposition in the smoke to inflammability, that holding a lighted candle to it a flame was quickly rekindled."—*New Experiments touching the Relation between Flame and Air*, 1671.

and the other liable, in part, to lose its elasticity\*; he extended his experiments on the generation of "factitious airs" to a variety of materials, multiplying them to such an extent that one of Cotes's hydrostatical lectures is filled with the repetition of them; he remarked the condensability of muriatic acid gas†, and the orange colour of nitrous acid gas ‡; and extricated from red lead, by the burning-glass, the gas§, which Priestley afterwards having obtained by the same method, was led by reasoning from the manner in which red lead is

\* "*Experiment 8.*—A mercurial gauge having been put into a conical glass whose bottom was covered with beaten coral, some spirit of vinegar was poured in, and then the glass stopper closing the neck exactly: on the working of the menstruum on the coral, store of bubbles were for a good while produced, which successively broke in the cavity of the vessel; and their accession compressed the confined air in the closed leg of the gauge three divisions, which I guessed to amount to about the third part of the extent it had before; but some hours after the compression made by this newly-generated air grew manifestly fainter, and the imprisoned gauge-air drove down the mercury again, till it was depressed within one division of its first station; so that in this operation there seemed to have been a double compressive power exercised, the one transient by the brisk agitation of vapours, the other durable from the aerial or springy particles either produced or extricated by the action of the spirit of vinegar on the coral."—*Phil. Trans.* 1675.

† "May 26, 1676.—Sal-ammoniac was put into a receiver with a sufficient quantity of oil of vitriol. Then the air being exhausted, the salt was put into the oil, whereupon a great ebullition presently followed, and the mercury in the gauge showed a good quantity of air to be generated; but this by the same gauge soon after appeared to be destroyed again. The experiment was repeated, and both the production and destruction were slower than before. It was confirmed by these trials that some artificial airs may be destroyed, but why this destruction happens sometimes sooner and sometimes slower, may perhaps seem worthy of further inquiry."—*2nd cont. Phys. Mech. Expts.* 1676.

‡ "We put an ounce of such strong spirit of nitre as is above mentioned into a moderately large bolt head, furnished with a proportionable stem, over the orifice of which we strongly tied the neck of a thin bladder, out of which most part of the air had been expressed, and into which we had conveyed a small phial with a little highly rectified spirit of wine. Then this phial, that was before closed with a cork, being unstopped without taking off the bladder, a small quantity, by guess not a spoonful, of the alcohol of wine was made to run down into the spirit of nitre, where it presently produced a great commotion, and blew up the bladder as far as it would well stretch, filling also the stem and cavity of the glass with very red fumes, which presently after forced their way into the open air, in which they continued for a good while to ascend in the form of an orange-coloured smoke."—*New Experiments about Explosions*, 1672.

§ "September 4, 1678.—I exposed one ounce of minium in an open glass to the sunbeams, concentrated by a burning-glass, and found that it had lost three-fourths of a grain of its weight, though much of the minium had not been touched by the solar rays. May 29.—Repeated the same experiment, in a light glass phial sealed hermetically and exactly weighed,

manufactured, to identify with the oxygen of the atmosphere\*, just as Mayow identified with it the gas from saltpetre.

In giving to the gases which he discovered the title of "factitious *airs*," Boyle did not confound them with *common air*. The extracts which I have given sufficiently show that he used the word *air* generically, in the sense which he assigns to it in the following passage:—"If I were to allow acids to be one principle, it should be only in some such metaphysical sense as that wherein *air* is said to be one body, though it consist of the associated effluvioms of a multitude of corpuscles of very different natures that agree in very little, save in their being minute enough to concur in the composition of a fluid aggregate consisting of flying parts‡."

It would indeed be a great mistake in the history of science, to suppose that the notion of the air being a simple element prevailed among philosophers down to the days of Black. From the time of that remarkable revolution in the scientific mind of Europe which attended the revival of the mechanico-corpuseular philosophy, when the phænomena of nature were accounted for no longer by forms and qualities, but by the sizes and motions, the cohesions and disjunctions of the particles of bodies, the atmosphere came at once to be conceived of as a miscellaneous aggregate of the molecules of a variety of heavy substances thrown into an elastic state, or floating in an active medium of a still finer and more divided consistence.

"Tout corps invisible et impalpable," says Descartes, "se nomme *air*, à savoir en sa plus ample signification‡." "By air," says Dr. Wallis, "I find Mr. Hobbs would sometimes have us understand a pure æther, 'aërem ab omni terræ aquæque effluviis purum, qualis putatur esse æther,' to which I suppose answers the *materia subtilis* of Descartes, and M. Hugen's 'more subtile matter' than air: on the other hand, M. Hugen here by air seems to understand that feculent matter arising from those the earth's and water's effluvia, which

and the loss of weight came to  $\frac{1}{3}$ rd part of a grain. May 30.—I endeavoured to burn the same minium again, but such plenty of air was produced, that the glass broke into a hundred pieces."

\* "At the same time that I got the air above-mentioned from mercurius calcinatus and the red precipitate, I had got the same kind from red lead or minium. In this process that part of the minium on which the focus of the lens had fallen turned yellow. The experiment with red lead confirmed me more in my suspicion, that the merc. calcinatus must have got the property of yielding this kind of air from the atmosphere, the process by which that preparation and this of red lead is made being similar."—*Priestley's Experiments on Air*, vol. ii. p. 111.

† Reflections on the Hypothesis of Alkali and Acidum, ch. iv. 1676.

‡ *Œuvres*, tom. vii. p. 237.

are intermingled with this subtle matter. *We mean by air the aggregate of both these, or whatever else makes up that heterogeneous fluid wherein we breathe, commonly called air, the purer part of which is Mr. Hobbs's air, and the feculent of it is M. Hugens's air\*.*"

It is curious to trace the fortunes of this *materia subtilis*, from the naked condition in which it was first ushered into notice, to the figure which it now makes in the speculations of science.

Descartes was undoubtedly the first who formed the idea of a liquid medium grosser than heat, but more subtle than air, extending from the heavenly bodies to the earth, filling the aërial interstices with a continuous series of molecular globules, pervading the pores of glass, diamond, and the densest substances, without interruption, and propagating, by communication of impulses from one molecule to another, the movement, or rather the *pressure without locomotion*, simple and compound, which he considered as constituting light† and colours.

This was a grand conception, for which the philosophy of optics is under an obligation to the inventor greater perhaps than has been confessed. But the range of Descartes's views in physics was too limited to admit of his turning such a conception to its full account. He seems to have had no idea of intermittent or elastic forces, and did not even endow either

\* Extract of Letters from Dr. J. Wallis to the publisher, 1672, Phil. Trans. No. 91.

† Dr. Whewell takes Descartes's "hypothesis concerning light" to have been, "that it consists of small particles emitted by the luminous body," and considers this as "the first form of the emission theory" (Phys. Optics, ch. x.); and so the theory of the Dioptrics seems to have been understood by some of Descartes's cotemporaries; but he explains himself otherwise in his letters. "Je vous prie de considérer que ces petits globes dont j'ai parlé ne sont point des corps qui exhalent et qui s'écoulent des astres jusques à nous; mais que ce sont des parcelles imperceptibles de cette matière que V. R. appelle elle-même céleste, qui occupent tous les intervalles que les parties des corps transparents laissent entre elles, et qui ne sont autrement appuyées les unes sur les autres que le vin de cette cuve que j'ai pris pour exemple en la page 6 de ma Dioptrique, où l'on peut voir que le vin qui est en C tend vers B, et qu'il n'empêche point pour cela que celui qui est en E ne tend vers A, et que chacune de ces parties tend à descendre vers plusieurs divers endroits, quoiqu'elle ne se puisse mouvoir que vers un seul en même temps. Or j'ai souvent averti que par la lumière je n'entendois pas tant le mouvement, que cette inclination ou propension que ces petits corps ont à se mouvoir; et que ce que je disois du mouvement, pour être plus aisément entendu, se devoit rapporter à cette propension; d'où il est manifeste que, selon moi, l'on ne doit entendre autre chose par les couleurs que les différentes variétés qui arrivent en ces propensions." (*Œuvres*, tom. vii. p. 193). "J'admire que vous alléguiez les pages 4 et 5 afin de prouver que



his filaments of air, or his æthereal globules interposed between them, with attractive or repulsive powers.

The genius of Hook, so comprehensive of clear physical notions, soon lent to this luminiferous æther the mechanical attribute which it needed, and added the notion of *vibratory pulses*,—a notion which was instantly reduced by Newton to the form most competent to account for the phænomena\*, and on which Huygens founded, and Young with his illustrious coadjutors have gone far to finish, the mathematical fabric of the undulatory theory of light, as it is now commonly received.

So necessary indeed to any account of the phænomena of light and colours did the admission of such a medium appear, that Wallis, who not only rejected the use which Huygens and others proposed to make of it in explaining the extraordinary height at which mercury, purged of air, may be suspended in a tube, but denied it the properties of elasticity and weight, nevertheless did not scruple to say, “That there is in our air a body more subtle than the fumes and vapours mixed with it in our lower region *is very certain*: but whether that subtle body be, as Dr. Garden seems to suppose, much heavier than our common air, I much doubt, and rather think it is not, not having hitherto had any cogent experiment either to prove it heavy or elastic; but it may, for aught I know, be void as well of weight as spring, and what is found of either in our common air may be attributed to the other mixtures in it †.”

le mouvement des corps lumineux ne peut passer jusques à nos yeux, qu'il n'y passe quelque chose de matériel qui sorte de ces corps; car je ne fais en ces deux pages qu'expliquer la comparaison d'un aveugle, laquelle j'ai principalement apportée pour faire voir en quelle sorte *le mouvement peut passer sans le mobile*; et je ne crois pas que vous pensiez lorsque cet aveugle touche son chien de son bâton qu'il faille que ce chien passe tous le long de son bâton jusque à sa main, afin qu'il en sent les mouvements. Mais afin que je vous reponds *in formâ*, quand vous dites que le mouvement n'est jamais sans le mobile, *distinguo*; car il ne peut véritablement être sans quelque corps, mais il peut bien être transmis d'un corps en un autre, et ainsi passer des corps lumineux vers nos yeux par l'entremise d'un tiers, à savoir, comme je dis en la page 4, par l'entremise de l'air et des autres corps transparents, ou comme j'explique plus distinctement en la page 6, par l'entremise d'une matière fort subtile qui remplit les pores de ces corps et s'étend depuis les astres jusques à nous” (p. 240).

\* Phil. Trans., No. 88, p. 5088. An. 1672. “The most free and natural application of this hypothesis I take to be this: That the agitated parts of bodies, according to their several sizes, figures, and motions, do excite vibrations in the æther” &c.

† Phil. Trans. No. 171, p. 1002.

[To be continued.]

XXIII. *On a Proposition relating to the Theory of Equations.* By JAMES COCKLE, M.A., of Trinity College, Cambridge; of the Middle Temple, Special Pleader\*.

1. **L**ET  $x$  be the root of the general equation of the  $n$ th degree, and

$$y = \Lambda^I x^{\lambda'} + \Lambda^{II} x^{\lambda''} + \Lambda^{III} x^{\lambda'''} + \Lambda^{IV} x^{\lambda^{IV}}; \dots \quad (a.)$$

also let  ${}_m Y$  be composed of symmetric functions of, and be homogeneous and of the  $m$ th degree with respect to  $y$ ; then, if  $n > 2$ ,  ${}_2 Y$  may be reduced to the form

$$(a'_1 \Lambda^I + a''_1 \Lambda^{II} + b')^2 + (a''_2 \Lambda^{II} + b'')^2, \dots \quad (b.)$$

where  $b'$  and  $b''$  are not both zero.

2. For, let

$$\Lambda^{III} x_n^{\lambda'''} + \Lambda^{IV} x_n^{\lambda^{IV}} = l' x_n^{\lambda'} + l'' x_n^{\lambda''}, \dots \quad (c.)$$

then if  $y_r = (\Lambda^I + l') x_r^{\lambda'} + (\Lambda^{II} + l'') x_r^{\lambda''} + l_r, \dots \quad (d.)$

$$l_n = 0. \dots \quad (e.)$$

Now  ${}_2 Y$  is to be reduced, by means of (d.), to the form (b.), independently of  $\Lambda$ , or, what is the same thing, of  $\Lambda + l$ ; but†

$${}_2 Y = (b.) + [l_1 \dots l_{n-1}]^2, \dots \quad (f.)$$

[ ]<sup>m</sup> denoting a homogeneous function of the enclosed quantities of the  $m$ th degree. And, if  $n - 1 > 1$ ,

$$[l_1 \dots l_{n-1}]^2 = 0 \dots \quad (g.)$$

may be satisfied without making the  $l$ 's zero.

3. Following a notation similar to that used in my last paper‡, let  $(p, q)$  represent the coefficient of  $\Lambda^{(p)} \Lambda^{(q)}$  in the development of

$$t p_2 - s p_1^2 = {}_2 Y = 0, \dots \quad (h.)$$

$p_2$  and  $p_1$  being respectively the coefficients of the third and second terms of the transformed equation in  $y$ ; then, if (h.) be reducible to the form (b.), we have

$$\dots + \dots + b' \pm \sqrt{-1} \cdot b'' = 0; \dots \quad (i.)$$

and both the values of the above expression can only vanish when  $b' = 0 = b''$ . Substitute for  $b'$  and  $b''$ , equate each expression to zero, and eliminate  $\frac{\Lambda^{III}}{\Lambda^{IV}}$  between the two; then we

have  $(1, 3)(2, 4) - (1, 4)(2, 3) = 0, \dots \quad (j.)$

where, for instance,

$$(1, 3) = t \sum (x_1^{\lambda'} x_2^{\lambda''}) - 2s \sum (x_1^{\lambda'}) \cdot \sum (x_1^{\lambda''}); \dots \quad (k.)$$

\* Communicated by the Author.

† For the process, see par. 3 of the place which I have before cited, at the first line of p. 126 of vol. xxvii. of the Phil. Mag. S. 3.

‡ Phil. Mag. S. 3. vol. xxvii. p. 292.

so that, on developing, we shall have on writing  $\lambda' . \lambda''$  for  $\Sigma (x^{\lambda'}) . \Sigma (x^{\lambda''})$ , &c.,

$$0 = (t-2s) \times \{ \lambda' . \lambda^{iv} . (\lambda'' + \lambda''') + \lambda'' . \lambda''' . (\lambda' + \lambda^{iv}) - \lambda' . \lambda''' . (\lambda'' + \lambda^{iv}) - \lambda'' . \lambda^{iv} . (\lambda' + \lambda''') \} + t \{ (\lambda' + \lambda''') . (\lambda'' + \lambda^{iv}) - (\lambda' + \lambda^{iv}) . (\lambda'' + \lambda''') \}. \quad (1.)$$

Let  $t = 2n$ , and  $s = n - 1$ , then, if  $n < 3$ , the last equation is identically true, but not in any other case. The method of the two first paragraphs, consequently, detects every case of failure; the last-mentioned instance of which is connected with the fact that, implicitly at least, every expression of the form (a.) contains in its right-hand side a term free from  $x$  which, with the above values of  $t$  and  $s$ , vanishes from  ${}_2Y$ . These values are those which occur in exterminating the 2nd, 3rd, and  $r$ th terms of an equation.

4. If, in the case of  $n=2$ ,  $t=4$ , and  $s=1$ , we reject in (g.) the solution  $l_1=0$ , we are conducted to

$$(x_2^{\lambda'} - x_1^{\lambda'})^2 (x_2^{\lambda''} - x_1^{\lambda''}) = 0, \quad . . . \quad (m.)$$

having multiplied by the coefficient of  $\Lambda^{1/2}$  before commencing our operations. This agrees with what we have inferred from (1.).

5. It seems to follow from this, that biquadratics can be reduced to a binomial, and equations of the fifth degree to a trinomial form, by an expression for  $y$  consisting of four terms, determinable by one linear\*, one quadratic, and one cubic equation.

6. At p. 384 of the 26th vol. of this work, I have only alluded to the equation (3.), which, for cubics, conducts to the reducing equation

$$\xi^2 + \xi \Sigma \left( \frac{\phi'_1}{\phi_1} \right) + \frac{\phi'_1 \phi'_2}{\phi_1 \phi_2} = 0; \quad . . . \quad (3.)'$$

and to a similar one for biquadratics; but if we discuss the equation  $\phi \{ (\Lambda x^\lambda + M x^\mu)^{-1} \} = 0, . . . \quad (3.)''$

it will be found that, though in appearance more complicated, it is in reality simpler than the former, inasmuch as the case of  $\lambda=0$  is not excluded; and if  $\lambda=0$  and  $\mu=1$ , we have the form actually taken by the reducing equation in my solution of a perfect cubic at p. 248 of vol. ii. of the Cambridge Mathematical Journal.

Devereux Court, Temple Bar,  
December 29, 1845.

JAMES COCKLE, Jun.

\* The 'base' equations are linear, as will be seen on referring to my definition at note † of p. 126 of this (27th) vol. If the roots of the trinomial equation of the fifth degree are given by the expression  $b \psi(a)$ , then  $\psi$  is contained in the solutions of the functional equation  $\psi^2(a) - a = 0$ .

XXIV. *On Fresnel's Theory of Double Refraction.* By R. MOON, M.A., *Fellow of Queen's College Cambridge, and of the Cambridge Philosophical Society.*

[Continued from vol. xxvii. p. 559.]

**A**FTER proving, as he imagines, the existence of the three axes of elasticity, Fresnel enters into the most elaborate calculations as to the motion of the originally disturbed particle, and then proceeds to discuss the laws according to which the disturbance is transmitted from it to the rest of the medium. His labours in this respect are perfectly futile. The motion of the original particle, which is of the most simple character, is altogether different from what he supposes; and as to the laws according to which the disturbance is communicated from it to the rest of the medium, *no disturbance whatever can be propagated.*

As to the first point, Sir John Herschel proceeds:—"Suppose now any molecule set in vibration," in a plane passing through the centre of the surface of elasticity, "then at any period of its motion it will not be urged directly to its point of rest; but obliquely so that it will not describe a straight line, but will circulate in a curve more or less complicated; its motion however will always be resolvable into two vibratory rectilinear ones at right angles to each other, one parallel to the greatest, and the other to the least diameter of the section," which diameters it is shown, and this incontestably, are at right angles to each other. "Each of these vibratory motions will, by the laws of motion, be performed independently of the other; and therefore the motion propagated through the crystal will affect every molecule of it in the same way as if two separate and independent vibrations (at right angles, as above) were propagated through it with different velocities."

It is perfectly true that "the motion of the particle will always be resolvable into two vibratory rectilinear ones at right angles to each other, one parallel to the greatest, and the other to the least diameter of the section." But it is not true, as Fresnel quietly assumes, that the motion will be the same as if two separate disturbances were communicated, one in the direction of the greatest, and the other of the least diameter of the section. The distinction between the two cases is very palpable. We may resolve the actual force on the particle into two, one parallel to the greatest and the other to the least diameter of the section; and so the motions of the particle parallel to those lines may be determined; but these motions

respectively are not the same as if we calculate the effect of a disturbance communicated in the direction of the greatest, and then of another communicated in the direction of the least diameter. In the latter case, if  $u$  and  $v$  be the co-ordinates of the particle in the plane of the section, respectively parallel and perpendicular to the greatest diameter, the equation of motion parallel to that line is

$$\frac{d^2 u}{dt^2} = A u. \quad . . . . . (1.)$$

In the former, if  $\alpha \beta \gamma$  be the inclinations of the greatest diameter to the axes of elasticity, we have

$$\frac{d^2 u}{dt^2} = a^2 x \cos \alpha + b^2 y \cos \beta + c^2 z \cos \gamma; \quad . . (2.)$$

but if  $r$  be the radius vector of the particle,

$$u = r \cos \theta = x \cos \alpha + y \cos \beta + z \cos \gamma,$$

from which it is obvious that equations (1.) and (2.) can never be identical. This single circumstance would alone be sufficient to condemn the whole theory; I mention it however chiefly to show the gross fallacies which have been unhesitatingly received into it.

As to the second point which I have asserted, that *no disturbance whatever will be propagated* from the originally disturbed particle, a circumstance which if true must scatter the whole theory to the winds, I must say I approach the discussion of it with considerable pain, when I reflect that a result so immediately and incontrovertibly flowing from Fresnel's assumptions should so long have been overlooked or disregarded; and this when the theory has for years been subjected to the scrutiny of the ablest philosophers of this and of other countries.

Assuming Fresnel's proof of the axes of elasticity to be genuine, we get the following equations for determining the motion of the disturbed particle:—

$$\frac{d^2 x}{dt^2} = - a^2 x,$$

$$\frac{d^2 y}{dt^2} = - b^2 y,$$

$$\frac{d^2 z}{dt^2} = - c^2 z,$$

from which we obtain

$$\begin{aligned}x &= A \cos (a t + B), \\y &= A_1 \cos (b t + B_1), \\z &= A_{11} \cos (c t + B_{11}),\end{aligned}$$

where  $A, A_1, A_{11}, B, B_1, B_{11}$  are constants to be determined from the initial circumstances of the motion of the particle. From these equations, coupled with the fact which Fresnel assumes in his demonstration of the axes of elasticity, viz. that the change of position of the surrounding particles from the state of rest does not affect the forces upon the disturbed particle, we gather, that (1) without some special interposition of providence directed to each individual particle, *it would never move at all, whatever might be the state, whether of rest or motion, of the other particles around it*; and (2) that *once in motion, it would vibrate for ever without the least reference to or influence upon the other particles*. In my former paper, I said that Fresnel was driven to make an assumption as to the velocity of propagation, which rested only on the analogy of a case most widely differing from that under consideration. I now show that it is futile to talk of the velocity of propagation, when on his own showing *no wave whatever can be propagated*.

I purpose, in a future paper, to consider Fresnel's expressions for the intensity of the reflected and refracted rays when polarized light is incident on a surface.

Liverpool, December 3, 1845.

XXV. *Reply to some Remarks contained in Prof. Young's recent paper "On the Evaluation of the Sums of Neutral Series."* By R. MOON, M.A., Fellow of Queen's College, Cambridge, and of the Cambridge Philosophical Society\*.

**I**N a paper published in this Journal some months ago, upon the symbols  $\sin \infty$  and  $\cos \infty$ , I entered upon the discussion of the value of the series  $1 - 1 + 1 - 1 + \&c.$  continued to infinity, which I then showed to be 1 or 0 indifferently, in opposition to the commonly received opinion, which would make it equal to  $\frac{1}{2}$ . Prof. Young appears to be partly of my opinion in this respect, but seems to think I have made a mistake in supposing this to hold in all cases; for he appears to be of the opinion, that when the above series is considered as the limit of the converging series  $1 - x + x^2 + \&c.$ , where  $x$

\* Communicated by the Author.

is less than 1, it is still equal to  $\frac{1}{2}$ . I am not sorry to have the opportunity afforded me of expressing more fully my views upon this point.

It is somewhat difficult to conceive how, by considering the series  $1 - 1 + 1 - \&c.$  as the limit of another series, or by considering it in any other point of view, we can make its value different from what it is. *If it be* the limit of the series  $1 - x + x^2 - \&c.$ , where  $x$  is less than 1, and if moreover the limit of this last is  $\frac{1}{2}$ , it follows incontrovertibly that  $1 - 1 + 1 - \&c.$  must in all cases  $= \frac{1}{2}$ . The mistake here arises from calling  $1 - 1 + 1 - \&c.$  the limit of  $1 - x + x^2 - \&c.$ , where  $x$  is less than 1. It is no such thing. It is indeed the value which that series assumes when the limiting value is given to the variable; but it does not thence follow, nor is it the fact, that the one series is the limit of the other. We might expect the case to be otherwise, but it is not. Prof. Young himself admits, that without exception

$$1 - x + x^2 + \&c. (-x)^n = \frac{1}{1+x} + \frac{(-x)^{n+1}}{1+x}; \dots (a.)$$

and this holding always will hold in the limit when  $x = 1$ , which gives us, when  $n$  is infinite,

$$1 - 1 + 1 - \&c. = \frac{1}{2} \pm \frac{1}{2} = 1 \text{ or } 0 \text{ indifferently.}$$

From the same original equation we likewise deduce this other, that when  $x$  is not greater than 1,

$$1 - x + x^2 - \&c. = \frac{1}{1+x},$$

*except* in the limit: whence it follows that the series  $1 - x + x^2 - \&c.$ , where  $x$  is not greater than 1, approaches  $\frac{1}{2}$  as its limit. Now this is not more incontrovertible than that  $1 - 1 + 1 - \&c.$  is equal to 1 or 0, from which it is evident that the series  $1 - x + x^2 - \&c.$ , where  $x$  is less than 1, does *not* approximate to the series  $1 - 1 + 1 - \&c.$  as its limit; for the limit of a quantity or ratio is that quantity or ratio to which it continually approximates, and from which, although it never actually reaches it, its difference can be made less than any assignable quantity. It is perfectly true then that the limit of the series  $1 - x + x^2 - \&c.$ , where  $x$  is less than 1, or of the series  $1 - (1-x) + (1-x)^2 - (1-x)^3 + \&c.$ , where  $x$  is greater

*Phil. Mag. S. 3. Vol. 28. No. 185. Feb. 1846.* L

than 0, is  $\frac{1}{2}$ , but let no one thence attempt to draw the inference that  $1 - 1 + 1 - \&c. = \frac{1}{2}$ ; for  $1 - 1 + 1 - \&c.$  is not the limit of either of the series, it is simply the form they respectively assume when the variable has its limiting value, which is a very different thing, as we have seen.

The broad fact which, although as clear as the sun at noon day, so many seem to hesitate to admit, is, that when  $x$  is very small, *if it be an actual magnitude*,

$$1 - (1 - x) - (1 - x)^2 + \&c.$$

differs very little from  $\frac{1}{2}$ , but that when  $x$  vanishes, it assumes two values, 1 and 0. There is in this case no middle term between entity and non-entity. The idea is simple and the fact certain.

Prof. Young holds it to be an axiom, that "the value which suffices for all cases except the extreme case, will suffice for that too," or uses words to that effect. This is a most unwarrantable and false assumption. Take the case of the same series,

$$1 - x + x^2 - x^3 + \&c.,$$

where  $x$  is greater than 1. The value in this case, as is easily seen from equation (a.), is  $\pm \infty$  indifferently; and this holding always, except in the extreme case, when  $x = 1$ , it would follow, on Prof. Young's principle, that it holds in that too, and therefore that  $1 - 1 + 1 - \&c. = \pm \infty$ ; and he has before supposed it to be equal to  $\frac{1}{2}$ , which is absurd.

We may hence see the absurdity of any attempt to prove that  $1 - 1 + 1 - \&c.$ , considered as the limit of  $1 - x + x^2 - \&c.$ , where  $x$  is less than 1, is equal to  $\frac{1}{2}$ ; for by this nothing else can be meant than to prove that the one series is the limit of the other, which is contrary to the fact.

Prof. Young's attempt in this respect depends on the assumption that  $\left(1 - \frac{1}{\infty}\right)^{-\infty} = e$ , the base of the Napierian system, which is untrue, and which at any rate I challenge him to prove. Does he consider that  $(1 - 0)^{-\frac{1}{0}} = e$ ?

Liverpool, November 10, 1845.



Postscript.

I have just read Prof. Young's second paper, and without entering into all the windings of his argument, I shall proceed to animadvert upon such parts of it as refer to my own.

After some preliminary matter the Professor proceeds, "Let us now examine the series

$$\frac{1}{2} + A \cos \theta + A^2 \cos 2\theta + \&c. + A^n \cos n\theta,$$

so intimately connected with Fourier's integral, and which has already been the subject of consideration in Mr. Moon's paper before adverted to. This series, as there shown, or much more simply, by common division, arises from the development of the fraction

$$\frac{1 - A^2}{2(1 - 2A \cos \theta + A^2)}; \dots \dots [1.]$$

so that, taking account of the remainder of the division, the general equivalent of the series is this fraction minus

$$A^{n+1} \frac{\cos(n+1)\theta - A \cos n\theta}{1 - 2A \cos \theta + A^2} \dots \dots [2.]$$

Now confining our attention to the continuous values of A, it is obvious, upon the principles laid down in the former part of this paper, that in the extreme case of A = 1 and n = ∞, the fraction [2.] vanishes; and [1.] alone correctly represents the sum of the series in the limiting case."

What is meant by "confining our attention to the continuous values of A?" Can we draw any conclusion from the equation

$$\begin{aligned} & \frac{1}{2} + A \cos \theta + A^2 \cos 2\theta + \&c. + A^n \cos n\theta, \\ & = \frac{1 - A^2}{2(1 - 2A \cos \theta + A^2)} - A^{n+1} \frac{\cos(n+1)\theta - A \cos n\theta}{1 - 2A \cos \theta + A^2} \end{aligned}$$

other than the following, viz. that so long as A is positive, and differs from 1 by an actual magnitude, in the limit when n = ∞,

$$\begin{aligned} & \frac{1}{2} + A \cos \theta + A^2 \cos 2\theta + \&c. \text{ in inf.} \\ & = \frac{1 - A^2}{2(1 - 2A \cos \theta + A^2)}; \end{aligned}$$

and that when A ceases to differ from 1 by an actual magnitude, the same series

$$= \left\{ \frac{1 - A^2}{2(1 - 2A \cos \theta + A^2)} \right\}_{A=1} - \left\{ \frac{\cos(n+1)\theta - \cos n\theta}{2(1 - \cos \theta)} \right\}_{n=\infty} ?$$

L 2

Prof. Young appears to be haunted by the ghost of an argument, and it is somewhat difficult for others whose rest is not scared by the same phantom, to tell what he is driving at; but it appears to me that his difficulties, whatever they may be, arise from an inaccurate mode of expression which has crept into use, and of which he has not perceived the impropriety. Thus later on, we find him saying, "The real error, so frequently committed in analysis, consists in confounding

$$\frac{1}{2} + \cos \theta + \cos 2\theta + \&c. \text{ in } \textit{inf}.$$

with the limit of

$$\frac{1}{2} + A \cos \theta + A^2 \cos 2\theta + \&c. \text{ in } \textit{inf}.$$

and calling [1.], when  $A = 1$ , the sum of the former." Now I would observe that the expression, the limit of the series

$$\frac{1}{2} + A \cos \theta + A^2 \cos^2 \theta,$$

is a relative term, and in this case means the value to which the series tends, when the difference between  $A$  and 1 gradually diminishes; and from which value (or rather quantity) it can be made to differ by a quantity less than any that can be assigned. But there is no propriety in the expression, "the limit of the series when  $A = 1$ ." The phrase should be, *the particular value of the series when  $A = 1$* . It is perfectly true that so long as the difference between  $A$  and 1 is an *actual magnitude* (which phrase I use advisedly, as the abuse of the expressions *finite quantities* and *indefinitely small quantities* has led many people to believe that there is after all no essential difference between entity and non-entity), the series by diminishing that difference can be made to differ from

$$\left\{ \frac{1 - A^2}{2(1 - 2A \cos \theta + A^2)} \right\}_{A=1}, \text{ i. e. from } \frac{1}{2 \sin^2 \frac{\theta}{2}}$$

by a quantity less than any that can be assigned; but it does not thence follow, nor is it the fact, that when  $A = 1$  the series becomes  $= \frac{1}{2 \sin^2 \frac{\theta}{2}}$ ; for from the same evidence as

that by which we are led to the conclusion that when  $A < 1$  limit,

$$\frac{1}{2} + A \cos \theta + A^2 \cos 2\theta + \&c. \text{ in } \textit{inf} = \frac{1}{2 \sin^2 \frac{\theta}{2}},$$

we likewise deduce the fact that

$$\frac{1}{2} + \cos \theta + \cos 2\theta + \&c. \text{ in } \textit{inf.}$$

$$= \frac{1}{2 \sin^2 \frac{\theta}{2}} - \left\{ \frac{\cos (n+1)\theta - \cos n\theta}{2(1 - \cos \theta)} \right\} n = \infty.$$

Prof. Young indeed “considers it an axiom, that what holds for all but the extreme case will hold for that too,” but I must beg to submit that this is a matter of fact and not of opinion; and the fallacy of the principle in the present case I have sufficiently shown in the former part of this paper. With all due deference therefore to Prof. Young, I shall reassert, that Mr. De Morgan *is* in error in affirming (1.) to be the limit of the proposed series “when  $A = 1$ .” Omit the words “when  $A = 1$ ,” and I admit the proposition. Insert those words, and the fact expressed is untrue, if it be not wholly unmeaning.

Prof. Young says again, “It is easily proved that

$$\int_0^\infty e^{-ax} \cos x dx = \frac{a}{1+a^2}, \quad \int_0^\infty e^{-ax} \sin x dx = \frac{1}{1+a^2}, \quad (\alpha.)$$

from which it certainly follows, though the inference is denied by Mr. Moon, that in the limit, when  $a = 0$ , the true values of these integrals are 0 and 1.” I do not deny the inference, that the limits of the integrals when  $a$  is diminished indefinitely, so long as it continues an actual magnitude, are 0 and 1; but I do deny that the limits of the integrals are to be found by putting  $a = 1$  in the left-hand members of the two equations ( $\alpha.$ ), that is, I deny that

$$\int_0^\infty \cos x dx \quad \text{and} \quad \int_0^\infty \sin x dx$$

are the limits of

$$\int_0^\infty e^{-ax} \cos x dx, \quad \int_0^\infty e^{-ax} \sin x dx,$$

which is all I care to establish.

December 10, 1845.

### Postscript 2.

I have read Prof. Young’s third paper. Had my reply to his first paper been inserted in proper course, it is probable that he would have saved himself the trouble of writing, and the public of reading his last two papers. The staple of what I have to advance in opposition to this last is contained in my two previous notices. A few words, however, are still called for.

The Professor begins, "The general expression for the sum of the infinite series

$$1 - x + x^2 - x^3 + x^4 - \&c.$$

is

$$S = \frac{1}{1+x} - \frac{x^\infty}{1+x}."$$

What may be meant by the recondite symbol  $\infty$ , I do not profess to understand. But it appears to me that

$$S = \frac{1}{1+x} - \frac{(-x)^\infty}{1+x};$$

and I am induced to think that Prof. Young himself will come to be of the same opinion when he again undertakes to examine the subject.

It is unnecessary for me to reply to the argument of the present paper, which appears to rest upon one of Prof. Young's previous fallacies which I have elsewhere exposed. I shall merely advert to the conclusion at which he ultimately arrives, "that it is indisputably true that the extreme of the *convergent* cases of the above series S, usually written in the form

$$1 - 1 + 1 - 1 + \&c.$$

is  $\frac{1}{2}$ , and that the extreme of the *divergent* cases, usually written in the same form, is really infinite, as stated in his former paper."

I am afraid that Prof. Young will be apt to mystify both himself and his readers by talking about "the extreme of the divergent cases" and "the extreme of the convergent cases." If these "extremes" are usually written under the above form, I can only say that such usage is "extremely" improper. But let Prof. Young define the terms he uses. What is meant by "the extreme of the *convergent* series" for example? Is it the value of the series

$$1 - (1-x) + (1-x)^2 + (1-x)^3 - \&c. \dots (\alpha.)$$

when  $x = 0$ ? If so, I beg to assure him that the extreme value of the convergent series is *not*  $\frac{1}{2}$ , but 1 or 0 indifferently. The fact is, it is absurd to talk of "extreme values" in these cases. If  $x$  be made ever so small, there will always be some smaller value which might be assigned to it; so that it is impossible to assign an extreme value to  $x$  so long as it is an existing magnitude, and the moment it ceases to be such the series ceases to be equal (or rather to approximate) to  $\frac{1}{2}$ . Twist and turn it as he may, Prof. Young will never be able to prove the series

$$1 - x + x^2 - x^3 + \&c. = \frac{1}{2}.$$

It may be made to differ from that quantity by a quantity less than any that can be assigned, but it is *never* actually to it—least of all is it so when  $x=0$ , in which case it assumes a totally different value.

It was not for nothing that Newton devised his method of limits, and to the present case it applies with peculiar clearness and beauty. But no longer to fight with shadows, I shall take up a definite position, and shall leave it to Professor Young to drive me from it if he can. I assert, then, that the series

$$1 - (1 - x) + (1 - x)^2 - (1 - x)^3 + \&c.,$$

so long as  $x$  lies between 0 and 1, and differs from each of them by an *actual* magnitude (I do not say a sensible magnitude, for the present is not a question of degree), approaches to  $\frac{1}{2}$  as its limit when  $x$  is made to diminish; that when  $x=0$ , the absolute value of the series is 1 or 0 indifferently; that when  $x$  is less than 0, it becomes  $\pm \infty$  indifferently; and I defy any man now living, or as the lawyers say, who may hereafter come *in esse*, to prove anything else, be it more or less concerning it.

Prof. Young considers that the conclusion, that “the extreme of the *divergent* cases is really infinite,” could “never have been anticipated from the theory hitherto prevalent.” Protesting as I do against the use of the term “extreme of the divergent cases,” I may say that long before Prof. Young either said or wrote a word upon the subject, I had shown that *all the divergent cases* have the value  $\pm \infty$  indifferently.

Again, Prof. Young says, “if he has been anticipated in any of these views, which are doubtless calculated to produce a reform in the existing theory, he hopes to be informed of the circumstance through the medium of this Journal.” I beg to assure him therefore, through the pages of this Journal, that *all his* views which are not erroneous (though what proportion that may be I confess myself unable to state, as I do not understand very clearly what they are) *have been anticipated* in my paper dated March 17, 1845, and published in the number of this Journal for June in the past year.

I now await Prof. Young’s answer, trusting that I may not be under the necessity of replying to any more of his papers till he has had an opportunity of reading some of mine.

January 3, 1846.

XXVI. *Remarks on a Paper by Mr. Moon on Fresnel's Theory of Double Refraction\**. By JESUITICUS.

THE hypothesis on which Fresnel's theory of double refraction is based, is the following:—

“That the displacement of a molecule of the vibrating medium in a crystallized body is resisted by different elastic forces according to the different directions in which the displacement takes place.”

This is not a mere speculative hypothesis, but is based on experiment. It is found that glass, possessing only the power of single or ordinary refraction, may be made by the application of heat, or by mechanical pressure, to possess that of double refraction.

It is further supposed that the medium is symmetrical with respect to three rectangular axes in space, but, in general, not symmetrical with respect to any other axis through the same origin. These axes are called the axes of elasticity. It is then proved, that if any particle of the æther be suddenly displaced, *the other particles remaining quiescent*, the force of restitution developed by such disturbance will not in general be in the direction of the displacement, but only when such displacement is in the direction of the aforesaid axes of elasticity. The elegant demonstration of Smith, quoted by Mr. Moon, is by Mr. Moon's own showing fully adequate to establish the theorem as I have enunciated it, which is doubtless the sense in which Fresnel (the illustrious Fresnel, “whose name is enrolled amongst those which pass not away,”) doubtlessly conceived it.

Any one who understands the subject must at once acknowledge that any theory of light must be, to a considerable extent, imaginative; and that theory which can explain the greatest number of facts ought to claim the attention of the philosopher more than any other. It is to this that the undulatory theory owes its great celebrity, and of all parts of the undulatory theory, that of double refraction is the most extraordinary. It ought to be regarded as a stupendous monument of human ingenuity. It must not be forgotten how admirably the properties of uniaxal crystals follow from the general investigation of the biaxal class; but above all, how from this same investigation, conical and cylindrical refraction were discovered by Sir William Hamilton. Such an unexpected refinement as this, which probably would never have been recognised by the mere experimentalist, undirected by the skill

\* Phil. Mag. S. 3. No. 183. vol. xxvii.

of so great an analyst, is surely no slight recommendation of the theory.

Mr. Moon subsequently gives a quotation from Airy's Tracts, concerning which he is by no means sparing in arrogant and supercilious criticism. But is it likely that Airy would make such a fool of himself as Mr. Moon earnestly endeavours to represent? It must be remembered, that at the time Airy's Tracts were published, very little of the undulatory theory was studied or known in Cambridge.

It was the part of this philosopher, therefore, to put everything as much as possible in the clearest and most simple point of view. That there are, and will perhaps long continue to be, difficulties in the undulatory theory, none of its supporters will deny. None of those difficulties are *shirked* or glossed over in the Tract of Airy; he plainly acknowledges each as it arrives. He no doubt himself considered the part quoted by Mr. Moon more as an illustration than anything else. Those who wish to see the matter treated with all the analytical generality of which it is capable, are referred to a tract on the Reflexion and Refraction of Light at the Surface of two contiguous Media, by the late famous George Green, in the Cambridge Philosophical Transactions.

I have one word more with Mr. Moon. He says, that on substituting for  $u$ ,

$$u - \frac{du}{dx}h + \frac{d^2u}{dx^2} \frac{h^2}{1.2}, \&c.,$$

and for  $u'$ ,

$$u + \frac{du}{dx}h + \frac{d^2u}{dx^2} \frac{h^2}{1.2}, \&c.,$$

that  $h$  is considered small with respect to  $u$ .

Does Mr. Moon know anything of analysis? He was eighth wrangler in 1838, and therefore he ought to know something. His knowledge, however, has served him miserably on this occasion. The substitutions, stopping at  $h^2$ , merely require that  $h$  should be small in comparison with the *length of a wave*, not with respect to  $u$ .

JESUITICUS.

XXVII. *Observations on the subject of the Preceding Communications.* By the EDITORS.

ON the subject of the foregoing letter, the Editors are induced to subjoin a few remarks, as besides the attacks on Fresnel, they have also received from Mr. Moon one more paper, containing stric-

tures on the writings of another distinguished mathematician, besides the reply to Mr. Young which they now publish.

In the admission of mathematical articles, the Editors are obliged to consult both quantity and character, as follows :—

It is not in their power to admit any very great quantity of *pure* mathematics. The majority of the readers of the Magazine are more interested in other sciences, and the Magazine would soon cease to exist if it were more than sparingly supplied with articles on lofty mathematical subjects.

As to the character of their mathematical articles, the Editors are placed in a peculiar position. They do not themselves profess to be so conversant with the higher mathematics as to rely entirely on their own judgement. In the articles which they insert, they must be guided by opinions. If they occasionally insert an article in which the general opinions of mathematicians are controverted, it is because they feel that mathematicians themselves would occasionally like to see the manner in which dissent from generally received principles manifests itself; and because they know that such occasional insertion will not, in the eyes of those same mathematicians, make them, the Editors, appear to be assuming a side in controversies of the merits of which they are not sufficient judges.

But if the Editors were to lend their Magazine to an extensive system of attack upon any usual results and methods of mathematics, either pure or mixed, they feel that they could not escape the charge of presumption. Whatever might be thought of an occasional paper, they feel sure that a series of such papers would cause them to be considered, and justly considered, as expressing an opinion on matters in which their knowledge is but limited. They would make just the same answer to a proposition for as extensive a system of defence to be inserted in their Magazine. They would suggest to both assailants and defendants to carry their communications to quarters in which they will find more competent judges. The pages of the Philosophical Transactions, of the Memoirs of the Royal Irish Academy, of the Cambridge Philosophical Society, of the Cambridge Mathematical Journal, &c., are much fitter vehicles for extensive mathematical discussion than those of the Philosophical Magazine.

As to one point, however, the Editors feel that a responsibility rests upon themselves, namely, as to the tone and temper in which controversial communications are expressed. They are persuaded that the differing opinions of men of science upon difficult subjects may be fully conveyed without any deviation from the respect and courtesy with which public discussion ought to be conducted :—and they feel regret when anything which is objectionable in this respect obtains a place in their pages.



XXVIII. *Proceedings of Learned Societies.*

## ROYAL SOCIETY.

“**E**XPERIMENTAL Researches in Electricity.” By Michael Faraday, Esq., D.C.L., F.R.S. &c. Twentieth Series. Section 26th. “On New Magnetic Actions; and on the Magnetic Condition of all Matter.”

The following is the order in which the several divisions of the subject treated of in this section of the author’s researches in electricity succeed one another:—1. Apparatus required. 2. Action of magnets on heavy glass. 3. Action of magnets on other substances acting magnetically on light. 4. Action of magnets on the metals generally. 5. Action of magnets on the magnetic metals and their compounds. 6. Action of magnets on air and gases. 7. General considerations.

In giving an account of the contents of this paper, any attempt to follow the track of the author in the precise order in which he relates the consecutive steps of his progress in this new path of discovery, would fail of accomplishing its object: for, by adhering to such a course, it would scarcely be possible to comprise within the requisite limits of an abstract the substance of a memoir extending, as the present one does, to so great a length, and of which so large a portion is occupied with minute and circumstantial details of experiments; or to succeed in conveying any clear and distinct idea of the extraordinary law of nature brought to light by the author, and of the important conclusions which he has deduced.

One of the simplest forms of experiment in which the operation of this newly-discovered law of magnetic action is manifested, is the following:—A bar of glass, composed of silicated borate of lead, two inches in length, and half an inch in width and in thickness, is suspended at its centre by a long thread, formed of several fibres of silk cocoon, so as to turn freely, by the slightest force, in a horizontal plane, and is secured from the agitation of currents of air by being enclosed in a glass jar. The two poles of a powerful electro-magnet are placed one on each side of the glass bar, so that the centre of the bar shall be in the line connecting the poles, which is the line of magnetic force. If, previous to the establishment of the magnetic action, the position of the bar be such that its axis is inclined at half a right angle to that line, then, on completing the circuit of the battery so as to bring the magnetic power into operation, the bar will turn so as to take a position at right angles to the same line; and, if disturbed, will return to that position. A bar of bismuth, substituted for the glass bar, exhibits the same phenomenon, but in a still more marked manner. It is well known that a bar of iron, placed in the same circumstances, takes a position coincident with the direction of the magnetic forces; and therefore at right angles with the position taken by the bar of bismuth subjected to the same influence. These two directions are termed by the author *axial* and *equatorial*; the former being that taken by the iron, the latter that taken by the bismuth.

Thus it appears that different bodies are acted upon by the magnetic forces in two different and opposite modes ; and they may accordingly be arranged in two classes ; the one, of which iron is the type, constituting those usually denominated *magnetics* ; the other, of which bismuth may be taken as the type, obeying a contrary law, and therefore coming under the generic appellation of *diamagnetics*. The author has examined a vast variety of substances, both simple and compound, and in a solid, liquid, or gaseous form, with a view to ascertain their respective places and relative order with reference to this classification. The number of simple bodies which belong to the class of magnetics is extremely limited, consisting only of iron, which possesses the magnetic property in an eminent degree, nickel, cobalt, manganese, chromium, cerium, titanium, palladium, platinum and osmium. All other bodies, when either solid or liquid, are diamagnetic ; that is, obey the same law, with regard to magnetic action, as bismuth, but with various degrees of intensity : arsenic is one of those that give the feeblest indications of possessing this property. The following exhibit it in increasing degrees, according to the order in which they are here enumerated ; namely, ether, alcohol, gold, water, mercury, flint glass, tin, lead, zinc, antimony, phosphorus, bismuth. On the other hand, no gaseous body of any kind, or in any state of rarefaction or condensation, affords the slightest trace of being affected by magnetic forces. Gases may therefore be considered as occupying the neutral point in the magnetic scale, intermediate between magnetic and diamagnetic bodies.

The magnetic properties of compound bodies depend on those of their elements ; and the bodies are rendered either magnetic or diamagnetic according to the predominance of one or other of these conditions among their constituent parts. Thus iron is found to retain its magnetic power when it has entered into combination with other bodies of the diamagnetic class ; the two forces acting in opposition to one another, and the resulting effect being only that due to the difference in their power. Hence the oxides and the salts of iron are still in a certain degree magnetic, and the latter even when they are held in solution by water ; but the water may be present in such a proportion as that neither shall prevail ; and the solution, as far as respects its magnetic properties, will then be exactly neutralized. These saline solutions, prepared of various degrees of strength, also afford a convenient method of comparing the relative degrees of force, both magnetic and diamagnetic, of different bodies, whether solid or fluid, but more especially the latter, as they admit of the body under examination being suspended in another liquid, when its position of equilibrium will indicate which of the two substances has the strongest magnetic power.

In one respect, indeed, the diamagnetic action presents a remarkable contrast with the magnetic ; and the difference is not merely one of degree, but of kind. The magnetism of iron and other magnetics characterized by polarity ; that of diamagnetics is devoid of any trace of polarity ; the particles of two bodies of the latter class, when jointly under the influence of the magnetic forces, manifest

towards each other no action whatever, either of attraction or repulsion. It has long been known that the magnetism of iron is impaired by heat; and it has been generally believed that a certain degree of heat destroys it entirely. The author finds, however, that this opinion is not correct; for he shows that, by applying more powerful tests than those which had been formerly confided in, iron, nickel and cobalt, however high their temperature may be raised, still retain a certain amount of magnetic power, of the same character as that which they ordinarily possess. From the different temperatures at which the magnetic metals appear to lose their peculiar power, it had formerly been surmised by the author that all the metals would probably be found to possess the same character of magnetism, if their temperature could be lowered sufficiently; but the results of the present investigation have convinced him that this is not the case, for bismuth, tin, &c. are in a condition very different from that of heated iron, nickel or cobalt.

The magnetic phenomena presented by copper and a few other metals are of a peculiar character, differing exceedingly from those exhibited by either iron or bismuth, in consequence of their being complicated with other agencies, arising from the gradual acquisition and loss of magnetic power by the iron core of the electro-magnet, the great conducting power of copper for electric currents, and its susceptibility of being acted upon by induced currents of magneto-electricity, as described by the author in the first and second series of these researches. The resulting phenomena are to all appearance exceedingly singular and anomalous, and would seem to be explicable only on the principles referred to by the author.

Pursuing his inductive inquiries with a view to discover the primary law of magnetic action from which the general phenomena result, the author noticed the modifications produced by different forms given to the bodies subjected to experiment. In order that these bodies may set either axially or equatorially, it is necessary that their section, with reference to the plane of revolution, be of an elongated shape: when in the form of a cube or sphere they have no disposition to turn in any direction: but the whole mass, if magnetic, is attracted towards either magnetic pole; if diamagnetic, is repelled from them. Substances divided into minute fragments, or reduced to a fine powder, obey the same law as the aggregate masses, moving in lines, which may be termed *diamagnetic curves*, in contradistinction to the ordinary magnetic curves, which they everywhere intersect at right angles. These movements may be beautifully seen by sprinkling bismuth in very fine powder on paper, and tapping on the paper while subjected to the action of a magnet.

The whole of these facts, when carefully considered, are resolvable, by induction, into the general and simple law, that while every particle of a magnetic body is attracted, every particle of a diamagnetic body is repelled, by either pole of a magnet. These forces continue to be exerted as long as the magnetic power is sustained, and immediately cease on the cessation of that power. Thus do these two modes of action stand in the same general antithetical re-

lation to one another as the positive and negative conditions of electricity, the northern and southern polarities of ordinary magnetism, or the lines of electric and of magnetic force in magneto-electricity. Of these phenomena, the diamagnetic are the most important, from their extending largely, and in a new direction, that character of duality which the magnetic force was already known, in a certain degree, to possess. All matter, indeed, appears to be subject to the magnetic force as universally as it is to the gravitating, the electric, the cohesive and the chemical forces. Small as the magnetic force appears to be in the limited field of our experiments, yet when estimated by its dynamic effects on masses of matter, it is found to be vastly more energetic than even the mighty power of gravitation, which binds together the whole universe: and there can be no doubt that it acts a most important part in nature, and conduces to some great purpose of utility to the system of the earth and of its inhabitants.

Towards the conclusion of the paper, the author enters on theoretical considerations suggested to him by the facts thus brought to light. An explanation of all the motions and other dynamic phenomena consequent on the action of magnets on diamagnetic bodies might, he thinks, be offered on the supposition that magnetic induction causes in them a state the reverse of that which it produces in magnetic matter: that is, if a particle of each kind of matter were placed in the magnetic field, both would become magnetic, and each would have its axis parallel to the resultant of magnetic force passing through it; but the particle of magnetic matter would have its north and south poles opposite to, or facing the contrary poles of the inducing magnet; whereas, with the diamagnetic particles, the reverse would obtain; and hence there would result, in the one substance, approximation; in the other, recession. On Ampère's theory, this view would be equivalent to the supposition that, as currents are induced in iron and magnetics, parallel to those existing in the inducing magnet or battery wire, so, in bismuth and other diamagnetics, the currents induced are in the contrary direction. As far as experiment yet bears upon such a notion, the inductive effects on masses of magnetic and diamagnetic metals are the same.

## XXIX. *Intelligence and Miscellaneous Articles.*

### ANALYSIS OF A SUBSTANCE OCCURRING WITH DISTHENE.

BY M. A. DELESSE.

**I**N most mineralogical collections the disthene of Pontivy occurs, crystallized in large prisms of a sky-blue colour; they are often nearly 4 inches long and about  $\frac{4}{10}$ ths of an inch wide; the spaces occurring between the crystals are filled with a white lamellar, pearly substance, which is sometimes so intermixed with the cleavable faces of the prisms of disthene, that it is difficult to determine the limits of the two minerals.

This substance differs from any hitherto described, and possesses

the following characters : it has the form of small crystalline laminae, usually radiating from a centre ; it is sometimes separated from disthene by a thin stratum of yellowish oxide of iron, evidently resulting either from the recent decomposition of disthene, or of the rock in which it occurs. In fragments, the substance is yellowish-white, and translucent ; is cut by the knife, and formed by the agglomeration of a multitude of small laminae, indicating a radiated structure ; these laminae are perfectly transparent, but have no crystalline form.

The cohesion of this substance is slight, but still it is difficult to reduce it to a fine powder. When pulverized it has the appearance of small scales of a shining silvery-white colour, with a pearly lustre ; it is soft to the touch, but not unctuous like talc ; it is harder than talc, for it scratches it ; but it is not so hard as fluor spar ; its density is 2.792 ; after drying at 212° and heated in a tube, it yields water ; when dried over sulphuric acid *in vacuo*, it loses only a few thousandths of its weight, and retains its water, which is consequently in a state of combination ; heated on platina it swells and becomes milk-white ; when more strongly heated, it agglutinates and then fuses, but with difficulty, into a white enamel ; it is phosphorescent and emits a brilliant light ; with nitrate of cobalt it becomes of a pure blue colour, when strongly heated ; with borax it dissolves readily and perfectly, with a slight colour proceeding from iron ; with the salt of phosphorus it yields a colourless crystalline bead ; the solution is quite complete, no silica skeleton remaining ; with carbonate of soda effervescence ensues ; alumina is left unacted upon, even when excess of the carbonate is employed.

Neither hydrochloric acid nor aqua regia acts upon this substance, but when finely levigated and boiled with concentrated sulphuric acid, it is completely decomposed ; the silica remains in the granular state and retains the form of the scales. After calcination, it is not acted upon by the acid.

The qualitative analysis of this substance shows that it contains silica, alumina, a little iron and manganese, the last two not appearing to be in a state of combination, potash and water ; soda was not found to be present, which it is proper to state, for usually the two alkalies occur together. As the mineral possesses some of the characters of mica, fluorine was sought for but not found.

In determining the quantity of water, it was found requisite to heat it pretty strongly to separate the whole of it ; when only a part of it was expelled, it was found that on placing it in water for some days and then drying it by exposure to the air, it regained exactly as much water as it had lost ; when however it is strongly heated and loses all the water, it does not regain it by immersion.

Analysed by means of nitrate of barytes, this substance yielded—

Silica . . . . .	45.48
Alumina . . . . .	38.20
Potash . . . . .	11.20
Water . . . . .	5.24

---

100.12

with traces of iron and manganese.

We may therefore consider it as composed of—

Twelve eqs. of silica . . . . .	16 × 12 = 192	44·75
Nine . . . alumina . . . . .	18 × 9 = 162	37·76
One . . . potash . . . . .	= 48	11·20
Three . . . water . . . . .	9 × 3 = 27	6·29
	429	100·

*Ann. de Ch. et de Phys.*, Octobre 1845.

HYDRATED SILICATE OF MAGNESIA. BY M. A. DELESSE.

This substance is arranged at the Ecole des Mines with the mineral species which M. Breithaupt has named *Kerolite*, and it appeared to M. Delesse to require examination. It comes from Germany, its locality however is unknown; but it has evidently occurred in serpentine.

Its colour is yellowish-white, it is opaline and slightly transparent; its fracture resembles wax, and it is greasy to the touch; it is occasionally spotted with milk-white spots, which appear to be a different substance. Its specific gravity is 2·335; when slightly heated in a glass tube it becomes black and loses water; when strongly heated it becomes of a dead-white colour, and loses its transparency. The black colour appears to be owing to bitumen, for it disappears when the substance is strongly heated in a closed tube; this property belongs also to kerolite, mestaxite, saponite, &c.

When put into water after calcination it emits a great number of bubbles of gas, becomes hard, and is with difficulty acted upon by acids; whereas before heating it is scratched by calcspar and easily acted upon; it is completely infusible; with the salt of phosphorus it gives a skeleton of silica.

A qualitative analysis showed that this substance contains only water, silica, magnesia, a little alumina and traces of iron, which appear to be in the state of peroxide, and disseminated in small veins throughout the mass.

By analysis this mineral gave—

Silica . . . . .	53·5
Magnesia . . . . .	28·6
Alumina and a trace of oxide of iron	00·9
Water . . . . .	16·4
	99·4

*Annales des Mines*, 1844.

ANALYSIS OF THE ELIE PYROPE OR GARNET.

BY PROF. CONNELL.

This mineral, which is known to amateur collectors under the name of Elie ruby, is found on the sea shore at Elie, in the county of Fife, proceeding from the debris of trap-rocks. It has been long known to Scottish mineralogists, and has been regarded as one of the varieties of precious garnet, and is occasionally called pyrope. It is not crystallized, but occurs in angular grains, which evidently

have not come from any distance. Its other leading external characters, including transparency and colour, agree with those of precious garnet and pyrope, the colour approaching the deeper tint of the latter; its specific gravity is 3.661.

Twenty grains of this mineral in very fine powder, were fused with four times their weight of carbonate of potash; the mass was treated with muriatic acid, and no smell of chlorine observed. Silica was separated by the usual method; the precipitate obtained by ammonia was dissolved in muriatic acid, the solution boiled with excess of potash which took up the alumina, and the matter left by this alkali was dissolved in muriatic acid; to this solution tartaric acid and ammonia in excess were added, and a current of sulphuretted hydrogen, passed into it, threw down sulphuret of iron with a little sulphuret of manganese. The filtered liquid was evaporated to dryness, and the residue incinerated was pure white; it was carefully examined for yttria, which Dr. Apjohn, some few years ago, announced that he discovered in pyrope. This white matter was dissolved in muriatic acid, and muriate of ammonia and excess of ammonia were added; a gelatinous precipitate fell, which by ignition acquired a greenish-yellow tint, and magnesia was left in solution. The ignited precipitate was again dissolved in muriatic acid, and yielded a gelatinous precipitate by treatment with ammonia and its muriate; this was dissolved to a great extent by potash, leaving a substance which was principally oxide of iron, but gave a permanent fine, though pale emerald-green colour to salt of phosphorus, and therefore contained a trace of oxide of chromium. It was determined by a separate experiment that the iron contained in the mineral was entirely in the state of peroxide.

One hundred parts of this substance were found to consist of

Silica . . . . .	42.80
Alumina . . . . .	28.65
Peroxide of iron . . . . .	9.31
Protoxide of manganese . . . . .	0.25
Lime . . . . .	4.78
Magnesia . . . . .	10.67
Oxide of chromium, trace	
	96.46

The deficiency Prof. Connell conceives to be probably owing to some magnesia which might have escaped precipitation by the carbonate of potash.

Prof. Connell remarks, that even if the oxide of iron in this mineral were held to be protoxide (instead of peroxide, as he found it), there would be quite as much difficulty in bringing the result under the garnet formula as there is in bringing the leading analyses of Bohemian pyrope under it. This circumstance, as well as the general conformity between the above result and the analyses of pyrope, comprising those of Klaproth, Wachtmeister and Von Kobell, particularly as respects the considerable quantity of magnesia and the comparatively small quantity of oxide of iron, notwithstanding the

discrepancy as to the state of oxidation of the latter, tend to show a close connexion between the Elie mineral and pyrope. The occurrence of oxide of chromium in both minerals, and their specific gravity, which is 3·661 for the Elie mineral, 3·78 for pyrope, while that of precious garnet exceeds 4, lead to a similar view of this connection.—*Jameson's Journal*, Oct. 1845.

---

ANALYSIS OF METEORIC IRON FROM BURLINGTON, OSTEGO COUNTY, NEW YORK. BY MR. C. H. ROCKWELL.

In the year 1819 two or three masses of native iron, as it appeared to be, were procured from the farmer who first turned it over with his plough, in a field near the north line of the town of Burlington, Ostego County, New York. These consisted of remnants of an entire mass originally supposed to weigh between one and two hundred pounds, and found several years before. It had been in the forge of a country blacksmith, and the whole heated in order to enable him to cut off portions for the manufacture of such articles as the farmer most needed.

The mass was divided by broad laminæ, crossing each other at an angle of 60° and 120°, cutting up the surface into triangular and rhombohedral figures. It broke with a hackly fracture, and only with the greatest difficulty on the thinnest edges.

Two deep and broad sutures marked its two most regular opposite faces, made by the wedge or chisel by the smith, who severed it from the adjoining portion. It bore the marks of having been intensely heated in the forge, and numerous microscopic crystals, of a black colour and brilliant lustre, covered some parts of its surface; they resembled phosphate of iron, but were too small to be detached.

The specific gravity was 7·501; it dissolved quickly and completely in nitric acid, with the application of a gentle heat. The solution, treated with nitrate of silver, gave no cloudiness, showing the absence of chlorine; it yielded by the usual process for separating iron from nickel,

Iron.....	92·291
Nickel.....	8·146
	100·437

No trace of any other substances could be detected.—*Silliman's Journal*, vol. xlvi.

---

PREPARATION OF CHLORO-ACETIC ACID.

M. Malaguti recommends the following process for the preparation of chloro-acetic acid readily and in large quantity:—Let chlorine act upon sulphuric æther, by which sesquichloride of carbon is obtained, and then the water which is suffered to remain in the bottles with the rough product is merely a solution of chloro-acetic and hydrochloric acids; or perchloric æther is prepared, and by distilling it and causing the product of the distillation to mix with water, a solution of chloro-acetic and hydrochloric acids is obtained. In both



cases it is sufficient to employ a vacuum with sulphuric acid and potash to obtain chloro-acetic acid in a state of great purity. This process possesses the advantage of also obtaining as a secondary product, a considerable quantity of sesquichloride of carbon.—*Ann. de Ch. et de Phys.*, Jan. 1846.

---

COMPOSITION OF PHOSPHATE OF AMMONIA AND MAGNESIA.

BY M. FRESENIUS.

The erroneous statements contained in chemical treatises with respect to the ammoniaco-magnesian phosphate have given inaccurate results as to the proportions of magnesia indicated by this salt, and they have also prevented its being employed in estimating the quantity of phosphoric acid. M. Fresenius has discovered that the double salt in question is absolutely insoluble in free ammonia, so that it may be employed in quantitative analysis.

He ascertained the solubility of this salt in water, solution of ammonia, solution of hydrochlorate of ammonia, and in a mixed solution of ammonia and hydrochlorate. He found it dissolved by 15293 parts of water at the usual temperature, and requires a mean quantity of 44330 parts of ammoniacal water for solution, so that one part of magnesia, in the form of this salt, requires 120760 parts of water, and one part of phosphoric acid 70000. According to these statements this salt may be for a long time washed with ammoniacal water before dissolving a very minute fraction of a grain either of magnesia or phosphoric acid.

As to solution of hydrochlorate of ammonia, one part of the double salt is dissolved by 7548 parts of it, and by 15627 parts of a mixed solution of ammonia and hydrochlorate; sal-ammoniac therefore slightly increases the solubility of the salt; still however this solubility is so slight as to be inappreciable in estimating its quantity.

M. Fresenius has also performed some comparative experiments to ascertain if the double phosphate would answer for analyses.

He analysed a determinate quantity of very pure sulphate of magnesia; by calculation the magnesia was estimated at 34.01 per cent.; experiment gave 34.0 and 34.02 per cent.; the phosphoric acid of phosphate of magnesia was calculated at 19.90 per cent., while experiment gave 19.87.—*Journ. de Pharm. et de Ch.*, Dec. 1845.

---

COMPOSITION OF COMMON PHOSPHATE OF SODA.

M. Fresenius states that the undermentioned chemists found this salt to consist of

	Berzelius.	Malaguti.	Graham.	Clark.
Phosphoric acid	20.33	18.80	37.1	37.48
Soda . . . . .	17.67	16.71		
Water . . . . .	62.00	64.25	62.9	62.52
	100.00	99.76	100.00	100.00

M. Fresenius found 19.87 of phosphoric acid and 62.67 of water;

his results therefore agree with those of Berzelius, Graham and Clark.  
—*Journ. de Pharm. et de Ch.*, Dec. 1845.

ON SEVERAL NEW SERIES OF DOUBLE OXALATES. BY M. REES  
HEECE.

These salts were discovered in investigating the action of alkaline and earthy bases on the oxalates of the sesquioxides.

It is well known that the salts of lime produce but a slight precipitation of oxalate of lime in a moderately concentrated solution of the oxalates of the sesquioxides of iron and chromium, &c., and none in a very dilute solution of oxalate of chromium and potash, a salt discovered by Prof. Gregory, and in which there are 3 equiv. of oxalic acid combined with the alkaline base. A concentrated solution of the same salts gives rise to an abundant precipitate, which has been considered as oxalate of lime, but in which I found a considerable proportion of chromium. These were the facts which led me to pursue this inquiry.

The combination by means of which I have prepared the double salts which are the objects of this memoir, is an oxalate of chromium and ammonia, having the same formula as the salt of Mr. Gregory; but it is preferable to this on account of its great solubility.

A concentrated solution of this salt, mixed with its volume of chloride of strontium, barium or calcium, yields voluminous precipitates, which, separated from the mother-leys and recrystallized, have the following composition:—

Oxalate of chrome and barytes (A)	$3\text{C}^2\text{O}^3 + \text{Cr}^2\text{O}^3 + 3(\text{C}^2\text{O}^3\text{BaO}) + 12\text{HO}$ .
Oxalate of chrome and barytes (B)	$3\text{C}^2\text{O}^3 + \text{Cr}^2\text{O}^3 + 2(\text{C}^2\text{O}^3\text{BaO}) + 18\text{HO}$ .
Oxalate of chrome and strontian ...	$3\text{C}^2\text{O}^3 + \text{Cr}^2\text{O}^3 + 3(\text{C}^2\text{O}^3\text{SrO}) + 18\text{HO}$ .
Oxalate of chrome and lime .....	$2(3\text{C}^2\text{O}^3 + \text{Cr}^2\text{O}^3) + 3(\text{C}^2\text{O}^3\text{CaO}) + 36\text{HO}$ .

If oxide of iron is substituted for the oxide of chromium, we obtain the corresponding salts with an iron base, and which are represented by the following formulæ:—

Oxalate of iron and barytes .....	$3\text{C}^2\text{O}^3 + \text{Fe}^2\text{O}^3 + 3(\text{C}^2\text{O}^3\text{BaO}) + 7\text{HO}$ .
Oxalate of iron and barytes .....	$3\text{C}^2\text{O}^3 + \text{Fe}^2\text{O}^3 + 3(\text{C}^2\text{O}^3\text{BaO}) + 12\text{HO}$ .
Oxalate of iron and strontian .....	$3\text{C}^2\text{O}^3 + \text{Fe}^2\text{O}^3 + 3(\text{C}^2\text{O}^3\text{SrO}) + 18\text{HO}$ .

The oxalate of iron and of lime does not crystallize.

If alumina is substituted for the oxide of chromium, we obtain similar salts, which are represented by—

Oxalate of alumina and barytes.....	$3\text{C}^2\text{O}^3 + \text{Al}^2\text{O}^3 + 3(\text{C}^2\text{O}^3\text{BaO}) + 10\text{HO}$ .
Oxalate of alumina and barytes.....	$3\text{C}^2\text{O}^3 + \text{Al}^2\text{O}^3 + 3(\text{C}^2\text{O}^3\text{BaO}) + 30\text{HO}$ .
Oxalate of alumina and strontian .....	$3\text{C}^2\text{O}^3 + \text{Al}^2\text{O}^3 + 2(\text{C}^2\text{O}^3\text{SrO}) + 18\text{HO}$ .

The oxalate of alumina and lime cannot be isolated in a state of purity, on account of its insolubility.

These salts crystallize in small silky needles; those of the oxide of chromium are of a dark violet colour, those of iron of a greenish-yellow, and those of alumina of a brilliant white. They are soluble in about 30 times their weight of boiling water (excepting the salts of lime and oxide of chromium, of alumina and strontia, which are decomposed by water); they are scarcely soluble in cold. All the alkalis decompose them by precipitating the sesquioxide and earthy

oxalate. The salts of chromium however behave differently towards ammonia, which does not throw down the oxide of chromium, even when the barytes has been separated by sulphuric acid.

The iron salts are decomposed by the solar rays, with an abundant disengagement of carbonic acid, even when the crystals are dry.

I shall conclude this summary of my investigations by drawing attention to the importance of these salts in analysis; the more so as it was with this view I undertook them.

I think that I have succeeded in explaining the fact, long since known, of the solubility of the oxalate of lime in solutions of the sesquioxides. Iron, aluminum and chromium are always separated from their ores as sesquioxides; lime and strontia, in the state of oxalates; but we know that lime cannot be separated from the solution of the sesquioxide; this circumstance is owing to the formation of a double salt, of which the oxalate of lime forms a part. It is therefore necessary to precipitate the sesquioxide by ammonia, which leaves the lime free in the solution, and it is therefore difficult to prevent its being thrown down by the carbonic acid of the atmosphere, and thus affecting the weight of the sesquioxide.

It is especially in the case of alumina and oxide of chromium, that the error may be the greatest. This difficulty is easily avoided by the following process:—I will select as an instance a mineral containing iron and lime. It is dissolved in hydrochloric acid; then a suitable quantity of oxalic acid added, which, if the liquor is diluted, will not produce any precipitate; I now add some oxalate of ammonia in excess, which will precipitate the whole of the lime, which is separated by filtration, oxide of iron remaining in solution entirely free from lime; this is precipitated, in the ordinary manner, by ammonia.—*Comptes Rendus*, Nov. 17, 1845.

---

#### REACTION FOR THE DISCOVERY OF SULPHUROUS ACID.

BY M. HEINTZ.

The substance to be examined, dissolved in water or hydrochloric acid, is to be heated with a solution of protochloride of tin in dilute hydrochloric acid to ebullition. If the liquid contains much sulphurous acid, sulphuret of tin is precipitated; but if the quantity be small, no precipitation occurs; the liquid becomes yellow and exhales the odour of sulphuretted hydrogen. It is then requisite only to add a few drops of solution of sulphate of copper to obtain an immediate precipitate of the brown sulphuret of this metal.

This method of detecting sulphurous acid, it will be observed, is merely a modification of that proposed more than fifty years ago by Pelletier, and since recommended by M. Gerard.

It is preferable to the process of MM. Fordos and Gelis, which is based on the formation of sulphuretted hydrogen by the contact of metallic zinc and sulphurous gas, inasmuch as it does not require the use of an apparatus to disengage the gas.—*Journ. de Pharm. et de Ch.*, Janvier 1846.

## ANALYSIS OF THE MOLARES OF A FOSSIL RHINOCEROS.

M. E. I. Meyer, by employing the process of Wöhler, found that besides phosphate and a little carbonate of lime, these teeth contained 2·10 per cent. of fluor.

## EXPERIMENTS ON THE YOLK OF EGGS. BY M. GOBLEY.

The author remarks, that a German chemist of the name of John was the first who carefully examined the yolk of the egg, the chemists who preceded him regarding it merely as consisting of water, albumen, oil, gelatine and colouring matter. John concluded from his experiments, published in 1811, that the yolk of egg was composed of water, a sweet yellow oil, traces of free acid, which he presumed to be the phosphoric, a small quantity of reddish-brown matter, soluble in æther and in alcohol, gelatin, much of a modified albuminous substance, and sulphur.

In 1825, Prout found the yolk to be composed of 54 water, 17 albumen, and 29 oil; and that it contained besides sulphur, phosphorus, the chlorides of sodium and potassium, the carbonates of potash and soda, lime and magnesia, partly in the state of carbonates.

Chevreul was of opinion that the orange colouring matter of the yolk was due to the combination of two colouring principles, one yellow, approximating that of the bile, and the other red, resembling that of the blood.

Lastly, in 1829, M. Lecanu discovered in the oil of the egg, a fat, crystallizable, unsaponifiable matter, which he considered to be cholestrine.

Such, says M. Gobley, was the state of our knowledge respecting the yolk of the egg when he began his experiments; he states that the substances which he obtained from the yolk are,—

1. Water.
2. Albuminous matter or vitelline.
3. Oleine.
4. Margarine.
5. Cholesterine.
6. Margaric acid.
7. Oleic acid.
8. A peculiar acid containing phosphorus, which is in fact phosphoglyceric acid.
9. Lactic acid and extract of meat.
10. Various salts, as chloride of sodium, chloride of potassium, hydrochlorate of ammonia, sulphate of potash, phosphate of lime, and phosphate of magnesia.
11. Yellow and red colouring matter.
12. Azotized organic matter, which does not appear to be albumen.

The oleic, margaric, and phosphoglyceric acids appear, in the author's opinion, to be combined with ammonia.

In the opinion of Berzelius, the yolk of egg contains some volatile fatty acids, on account of the facility with which the yolk becomes rancid; M. Gobley has not been able to discover them, nor any gelatine, and sulphur was met with only in the albuminous matter.—*Journ. de Pharm. et de Ch.*, Janvier 1846.

METEOROLOGICAL OBSERVATIONS FOR DEC. 1845.

*Chiswick*.—December 1. Rain: cloudy: clear. 2. Clear and fine: heavy rain. 3. Overcast: showery: clear. 4. Clear: fine: heavy rain. 5—7. Clear: frosty. 8. Sharp frost: overcast: drizzly. 9. Fine. 10. Clear. 11. Cloudy: clear and windy at night. 12. Overcast: fine: clear. 13. Frosty and foggy: cloudy. 14. Foggy: hazy: drizzly. 15. Rain: fine. 16. Fine. 17. Overcast: slight drizzle. 18. Foggy: rain. 19. Densely and uniformly overcast: rain. 20. Clear: dark clouds, with rainbow. 21. Boisterous and densely clouded: clear and frosty at night. 22. Densely overcast: sleet: showery: very boisterous at night. 23. Cloudy and boisterous at night. 24. Cloudless, with bright sun. 25. Hazy: thick fog at night. 26. Cloudy. 27. Clear: fine: overcast. 28. Boisterous, with rain: clear. 29. Frosty: overcast. 30. Overcast: clear. 31. Very fine: heavy rain and boisterous at night.—Mean temperature of the month  $0^{\circ} \cdot 4$  above the average.

*Boston*.—Dec. 1. Cloudy: rain early A.M. 2. Fine: rain P.M. 3. Fine. 4. Fine: rain P.M. 5—7. Fine. 8. Fine: rain P.M. 9. Cloudy: stormy P.M. 10. Fine. 11. Stormy: stormy night. 12. Cloudy: rain early A.M. 13, 14. Fine. 15. Stormy. 16. Cloudy. 17. Cloudy: rain P.M. 18. Rain: rain early A.M.: rain all day. 19. Cloudy. 20. Cloudy: rain early A.M. 21. Windy: rain early A.M. 22. Windy and showery. 23. Stormy. 24. Fine. 25. Rain: rain early A.M. 26. Cloudy: rain P.M. 27. Fine. 28. Rain: rain early A.M. 29. Fine: rain P.M. 30. Windy: stormy P.M. 31. Fine.

*Sandwick Manse, Orkney*.—Dec. 1. Showers: sleet-showers. 2. Showers: sleet: clear: aurora borealis very brilliant. 3. Fine: clear: aurora borealis very brilliant. 4. Showers: hail: cloudy. 5, 6. Rain: cloudy. 7. Clear frost: clear. 8. Bright: cloudy. 9. Showers. 10. Cloudy: rain. 11. Showers. 12. Cloudy. 13. Cloudy: showers. 14. Rain. 15. Sleet-showers: rain. 16. Sleet-showers: showers. 17. Frost: cloudy: clear frost. 18. Frost: cloudy: snow-showers. 19. Showers: clear frost. 20. Frost: cloudy: sleet-showers. 21. Frost: bright: cloudy: thaw. 22. Showers. 23. Showers: clear. 24. Cloudy: showers. 25. Showers: cloudy. 26. Showers. 27. Snow-showers: sleet-showers. 28. Snow: frost. 29. Rain. 30. Showers: clear frost. 31. Cloudy: rain.

*Applegarth Manse, Dumfries-shire*.—Dec. 1, 2. Showers. 3. Showers of snow. 4. Frost: rain P.M. 5. Very heavy rain. 6. Showers. 7. Fair and fine: slight frost. 8. Frost: rain P.M. 9. Fine A.M.: rain P.M. 10. Fair, but damp. 11. Fair and clear: frost. 12. Frost. 13. Frost, hard. 14. Very wet P.M.: frost A.M. 15, 16. Heavy showers. 17. Fine A.M.: shower P.M. 18. Fine A.M.: frost P.M. 19. Frost A.M.: rain P.M. 20. Frost A.M. 21. Frost: clear. 22. Heavy showers. 23. Slight frost. 24. Frost A.M.: shower P.M. 25. Fine. 26. Heavy rain all day. 27. Heavy showers. 28. Fair and fine. 29. Heavy rain: frost. 30. Heavy rain. 31. Frost A.M.: rain P.M.

Mean temperature of the month *	39 $^{\circ}$ ·5
Mean temperature of Dec. 1844	33 ·8
Mean temperature of Dec. for 23 years	38 ·3
Mean rain in Dec. for 18 years	3 inches.

\* It would be worth while for the meteorological correspondents to note the particulars here stated in their reports.



LONDON, EDINBURGH AND DUBLIN  
**PHILOSOPHICAL MAGAZINE**  
AND  
**JOURNAL OF SCIENCE.**

[THIRD SERIES.]

MARCH 1846.

XXX. *On the Determination of the Temperature and Conducting Power of Solid Bodies.* By CHR. LANGBERG of *Christiania*.\*.

NOTWITHSTANDING the important progress which has been made in the mathematical theory of the phænomena of heat by the analytical researches of Fourier, Poisson and others, it is not to be denied that the influence which these have exerted upon the extension of our physical knowledge of the phænomena is very limited, and that only a few of the results obtained from mathematical theory have been demonstrated and proved by experiment. The reason of this is in a great measure owing to the want of accurate modes of ascertaining changes of temperature in solid bodies, without militating against too many of the conditions required by the mathematical theory.

Thus we are taught by mathematical analysis that one of the most important elements in the theory of heat, namely the conducting power of solids, can be ascertained by placing in connexion with a constant source of heat the end of a long, thin, homogeneous, cylindrical or prismatical rod, composed of the substance to be examined, and observing the temperature of this rod at different distances from the heated end; the difference between the observed temperatures on the rod and that of the surrounding air decreases in geometrical progression when the points of observation are at equal distances from each other.

For establishing these laws experiments have been instituted by Biot †, and more lately by Despretz ‡; those of the latter

\* Being an abstract from Poggendorff's *Annalen*, 1845, No. 9; communicated by Dr. Ronalds.

† *Traité de Physique*, tom. iv. p. 670.

‡ *Annales de Chimie et de Physique*, tom. xxxvi. p. 422. *Traité Élémentaire de Physique*, p. 210.

however, as far as they have been made public, appear to me to prove exactly the converse of what they are intended to do, as the temperatures observed decrease in a much quicker ratio than the geometrical progression would require, and the difference between the calculated and observed values is too constant to admit of the supposition that it is solely due to errors of observation. One source of the difference may probably arise from the Newtonian law of cooling having been made the basis upon which to found the mathematical deduction of the law in question. According to Newton's law, a heated body cools with a rapidity proportional to the degree of temperature to which it has been raised above that of the air surrounding it, and applies with correctness only to very slight differences of temperature. [In the experiments alluded to above, this difference amounts to  $60^{\circ}$  or  $70^{\circ}$  C.] Again, it is presumed that the power of conduction remains unchanged at different temperatures, which is certainly not probable; and the theory further requires that the heated rod should be infinitely thin, or at least so thin that its temperature in every part of a normal section should be exactly the same.

Despretz used in his experiments prismatical rods, the square section of which was 21 millimetres in breadth; holes were bored in these at 10 millimetres distance from each other, 6 millimetres in diameter, and 14 millimetres deep. When the rod had been brought into a horizontal position, these holes were filled with mercury, and in each the bulb of a thermometer was placed, the temperature of which, when it had become stationary, was considered as that of the section of the rod passing through the middle of the hole. As the breadth of the holes amounted to nearly one-third of the whole breadth of the rod, there is reason to fear that these large and frequent interruptions in the continuity of the rod might cause a considerable obstruction to the progress and distribution of the heat. The results show that the method employed in the experiments fulfilled very imperfectly the conditions required by the theory, and it remains therefore still uncertain whether the variations from the theoretical law which were observed are to be attributed to an inaccurate method of observation, or to an error in theory.

The importance of the law, forming as it does the basis of the mathematical theory of the phænomena of heat, as well as from its application for determining the conducting power of solid bodies, appeared to me sufficiently great, and induced me to seek a mode of observation not subject to the objections which I have raised above. The first requisite was therefore a mode of ascertaining correctly such slight differences be-



tween the temperature of the rod and the surrounding air as would come with accuracy within the scope of the Newtonian law of cooling, and this must be done with rods of indefinitely small diameter, having their continuity unimpaired by any holes bored into their substance. The thermo-electrical battery appeared to me to be an instrument admirably adapted for this purpose; and arranged in the manner which I shall presently describe, I hope that it will be found a much more accurate measure for observing uncombined heat in solid bodies than any other means that have yet been applied to that purpose.

The experiments were performed in the laboratory of Professor Magnus, who was kind enough to lend me the necessary apparatus, and to whose friendly guidance the successful result of the experiments is chiefly to be attributed.

I found by several preliminary experiments, that the same divergence in the needle of the multiplier was always attained when the end of a thermo-electrical battery, consisting of but few alternations, was connected in a similar manner with a body of constant temperature, and pressed against it with the same amount of force. It was always 2 to  $2\frac{1}{2}$  minutes before the needle of the multiplier became stationary; the connection might then be continued for an indefinite time without perceptibly affecting the position of the needle. In order to secure perfectly uniform contact, which is hardly possible with a battery composed of many alternations, I had one constructed of only two elements, bismuth and antimony, having therefore but one soldered point of contact at each end. The ends were filed off presenting facets, so that each presented a rectangular surface of 1.7 millimetre in length and 0.7 millimetre in breadth. The whole length of the bars was 36.3 millimetres; each bar was very thin, being 1.7 millimetre in width and 1.0 millimetre thick.

On a strong horizontal board, on which divisions had been marked, three uprights were erected, bearing each a forked arm in which were fixed two perpendicular glass rods drawn out to a point and placed opposite to each other, between which the metallic bars to be examined were clamped parallel to the horizontal divided board, and at about 24 centimetres above it; a fourth upright at the end of the board served to fix securely, with the help of a screw, the cool end of the rod during the experiment. To obtain for a length of time a uniform source of heat, hot water was used: the end of the rod to be heated, passed through a cork into the water by an opening made in the boiling vessel below the surface of the water.

By means of two double polished brass screens, through

holes in the centres of which the rod passed, the battery and that part of the rod to be examined were effectually protected from the radiant heat of the boiling vessel. The support of the battery was screwed to a sledge which could be moved along the edge of the horizontal divided board in a direction parallel with the metallic rod, by which means the relative distances of the points to be examined on the rod and their temperatures could be easily ascertained. In order that the battery should press with equal force each time against the rod, a spiral spring was so placed in the support of the battery as to force it upwards against the under side of the rod; or it could be brought into contact with the rod in a direction at right angles to it.

The observations were made in the following manner:— When the rod had attained a constant temperature, which seldom occurred until about  $2\frac{1}{2}$  to 3 hours after the commencement of the experiment, the sledge bearing the battery was so placed that the upper end of the battery bore perpendicularly upon the under side of the rod. The spiral spring was then allowed to force the end against that part of the rod the temperature of which was to be examined: the needle of the multiplier diverged immediately. I waited generally about two minutes to allow the needle to come to rest, and having noted the divergence, removed the battery. After each observation I allowed four minutes to elapse before the battery was again placed in contact with the rod, partly that the needle of the multiplier might return to  $0^\circ$ , and partly that the equilibrium of temperature in the rod, which might possibly have been disturbed by its contact with the battery, might again be restored. This latter precaution was however needless, as observations made upon the same part of the rod immediately the one after the other, were found to give the same deviations in the needle as when a space of time was allowed to elapse between each observation.

Thus far we have given very nearly the author's own words; but as our space will not allow us to follow him through each experiment, we shall here briefly add some of the precautions which were taken to avoid error, and then give the results to which the experiments have led.

To ascertain whether the battery itself, after being for some time in contact with the rod, might by becoming warm no longer indicate with correctness the difference of temperature between the rod and the surrounding air, it was left in connexion with the heated rod for three quarters of an hour, but during the whole of that time the divergence of the needle of the multiplier scarcely changed. The circumstance that the needle

of the multiplier seldom returned to precisely the point from which it had been deflected, sometimes becoming stationary a little to the right, sometimes to the left of its original position, the cause of which deviation could not be traced, would give rise to a slight error, perhaps 0·1 to 0·2 of a degree, upon the scale of the multiplier, which had been divided in the manner recommended by Melloni.

The divisions upon this scale not corresponding exactly with the temperatures they are intended to indicate, might also lead to a slight error; and although the observations were always made with the aid of a magnifier, still it is possible that from 0·1 to 0·2 of a degree should escape observation.

These three sources of error, which arose chiefly from the author's having used an imperfect instrument, added to another more important one, the effect, namely, of currents of air in the room cooling one part of the rod under examination more than another, the author calculates will not amount to more than 0·4° C. in the final result of any experiment. In fact it never actually amounted to so much.

The metals used in the experiments were copper, steel, tin and lead; they were all drawn out into cylindrical wires or thin rods, and their length was such that even in the middle of the rod no effect of the heating body was perceptible. The copper alone, being the best conductor, was slightly affected through its whole length.

The object of the experiments was not so much to determine the conducting power of the metals employed, as to submit the analytical law to the test of experiment; the metallic surface of the rods was therefore not protected, and remained unimpaired in the three first metals; the lead wire however soon became covered with a layer of oxide, which increased in thickness every time it was heated.

The results at which the author arrived were principally the following:—

The law of Biot,—that in a very thin, long metallic rod, one end of which is kept at a constant temperature above that of the surrounding air, after equilibrium of temperature has been established, the excess of temperature in any part of the rod above that of the surrounding air, decreases in geometrical progression as the part examined is removed by equal distances from the heated end—is not generally confirmed by the author's experiments, and is only true for most of the metals in the case of a very small excess of temperature. Among the metals examined copper was the only one for which the law held good, at least when the excess of temperature amounted to 30° C.

With tin, the law no longer applied when the excess was

4° C.; with steel when it amounted to 2° or 3°; and lastly, with lead, when 1° of difference existed, it was not accurate.

2. The reason for this want of accordance between the observation and the mathematical law is, that in establishing the latter the outward and inward power of conduction of bodies was considered as independent of the temperature. If these be considered as functions of the temperature, a proximate formula for the distribution of heat in the rod may be established, which would very nearly agree with my observations.

3. The conducting powers of bodies established by former philosophers with the aid of Biot's law are consequently incorrect, and can only be considered as partial approximations to the truth.

4. The constant coefficient for the conducting power with a difference of temperature equal 0, is therefore only to be determined in this manner; either by using Biot's law and observing the distribution of heat in the rod with very small differences of temperature, or more correctly, by ascertaining its value, as was done in the author's experiments by means of Poisson's formula.

5. That the method employed in the observations is accurate, and that the thermo-electrical battery will become in the hands of natural philosophers a more correct means of ascertaining the temperature of the surfaces of bodies than any other, and may be used in cases where the common thermometers cannot be employed.

XXXI. *On the Causes of the Semi-diurnal Fluctuations of the Barometer.* By THOMAS HOPKINS, Esq.\*

THAT the non-condensable gases and the aqueous vapour of the atmosphere, when it is at rest, press on the mercury of the barometer independently of each other, and constitute the general atmospheric pressure, is evident from their known laws of diffusion and independent existence while diffused through each other.

But that the facts and reasonings, commonly adduced, resting on those circumstances, together with the daily alterations of thermometric temperature, account for the two risings and the two fallings of the barometer, as is contended by some parties, cannot be admitted.

In certain parts, such as Canada, of which an account has been recently given by Colonel Sabine, the influence of the causes named may be sufficient to account for a considerable

\* Read to the Literary and Philosophical Society of Manchester, Dec. 30, 1845, and communicated by the Author.

portion of the semi-diurnal movements of the barometer which occur in that country; but these causes are not sufficient to produce the diurnal fluctuations in other places, such as Bombay, Calcutta and La Guayra. And there can be little doubt that the real causes, whatever they may be, which give rise to the double undulations in these tropical parts, produce them in places where they are less extensive, although the operation of the causes in the latter places may be weaker and more difficult to trace.

I have shown in my "Atmospheric Changes" that there is no reason to believe that the daily warming of the atmospheric gases by the direct influence of the sun produces any appreciable alteration in their pressure on the mercury of the barometer, as the effect of that warming on the whole column in the locality is so small, as to prevent much disturbance of atmospheric pressure; yet great influence has been attributed to solar heating near the surface in producing the semi-diurnal fluctuations that take place.

Colonel Sabine, in his Report on the Meteorology of Toronto at the meeting of the British Association in 1844, gives an explanation of the daily oscillations. He says, "As the temperature of the day increases, the earth becomes warmed and imparts heat to the air in contact with it, and causes it to ascend. The column of air over the place of observation thus warmed rises, and a portion of it diffuses itself in the higher regions of the atmosphere, where the temperature at the surface is less. Hence the statical pressure of the column is diminished. On the other hand, as the temperature falls, the column contracts, and receives in its turn a portion of air which passes over in the higher regions from spaces where a higher temperature prevails; and thus the statical pressure is augmented."

In the Athenæum of July 5, 1845, the Colonel is represented as having said at the then recent meeting of the British Association, that in Dr. Buist's Meteorological Report from Bombay, "the explanations thereby afforded of the diurnal variations of the *gaseous* pressure at Bombay, which, although at first sight more complex than at the stations of Toronto, Prague or Greenwich, he conceives to be equally traceable to variations of temperatures." Colonel Sabine therefore, after having examined Dr. Buist's meteorological registers, retains the opinion that the semi-diurnal alterations of the *gaseous* pressure are produced by alterations of temperature, as that temperature is shown by the thermometer.

As I propose to examine this theory and to compare it with another, it will be convenient to designate the two by distinct

names. I shall therefore call the Colonel's "*the temperature theory*," and the other, "*the condensation theory*." Both of these rest on alterations of temperature; but the former depends on the temperature found by thermometric measurement near the earth's surface, and the latter on the temperature which must be produced by condensation of vapour in a higher part of the atmosphere, of which we have no direct measure.

The semi-diurnal fluctuations of the barometer are the greatest within the tropics; and as details of those at Bombay have not yet been published, we will proceed to examine accounts furnished by Kaemtz in his valuable work on Meteorology. In page 248 of that work we have the following tables of the hourly heights of the barometer:—

TABLE I.

Mean height of the barometer expressed in millimetres for all hours, and in different places.

Places . . .	Gt. Ocean.	Cumana.	La Guayra.	Calcutta.	Padua.	Halle.	Abo.	Petersburg.
Latitude . .	0° 0'	10° 23' N.	10° 36' N.	22° 35' N.	45° 24' N.	54° 29' N.	60° 57' N.	59° 66' N.
Observers .	Horner.	Humboldt.	Boussingault.	Balfour.	Ciminello.	Kaemtz.	Hallstroem.	Kupffer.
Noon	752·35	756·57	759·41	759·61	757·02	753·29	759·31	759·47
1	751·87	755·99	758·91	759·22	756·85	753·11	759·29	
2	751·55	755·47	758·41	758·39	756·67	752·99	759·27	759·38
3	751·15	755·14	758·12	758·12	756·54	752·89	759·25	
4	751·02	754·96	758·05	757·91	756·47	752·84	759·25	759·32
5	751·31	755·14	758·10	757·93	756·46	752·86	759·27	
6	751·71	755·41	758·40	758·01	756·50	752·91	759·29	759·31
7	751·93	755·81	758·90	758·02	756·63	753·02	759·34	
8	752·35	756·21	759·19	758·54	756·79	753·14	759·39	759·32
9	752·74	756·59	759·69	759·24	756·92	753·24	759·44	
10	752·85	756·87	759·93	759·33	757·02	753·31	759·47	759·36
11	752·86	757·15	759·98	759·09	757·02	753·29	759·47	
Midnight	752·47	756·86	759·64	758·80	757·01	753·23	759·41	759·35
13	752·20	756·53	759·34	758·62	756·90	753·14	759·33	
14	751·77	756·21	759·05	758·57	756·84	753·05	759·24	759·32
15	751·63	755·89.	758·81	758·49	756·78	752·99	759·14	
16	751·32	755·66	758·68	758·47	756·74	752·99	759·07	759·32
17	751·65	755·79	758·85	758·44	756·75	753·34	759·03	
18	751·95	756·18	759·32	758·68	756·79	753·12	759·04	759·39
19	752·84	758·58	759·94	759·16	756·89	753·24	759·08	
20	752·95	756·98	760·50	759·88	757·01	753·37	759·15	759·49
21	753·16	757·31	759·63	760·11	757·08	753·44	759·21	
22	753·15	757·32	760·50	759·19	757·14	753·46	759·29	759·51
23	752·80	757·01	759·99	759·09	757·07	753·40	759·32	

From an examination of this table, it will be seen that the fluctuations are the greatest within the tropics, and they diminish, though not invariably, with the increase of latitude.

The first column exhibits the fluctuations at the equator in the great ocean. The range extends beyond two millimetres, and the descent from 9 in the morning till 4 in the afternoon gives the whole extent of the range.

The two next columns show the alterations at Cumana and La Guayra, both above  $10^{\circ}$  north latitude, and the ranges are nearly equal to that at the equator.

The fourth column shows the changes at Calcutta to be nearly as great as in the preceding places, but both this and the La Guayra columns exhibit singular irregularities in the earlier parts of the mid-day descents.

In the Padua column,  $45^{\circ}$  north, the fluctuation is much reduced in the extent of its range, but retains the same general character.

In Halle, in latitude  $54^{\circ}$ , the alterations do not differ materially from those at Padua.

The changes are very small in Abo and Petersburg, and in the former place the second rise attains a greater height than the first.

To these it is desirable that we should add the following table (p. 170) of the height of the dry- and wet-bulb thermometers, and the difference between the two,—with the dew-point and the height of the barometer at Plymouth for three years, as furnished by Mr. S. Harris, and published in the Ninth Report of the British Association (p. 167).

In all these places the temperature shows only a single fluctuation, such as is seen in the table of the thermometer at Plymouth, namely one rise generally from about 5 A.M. to 1 or 2 P.M., and one fall from that time until 5 the following morning. Now, if the temperature of the atmosphere, as marked by the thermometer, caused the diurnal fluctuations in the way supposed, we ought to have in all these places one undulation in the twenty-four hours instead of two,—the rise of temperature causing a decline of the barometer during the hotter part of the day, and the fall of temperature producing a rise of the barometer in the colder part. Yet Colonel Sabine himself says that at Bombay, where there is only one rise and one fall of temperature, there are two risings and two fallings of the barometer! And these movements of the barometer take place not only when that instrument is taken as the measure of the whole pressure of the atmosphere, but also when the vapour pressure is deducted, and the mercury of the barometer is taken as the measure of the gaseous pressure alone. These facts are opposed to, and are irreconcilable with, the temperature theory.

TABLE II.

Table of the heights of the dry- and wet-bulb thermometers, and the difference between the two, together with the dew-point and height of the barometer at Plymouth for three years.

Hour.	Thermo- meter.	Wet-bulb thermo- meter.	Difference.	Dew- point.	Barometer.
1 A.M.	47.52	46.20	1.32	45.00	29.8017
2	47.33	46.03	1.30	44.75	29.7993
3	47.11	45.92	1.19	44.75	29.7944
4	47.00	45.66	1.34	44.25	29.7928
5	46.98	45.77	1.21	44.75	29.7928
6	47.41	46.01	1.40	44.50	29.7960
7	48.44	46.83	1.61	45.25	29.8002
8	49.68	47.51	2.17	45.00	29.8032
9	51.30	48.50	2.80	45.26	29.8048
10	52.84	49.45	3.39	46.25	29.8061
11	53.00	50.02	3.88	46.75	29.8045
12	54.51	50.40	4.14	46.75	29.8002
1 P.M.	55.83	50.55	4.28	46.75	29.7957
2	54.77	50.44	4.33	46.75	29.7922
3	54.25	50.24	4.01	46.75	29.7908
4	53.45	49.80	3.65	46.75	29.7895
5	52.27	49.06	3.21	46.25	29.7938
6	51.24	48.46	2.78	46.00	29.7970
7	50.28	47.90	2.38	45.75	29.8019
8	49.44	47.51	1.93	45.75	29.8061
9	48.83	47.17	1.66	45.60	29.8094
10	48.48	46.93	1.55	45.60	29.8099
11	48.10	46.66	1.44	45.00	29.8092
12	47.80	46.43	1.37	45.00	29.8065
Mean ..	50.32	47.89	2.43	45.60	29.7999

As however the aqueous vapour of the atmosphere presses on the mercury of the barometer separately and independently, it has been attempted to be shown that the variable pressure of the vapour arising from difference in the quantity in the atmosphere at different periods of the day, combined with change of the gaseous pressure resulting from alteration of surface temperature, and that the two causes acting together produced the double undulation of the barometer; to this view therefore we will direct our attention.

The temperature near the surface of the earth at Plymouth, as well as at the other places, rises from about 5 in the morning till about 2 in the afternoon; and when the wet bulb, as well as the dry thermometer, is used, as it was at Plymouth, it is seen that the temperature of the latter rises more than that of the former, or of the dew-point, and evaporation must consequently become progressively more active;



there must therefore be successively more water evaporated and thrown into the atmosphere to be added to its weight. And according to the temperature theory, this water, now converted into vapour, must, up to say 10 o'clock, press with sufficient force on the mercury to counteract the lightening influence of the rising temperature, as during that time the barometer rises.

From 10 until 1 o'clock, as the temperature rises still higher, as compared with the wet-bulb thermometer and the dew-point, evaporation must go on increasing, and the increase of vapour pressure ought to continue; but it appears from the table not to do so, as the mercury of the barometer falls instead of continuing to rise; we have therefore to try to ascertain what can be the cause of this fall, while additional vapour is passing into the atmosphere.

Those who advance the temperature theory, say that the fall of the barometer is caused by the increasing temperature of the atmosphere produced by the action of the sun on the surface of the earth, and the air near to it; and they must maintain that this increase is sufficient, not only to lighten the atmosphere enough to cause the fall of the barometer, but also in addition to counteract the influence of the increased vapour pressure. Now at Plymouth the temperature rises from 5 to 10 A.M. nearly  $6^{\circ}$ , and may be supposed to lighten the atmosphere to a certain extent; at the same time evaporation throws vapour into the atmosphere. We are, however, required to suppose that the vapour produces so much greater effect by pressing on the mercury, than the heating of the atmosphere does in reducing atmospheric pressure, that the whole pressure becomes greater and the mercury rises. But after 10 o'clock the temperature continues to rise, but in a smaller degree, say nearly  $3^{\circ}$ , and vapour must be more abundantly thrown into the air, as is shown by the extent to which the wet-bulb thermometer is kept down; yet the barometer, instead of continuing to rise, suddenly turns and falls, and continues falling from 10 to 1 o'clock, the time of the highest temperature! So that according to this theory, from 5 to 10 o'clock, the sun heats the air nearly  $6^{\circ}$  and produces some vapour; and the two influences acting together cause the barometer to rise, but from 10 to 1 the sun heats the air about  $3^{\circ}$ , and must throw much additional vapour into the atmosphere; and then these two influences still acting together cause the barometer to fall! This is attributing opposite effects to the same causes, and must be presumed to be erroneous.

But let us examine the valuable Plymouth tables a little more minutely. The first column gives the temperature as

shown by the ordinary thermometer; the second, the temperature of the wet-bulb thermometer, as kept down by the cooling influence of evaporation; and the third gives the difference between the two first. Now as this difference arises from the extent of the evaporation, the numbers of the difference may be taken to express the force and amount of evaporation, and to indicate the additional vapour that is discharged into the atmosphere. This force or amount at 5 o'clock in the morning is  $1^{\circ}21$ , from which time it increases to  $3^{\circ}39$  at 10 o'clock. So that during this time, five hours, the increase in the force of evaporation is  $2^{\circ}18$ ; and this in the temperature theory must be held to be sufficient to overcome the lightening effect of a rise of  $5^{\circ}86$  of temperature, and *also* to raise the mercury of the barometer to the full extent of the morning rise! After this time, from 10 to 1 o'clock, the temperature rises further from  $52^{\circ}84$  to  $55^{\circ}83$  or  $2^{\circ}99$ ; and during the same period the force of evaporation increases  $1^{\circ}89$ , that is, from  $3^{\circ}39$  to  $4^{\circ}28$ . Thus we are required to believe, that from 5 to 10 in the morning,  $2^{\circ}18$  of evaporation overcame the lightening influence of  $5^{\circ}86$  of temperature, and in addition raised the mercury of the barometer; and from 10 to 1 in the day,  $1^{\circ}89$  of evaporation not only failed to overcome the lightening effect of  $2^{\circ}99$  of temperature, but allowed this relatively small amount of temperature to produce the further result of a fall of the mercury of the barometer. Or put in a tabular form, say that from

5 to 10 o'clock,  $5^{\circ}86$  of temperature and  $2^{\circ}18$  of evaporation caused a rise.  
 10 to 1 o'clock,  $2^{\circ}99$  of temperature and  $1^{\circ}89$  of evaporation caused a fall!

That is, where temperature, the influence which lightens the atmosphere, is relatively great and should cause a fall, the mercury of the barometer rises; and where the influence of temperature is relatively small and should cause the vapour to produce a rise, the mercury falls! This must be erroneous.

In the same place, at Plymouth, from 1 o'clock until 4 P.M., as may be seen in the table, the temperature falls; and as far as that temperature acted the atmosphere would of course become heavier. At the same time evaporation shows vapour is passing into the atmosphere; it ought therefore to follow that the barometer should rise, and considerably too, through the operation at the same time of both the causes which are supposed to contribute to the production of a rise. But the barometer does not rise; on the contrary, it falls, and continues falling until 4 o'clock. These facts and reasonings prove that neither the daily variations of surface temperature, nor the different amounts of vapour pressure, nor both taken to-

gether, are adequate to the production of the fall of the barometer from 10 to 4 o'clock in the day.

And if we proceed with our inquiries into the next period of six hours, that is, from 4 to 10 P.M., we meet with facts that do not harmonize with the temperature theory. During the whole of this time, it is true the temperature falls and the barometer rises: but the vapour pressure must have diminished according to the temperature theory, as the dew-point, the measure of vapour pressure, falls; and the lowering of the dew-point after 4 o'clock showed that vapour was then condensing in the lower part of the atmosphere. So that here it becomes necessary to suppose that the atmosphere cools enough, not only to raise the barometer to the full extent of its daily range, but also to counteract the reduction which takes place at the same time in the vapour pressure. Again, from 10 at night, although the atmosphere continued to cool, the barometer did not continue to rise, but once more fell, which fall is attributed to a diminution of vapour pressure. Thus from 4 to 10 in the afternoon and evening, cooling the atmosphere is represented as more powerful than reduction of vapour pressure; and from 10 in the evening to 4 in the morning, reduction of vapour pressure is supposed to be more powerful than cooling the atmosphere. The two forces, we are required to believe, do not merely neutralize each other, but each in its turn exercises a paramount influence, and for the time determines an absolute rise or a fall of the barometer; and this we are called upon to admit without any satisfactory or even plausible evidence being adduced to prove it.

What has been here advanced applies with the greatest force to the semi-diurnal fluctuations in atmospheric pressure which take place within the tropics. Aqueous vapour exists in the atmosphere in larger proportions in that part of the world than it does in higher latitudes; and it is to the daily condensation of that vapour in the atmosphere, and its subsequent evaporation there, that we are really to attribute the great deviation of the movements of atmospheric pressure from the daily march of temperature. If no vapour existed in the atmosphere, the alteration of pressure would be very little, and it would be the reverse of temperature. As the atmosphere became warmer, the pressure would be less; as it became colder, the pressure would be more. And the hourly variation in the quantities of vapour actually found in the atmosphere which arises from alteration of surface temperature, only introduces another element of pressure into the inquiry, which is simple in its character,—the vapour increasing or diminishing with an increase or diminution of temperature.

If the two were equal while acting in opposite directions, they would balance each other. But the separate action of these two causes cannot produce such a double undulation of the mercury of the barometer as that which occurs daily in the tropical regions and at Plymouth.

The double undulation which takes place may be thus accounted for. When the sun acts with force on the surface of the earth in the morning, it heats that surface, and the air near it; increases evaporation of moisture from wet surfaces, and sends forth vapour, which presses on the mercury of the barometer and causes it to rise. The lower part of the atmosphere being heated also rises at the same time, probably in separate vertical streams, until it reaches a height where its expansion and consequent cooling is sufficient to condense a part of the vapour which it contains. A cloud is then formed, and the heat which has been evolved in the condensation of the vapour makes the cloud lighter than the adjoining air. The vapour in the upper part of the air being thus removed by conversion into water, no longer presses as vapour, or with the same force on that below; and the lower vapour consequently rises more freely to the height of the cloud. Both the air and vapour are also (speaking in popular language) drawn up by the ascending cloud, and fresh air flows in from adjoining low levels, forming what in some parts is called the sea breeze. Cloud more or less thick is now formed, more heat is liberated, and a larger mass of air heated, which being forced upwards expands and makes the whole atmospheric column lighter, and reduces the pressure on the surface below. Under ordinary circumstances this process proceeds while the sun acts with considerable power on the surface of the earth, which is generally from 10 A.M. to 4 P.M., when day-cloud ceases to form. In this way, from 10 in the morning till 4 in the afternoon, the barometer is caused to fall, through the condensation of vapour in the upper part of the atmosphere making the column of air warmer and lighter. But now as vapour no longer ascends, cloud ceases to form, but that cloud which had been formed remains suspended in the air, where it begins to cool from the influence of evaporation of the particles of water that form the cloud. When it cools sufficiently, it becomes heavier and sinks, and additional air flows towards and over it, increasing the weight of the whole column in the locality and causing the barometer to rise. By 10 the heavy air produced by cloud evaporation has partly descended and diffused itself on the surface of the earth, forming what is called the land breeze; and during the same time the cold of the surface condenses some of the va-

pour into dew, when the atmosphere becomes somewhat lighter up to about 4 or 5 in the morning.

As we proceed from the equator towards higher latitudes, we find less vapour in the atmosphere, and its influence on atmospheric pressure is less marked. At Padua the fall of the barometer from 10 to 4 in the day is not much more than one-fourth the extent that it is at the equator, and at St. Petersburg it is very small. In situations where there is not sufficient vapour in the atmosphere to form any daily cloud, it is to be presumed that if a barometrical registration were to be made, there would be no double movement exhibited showing a fall from 10 A.M. to 4 P.M., and a rise from 4 to 10 P.M., because there would be no condensation and warming to produce the former, nor evaporation and cooling to cause the latter.

The heating effects of condensing vapours may however be traced even in comparatively dry latitudes, such as that of Toronto, as shown in Col. Sabine's report to the British Association in 1844. There was no fall of the barometer at that place from 4 to 10 in the morning, although the temperature had risen from  $39^{\circ}20$  to  $46^{\circ}35$ , above  $7^{\circ}$ ; but in the middle of the day, from 10 to 4, with an increase of temperature from  $46^{\circ}35$  to  $50^{\circ}55$ , being only  $4^{\circ}20$ , the gaseous as well as the general atmospheric pressure was materially reduced! notwithstanding that the increase in the quantity of vapour during this time must have been as great as it was in the preceding period; and if this increased quantity had remained in the atmosphere, its pressure must have been added to that which previously existed. We are then obliged to suppose that the reduction of the pressure which took place immediately after 10 o'clock, arose from a cause which came into operation at that time; and that cause it is contended can be found only in the heating of the atmosphere by the condensation of vapour.

The great defect of the temperature theory is, that it fails to account for the fall of the barometer from 10 A.M. to 4 P.M., and its subsequent rise from 4 to 10 P.M., though this is the oscillation for which we have particularly to account; whilst the theory here maintained points out the cause of these, as well as the other diurnal, and also of the casual movements of the barometer. We are therefore at liberty to conclude that the semi-diurnal fluctuations of the barometer can be accounted for only on the condensation theory.

XXXII. *On the Principles to be applied in explaining the Aberration of Light.* By the Rev. J. CHALLIS, M.A., Plumian Professor of Astronomy in the University of Cambridge\*.

THE aberration of light having been brought before the notice of the readers of this Journal by several recent communications, I am unwilling to let the subject drop without saying a few more words respecting the principles to be applied in the explanation of the phænomenon, which possibly may appear, after all that has been said, to be involved in uncertainty. I propose to answer the question, Is the aberration of light to be attributed to known causes, or must we, to explain it, have recourse to hypothesis?

The first attempts to explain aberration referred it to the combined effect of the motion of the earth and the temporary transmission of light, and accordingly proceeded on the principle of attributing it to known causes. It must, however, be admitted that every attempt to show *how* the observed effect resulted from these causes, what was the particular *modus operandi*, was unsatisfactory. Some idea appropriate to the subject was still wanting. This idea I consider that I have succeeded in supplying. I have argued, as had not been argued before, that because the direction of a celestial object is necessarily referred to the direction of a terrestrial object, light from the one as well as light from the other must be taken account of in considering the question of aberration. It is self-evident, that if at any instant two objects appear in the same direction, whatever course the light from the more distant may have taken before it reaches the nearer, it subsequently pursues a common course with light from the latter, and the two portions of light enter the eye at the given instant simultaneously. The direction in which the light comes is therefore judged to be the same as the direction at that instant of the nearer object from the eye. But during the interval the light takes to pass from the nearer or terrestrial object to the eye, this object is carried by the earth's motion away from the direction of the progression of light, and the two directions, at the time they are judged to be coincident, are in reality separated by a certain angle. This angle is aberration. I may refer to my communication in the February Number for a proof, which I venture to say is as cogent as any proof in the elements of geometry, that according to the principles just stated, an astronomical instrument employed to measure the *earth's way*, as it is called, would measure a smaller angle.

\* Communicated by the Author.

The difference, or aberration, is readily calculated from knowing by observations of the eclipses of Jupiter's satellites, the ratio of the earth's velocity to the velocity of light. Being so calculated the amount is found to be the same as the amount of aberration independently determined by astronomical observation. It follows from this accordance, not only that the aberration of light is entirely accounted for on these principles, but also, as a corollary, that the direction of the progression of light from a star, as it enters the eye, is the true direction of the star. Whether it be the star, or the terrestrial object to which it is referred, that is *seen* in its true place, is a curious question, not readily answered, and not in the least degree necessary to be answered in the present inquiry.

Sufficient reasons have now, I think, been adduced for coming to the conclusion, that the question I proposed to consider must receive the following categorical answer:—The aberration of light is entirely due to known causes, viz. the motion of the earth and the temporaneous transmission of light, and does not require for its explanation any hypothesis whatever.

What then becomes of the theories which have been framed to account for aberration on the hypothesis of certain motions of the æthereal medium? As explanations of aberration they can be of no value, it being an acknowledged principle in philosophy, that an hypothesis is not to be sought for to explain what may be explained by known causes. All that is left for the theorist to do, supposing, as it appears necessary to suppose, that the æther is in some way put in motion by the motion of the earth, is to show that *no* aberration results from such motion, the whole being attributable to the earth's motion. This problem I have considered in my two former communications, not because it was necessary to do so to complete the explanation of aberration, but with the view of removing an objection that might be raised against the undulatory theory of light. By taking account both of the light from the star and the light from the terrestrial object to which the star's direction is referred, I found that no aberration would result from the motion of the æther, provided it satisfied certain not improbable analytical conditions. A different conclusion would be arrived at by the same reasoning, if the light from the star, as is commonly done in treating of aberration, were alone considered.

With these remarks I dismiss the subject of aberration, having attained the object I had in view in taking it up, if I have succeeded in extricating the explanation of the phenomenon from hypothesis and conjecture, and placing it on its true basis.

XXXIII. *On the Cause of the Circulation of the Blood.* By JOHN WILLIAM DRAPER, M.D., *Professor of Chemistry in the University of New York.*

AMONG physiological problems there is none of greater interest, or of more importance in its relations to the well-being of man, than that which proposes to determine the true cause of the circulation of the blood, and the various other liquids which pass from one portion of living systems to another. Unquestionably one of the most important discoveries ever made by any physician was that of the route of the circulation by Harvey. The clearness with which he and his successors developed that doctrine not only fully established his views, but gave rise to a serious error which is scarcely removed in our times.

That error relates to the action of the heart. These earlier writers regarded the circulation of the blood as a hydraulic phenomenon, supposing that the heart simulated exactly the action of a pumping machine. It is now on all hands conceded that this organ discharges a very subsidiary duty. The whole vegetable creation, in which circulatory movements of liquids are actively carried on without any such central mechanism of impulsion; the numberless existing acardiac beings belonging to the animal world; the accomplishment of the systemic circulation of fishes without a heart; and the occurrence in the highest tribes, as in man, of special circulations which are isolated from the greater one, have all served to demonstrate to physiologists that they must look to other principles for the cause of these remarkable movements.

When we reflect how large a portion of the human family is destroyed by diseases dependent on derangements of the circulation, and to how great an extent the practice of medicine, as a scientific pursuit, must depend on just views of this important function, a natural philosopher can scarcely be more profitably employed than in attempting a solution of this problem.

I am persuaded that the phenomenon may be accounted for upon physical principles in a satisfactory manner; that we can co-ordinate together, and arrange as examples of one common law, the various forms of circulatory movements, whether they occur among vegetables or animals, among insects, or fishes, or mammals; and that the facts which we meet in derangements of these motions, or their cessation, as in fainting, coughing, and the different forms of disease, or such as take place after hanging, the inhalation of protoxide of nitrogen, or alcoholic drunkenness, or in that most remark-



able of all results, the restoration from death by drowning; in all these and many other such cases we can give the most felicitous explanation.

The principal facts which I design here to establish are,—

First. The systemic circulation is due to the de-oxidation of arterial blood.

Secondly. The pulmonary circulation is due to the oxidation of venous blood.

And, in conclusion, I shall offer some explanatory remarks on the phænomena of the coagulation of the blood.

Several physiologists have already made an approach to the doctrine which will be developed in this memoir. Among well-informed writers it is conceded that we must look to the relations between the blood and the tissues for the true cause of the circulation. Thus Dr. Alison attributes the effect to a “series of vital attractions and repulsions,” created by the operations to which the blood in the capillaries is subservient, an idea which Dr. Carpenter has rendered more explicit, by suggesting that these forces may not be “essentially different from those which are witnessed in Physics and Chemistry” (Carpenter’s *Human Physiology*, vol. ii. p. 417). But these views do not communicate a definite idea of the true mechanism of the motion, nor do they exhibit that phænomenon as clearly connected with well-known chemical changes occurring in living systems. Should it appear, as I shall endeavour to prove, that the circulation is a necessary result of the alternate oxidation and deoxidation of the blood, we exchange at once a loose and ill-defined conception for a precise and definite fact.

It will be perceived that I speak of the oxidation and de-oxidation of the blood as the great facts to be regarded, and leave out of consideration the spontaneous changes which that fluid itself undergoes; those minor effects which it impresses on the tissues, and those which they reciprocally impress on it. For the blood experiences in the systemic circulation an incessant change, discharging a double function. Its plasma serves for nutrition, its discs for the production of heat. But whilst the final function of the plasma and discs is different, there is an intimate relationship between them. It is from the plasma that the discs arise, and at its expense they grow. Moreover, the tissues themselves, in their metamorphoses, impress changes on the blood; the cells of which they are composed have an ephemeral existence, they dissolve, and the circulating fluid removes their remains and forms new ones in their stead.

I doubt very much whether animals obtain ready-formed

fibrine from the vegetable world. During the incubation of an egg we see this substance arising from albumen, and the analogy is probably continued in higher forms of existence. Neither is it by any means certain that fibrine exists in a state of solution in the blood. But, as we shall presently see, the probabilities are that it coagulates as it is produced by the metamorphosis of the blood, that metamorphosis being originally due to the act of respiration. Under an accelerated respiration, the discs oxidize with corresponding rapidity and the amount of fibrine increases; but if the supply of oxygen be limited, there is a restraint on the change of the discs, and the amount of fibrine declines.

The ultimate products of these metamorphoses include of course all the results of the intervening stages, and those ultimate products are chiefly water, ammonia and carbonic acid. We are justified therefore in these physiological discussions in looking at the whole process as one of oxidation, and neglecting intermediate metamorphoses we regard only the final action, and that action is the transmutation of oxygen into carbonic acid, of hydrogen into water, of nitrogen into ammonia.

*Explanation of the General Physical Principle.*—If, in a vessel containing some water, a tube of small diameter be placed, the water immediately rises to a certain point in the tube and remains suspended.


Let the tube be now broken off below that point, and replaced in the cup of water; the liquid rises as before, but though it reaches the broken extremity it does not overflow. A capillary tube may raise water to its highest termination, *but a continuous current cannot take place through it.*

Now, suppose a rapid evaporation of the liquid to ensue from the broken extremity of the tube, as fast as the removal of one portion is accomplished others will rise through the tube, and in the course of time the vessel will be emptied. By evaporation from the upper extremity a continuous current is established; a spirit-lamp, with its cap removed, is an example of this fact.

Or, if the liquid which has risen to the upper end of the tube be of a combustible nature, oil for example, and be there set on fire, as the process of combustion goes on a current will be established in the tube, as in a common oil-lamp in the act of burning.

The principle which I wish to draw from these well-known facts is, that though ordinary capillary attraction cannot determine a continuous flow of a liquid through a tube, there are very many causes which may tend to produce that result.

Let  $a, b, c$  be a capillary tube filled with a certain liquid, between which and the tube there are at different points affinities differing in intensity. Suppose at  $a$  the affinity between the liquid and the tube is intense, that it becomes feebler and feebler towards  $b$ , and at  $c$  has ceased altogether. Under these circumstances there will be a continuous flow through the tube from  $a$  to  $c$ .



To make this quite plain, let us imagine the tube  $a c$  to be formed of combustible matter of any kind, and at the point  $a$  an oxidizing liquid enters it. The liquid, as it passes along the tube, exerts its oxidizing agency, which at the expense of the tube is gradually satisfied. In successive portions of such a tube the affinity is constantly declining. It is greatest at  $a$ , diminishes as it passes along, and ceases altogether at  $c$ . Under these circumstances there will be a constant flow along the tube.

A tube with an included liquid which is thus incessantly varying in its relations will give rise to a continuous movement. At the point of entrance, the liquid, powerfully attracted by the tube, rises with energy; but the chemical changes that set in, satisfying and neutralizing that attraction, to use a common expression, it loses its hold on the tube as it goes, and new quantities, arriving behind, continuously press out those which are before them.

These various results may be expressed in the following general terms.

If a given liquid occupies a capillary tube, or a porous or parenchymatous structure, and has for that tube or structure at different points affinities which are constantly diminishing, movement will ensue in a direction from the point of greater to the point of less affinity.

Or thus:

If a given liquid occupies a capillary tube, or a porous or parenchymatous structure, and whilst in that tube or structure changes happen to it, which tend continually to diminish its attraction for the surface with which it is in contact, movement will ensue in a direction from the changing to the changed fluid.

*Application of this principle to the Circulation of the Blood.*

—Let us now apply these principles to some of the circulations which take place in the human system, and select for that purpose the four leading forms, the systemic, the pulmonary, the portal and the placental circulation.

**THE SYSTEMIC CIRCULATION.**—The arterial blood, which moves along the various aortic branches, contains oxygen

which has been obtained in its passage over the air-cells of the lungs, an oxidation which is indicated by its bright crimson tint. On reaching its final distribution in the tissues, it effects their oxidation, producing heat; and as it loses its oxygen, and receives the metamorphosed products of the tissues, it takes on the blue colour characteristic of venous blood.

If now we contrast the relations of arterial and venous blood to the tissues, it is obvious that the former, from the fact that it can oxidize them, must have an intense affinity for them; but the latter, as it is the result of that action after all affinities have been satisfied, must have an attraction which is correspondingly less.

Arterial blood has therefore a high affinity for the tissues; venous blood little or none. But the change from arterial to venous blood takes place in the manner I have just indicated; and therefore, upon the first of the foregoing general rules, motion will take place, and in a direction from the arterial to the venous side.

By the deoxidizing action of the tissues upon the blood, that liquid ought upon these principles to move from the arteries into the veins, in the systemic circulation. The systemic circulation is therefore due to the deoxidation of arterial blood.

**THE PULMONARY CIRCULATION.**—In this circulation venous blood presents itself on the sides of the air-cells of the lungs, not to carbonaceous or hydrogenous atoms, but to oxygen gas, which being the more absorbable of the constituents of the air, is taken up and held in solution by the moist walls of those cells. Absorption of that oxygen takes place, and arterialization is the result. The blood from being blue turns crimson.

What now are the relations between venous and arterial blood and oxygen gas? For that gas venous blood has a high affinity, as is shown by its active absorption; but this affinity is satisfied and has ceased in the case of arterial blood.

The change from venous to arterial blood, which takes place on the air-cells which are charged with oxygen gas, ought upon these general principles to be accompanied by movement in a direction from the venous to the arterial side.

The pulmonary circulation is due to the oxidation of venous blood, and ought to be in a direction from the venous to the arterial side. These considerations therefore explain the cause of the flow in opposite directions in the systemic and the pulmonic circulation; in the former the direction is from the arterial to the venous side, in the latter from the venous to the arterial. It arises from the opposite chemical reactions

which are taking effect in the system and in the lungs; in the former, as respects the blood, it is a de-oxidation, in the latter an oxidation.

**THE PORTAL CIRCULATION.**—Two systems of forces conspire to drive the portal blood out of the liver into the ascending cava.

1st. The blood which is coming along the capillary portal veins, and that which is receding by the hepatic veins, compared together as to their affinities for the structure of the liver, have obviously this relation—the portal blood is acted upon by the liver, and there are separated from it the constituents of the bile; the affinities which have been at work in producing this result have all been satisfied, and the residual blood over which the liver can exert no action constitutes that which passes into the hepatic veins. Between the portal blood and the structure of the liver there is an energetic affinity, betrayed by the circumstance that a chemical decomposition takes place, and bile is separated; and that change completed, the residue, which is no longer acted upon, forms the venous blood of the hepatic veins. In the same manner, therefore, that in the systemic circulation arterial blood in its passage along the capillaries becomes deoxidized, in consequence of an affinity between its elements and those of the structures with which it is brought in contact, and drives the inert venous blood before it, so too, in the portal circulation, in consequence of the chemical affinities and reactions which obtain between the portal blood and the substance of the liver, affinities and reactions which are expressed by the separation of the bile, that blood drives before it the inert blood of the hepatic veins.

2nd. The blood of the hepatic artery, after serving for the economic purposes of the liver, is thrown into the portal plexus. Hence arises a second force. The pressure of the arterial blood in the hepatic capillaries upon this is sufficient not only to impel it into the capillaries of the portal veins, but also to give it a pressure in a direction towards the hepatic veins; for any pressure which arises between the arterial blood of the hepatic, and its corresponding venous blood, must give rise to motion towards the hepatic veins, no regurgitation can take place backward through the portal vein upon the blood arriving from the chylipoietic viscera, because along that channel there is a pressure in the opposite direction, arising from the arterial blood of the aortic branches. The pressure therefore arising from the relations of the hepatic arterial blood conspires with that arising from the portal blood, and both together join in giving rise to motion towards the ascending cava.

**THE PLACENTAL CIRCULATION.**—The umbilical arteries carry in their spiral courses, as they twist round the umbilical vein, the effete blood of the fœtus, and distribute it by their ramifications to the placenta. In that organ it is brought in relation with the arterial blood of the mother, which oxidizes it, becoming by that act deoxidized itself. The fœtal blood now returns along the ramifications of the umbilical vein, and finally is discharged from the placenta by that single trunk.

That this is truly a change similar to that which is accomplished in the adult lungs, is shown by the circumstance that the blood of the umbilical arteries becomes brighter on its passage into the umbilical vein.

As the venous blood of the fœtus is thus oxidized by the arterial blood of the mother, movement must of necessity ensue in it, on the same principle that it ensues in the adult lung, and must take place in the same direction, that is to say, from the venous to the arterial side.

The fœtal circulation offers a very close resemblance to the circulation of fishes, and is merely a refined variety of that type. The true difference is that in fœtal life the condition of immobility is observed. In fishes the venous blood is brought to the gills, and subjected in their fibrillary tufts to the oxidizing agency of the air dissolved in the surrounding water. In these organs it therefore becomes arterialized, and is pushed into the pulmonary veins. These empty directly into the aorta, no systemic heart intervening, and the mechanical impulse received by the blood during its oxidation is found sufficient to carry on the aortic circulation: the heart therefore may be and is dispensed with. A fish, by spontaneously changing its position, or by the mechanical establishment of currents in the surrounding medium, can obtain new surfaces of water for the oxidation of its blood; but for the motionless fœtal mammalian a higher mechanism is required, a mechanism which can bring the oxidizing-maternal-arterial blood in relation with the branchial or placental vessels. It is true an intricate apparatus consisting of five different classes of vessels is the result, but the play of that apparatus is precisely the same as in the simpler contrivance of fishes.

*Of the Mechanical Force with which these Motions are accomplished.*—The force by which these motions are established is not alone in the proper direction, but also of sufficient intensity. Some years ago I made experiments with a view of establishing this point. Some of them are inserted in the *Phil. Mag.* for Oct. 1838. I found that water, under such circumstances as are here considered, would pass through a piece of peritoneum, though resisted by a pressure of nearly

two atmospheres; and the same facts were observed even in the case of gases. Thus sulphurous acid gas would pass through a piece of India rubber against a pressure of seven and one-third atmospheres; carbonic acid against a pressure of ten atmospheres; and sulphuretted hydrogen, though resisted by more than twenty-four atmospheres.

*Explanatory Remarks on the Coagulation of the Blood.*—When blood recently drawn is kept in a vessel for a space of time it spontaneously separates into two well-defined portions, the one liquid and the other a soft solid—the serum and the clot.

Physicians generally regard this as due to the death of the blood. Whilst it is in the system it is under the influence of the vital force; but when removed it spontaneously undergoes the change in question, and, unable to keep its primitive condition, coagulates and dies. Accordingly this partial solidification of the blood is looked upon as a mysterious phænomenon, and though from time to time many experiments have been made and explanations offered, that which refers it to the presence or absence of the vital principle appears to be most generally received.

But it is very doubtful whether any such special power as a vital force exists. In the instance under consideration I cannot comprehend how a loss of vitality in the blood can in any manner elucidate or indeed have anything to do with the fact of its coagulation.

It appears to me that what occurs to the blood when drawn is precisely the same as that which occurs to it continually when in the system. If its fibrine coagulates in the receiving cup, it tends equally so to do in the peripheral circulation. I can see no difference in the two cases. And if this be true, it obviously is a fruitless affair to be seeking for an explanation of a difference in habitudes in and out of the system, when those differences in reality have no existence in nature.

If, when blood flows into a cup, we could by any mechanism withdraw the particles of fibrine as they agglutinate together, the phænomenon of coagulation would never be witnessed; and this is precisely the result in the living mechanism. The fibrine, as it passes into the proper condition, is removed by a series of events which will be hereafter explained. But whether it be in those states which physiologists designate living or dead, it exhibits continually the same tendency.

When we remember that the average amount of fibrine in blood scarcely exceeds one-five-hundredth part of its weight, and that this minute quantity is sufficient, by entangling the blood-discs, to furnish so voluminous a clot, we have little

difficulty in understanding the cause of the false importance which has been attached to the fact of its coagulation. When we also remember that the phænomenon is one which, far from taking effect instantaneously, requires a considerable length of time, and estimate duly the demand that is made for fibrine by the system upon the blood, we shall have no difficulty in perceiving the truth of the observation which I thus wish to bring into a clear point of view,—that the tendency to coagulation in the system is as great as it is out of it, and that the true difference in the two cases is, that in the former the resulting solid is taken up and appropriated to the wants of the œconomy; in the latter it remains undisposed of, and, entangling the blood-discs in its meshes, produces a voluminous and therefore deceptive clot.

It is with this matter of the coagulation of blood precisely as it was formerly with putrefaction. Many of the older physiologists defined a living body as a mechanism having the quality of resisting external changes. After death its parts were ultimately resolved into water, ammonia, and carbonic acid. But better views on these topics are now entertained, and we know that the living body undergoes these putrefactive changes just as much as the dead, but then in it there are appointed routes by which the resulting bodies may escape; the carbonic acid through the lungs, the nitrogenized compounds through the kidneys, the water through both these organs and the skin. It is in this as in the coagulation of the blood, there is no difference in the chemical changes taking place, the difference consists in the disposal finally made of the resulting products.

That coagulation tends to take place equally in the living system as out of it, there is abundant proof. What are all the muscular tissues which constitute by far the larger portion of the soft parts, but fibrine which has thus been separated from the blood? And those muscular tissues every moment are wasting away, and giving origin to the metamorphosed products that we find escaping from the lungs, the kidneys, the liver; from what source then do they repair their waste, if not from fibrine coagulated from the blood during the act of life? Every muscular fibre is a living witness against the doctrine that it is death that brings on the coagulation of the blood.

That the truth of this view, which at first sight may appear indefensible, may be more clearly made out, let us consider under what circumstances the blood is placed whilst moving in the system. We have to remember that coagulation is not an instantaneous phænomenon, but one which requires a con-



siderable lapse of time. And now, assuming the doctrine which I am advancing to be true, there are very obvious reasons that the blood, so long as it moves in the system, has its tendency to coagulate satisfied in a very partial manner. Let us observe its course. It leaves the left ventricle of the heart, one pulse-wave succeeding another with rapidity, and is distributed through all the aortic branches. It takes but a few seconds for this movement to be complete, a period far too short to allow coagulation to take place; it now passes on through the capillaries, or moves through parenchymatous structures; and here, even though a great delay may occur, inasmuch as the passages are so sinuous and often so minute that the discs can move but in a single file at a time, how is it likely, under such circumstances, that coagulation should ensue? For that to take place, it is needful that there should be a free communication throughout the mass, that each particle of fibrine brought into relation with those around it may exert its plastic power and join itself to them. But in the peripheral circulation it is isolated, the cells over which it is moving, or the narrow tubes through which it goes, protect it from other particles around, and on escaping into the commencement of the venous trunks, it is hurried in the torrent of the circulation at once to the heart. Without delay the right auricle and ventricle pass it forward to the lungs, and if any tendency to set had been exhibited during the brief moment of its passage, it is again distributed upon the capillaries of the lungs, and is situated precisely as it was when in the capillaries of the peripheral system.

In this manner I regard the coagulation of blood as a simple mechanical result, having no connexion with life or death, or the fictitious principle of vitality. At the two extremes of the circulation, the peripheral and the pulmonary, there is a sorting process continually going on. If a man were to agitate a quantity of this liquid in a tube, having a contrivance at each extremity to keep the particles of fibrine as they passed apart from one another, their plastic tendency to cohere could never be satisfied, and coagulation could never ensue. And this condition of things is, to a certain extent, approximated to in the mechanism of the body.

It thus appears that by the intervention of two capillary circulations, one in the lungs and the other in the system, the coagulation of blood must be greatly retarded, though the tendency to produce that result is quite as great as when the fluid is removed from the system. And with such an obvious explanation before us, why should we resort to any occult agency, or envelope the phenomenon in mystery, when it is plainly a mechanical affair?

Physiologists have never given a full value to the facts, that the setting of the blood requires time and a free communication through all parts of the fluid mass. If it be subjected incessantly to a mechanism which divides it into portions of inconceivable tenuity, and every moment isolates each particle from all its fellows, its coagulation must be greatly restrained. It is upon the same principle that the expressed juices of carrots and turnips deposit a fibrinary clot, as M. Liebig and others have observed. Whilst they are enveloped in the cells of those vegetables coagulation cannot take place, for each granule of fibrine is shut out from the others. What need is there to resort to a vital principle to explain for the human œconomy a result which equally obtains in the case of those humble plants, or why with some physiologists impute to the nervous system the quality of maintaining fluidity in the blood? These vegetables have no nerves.

The application of the principles here set forth furnishes a very felicitous explanation of a great number of effects which we witness, to some of which I may briefly refer. It is well known that after ordinary death, whilst the arteries are empty, the systemic veins and also the right cavities of the heart are full of venous blood. The reason is clear, although the ordinary theory, that the heart acts like a pumping machine, fails, as is well known, to explain it. As long as arterial blood is deoxidizing it will move to the venous side, a movement which must continue until the arteries are empty.

But it may be asked, why do not the right auricle and ventricle relieve the veins, and by their hydraulic action in the last moments of life push the accumulating blood through the pulmonary system? Again the reason is clear. *Movement through the lungs cannot take place except when oxidation is going on.* The systemic capillaries continuing their action long after the last breath is drawn, they make the blood accumulate in the veins, and from them there is no escape.

In the same way, in fainting, the blood leaving the arteries accumulates on the venous side, and as its flow is dependent on the push of the arterial blood entering the capillaries, so soon as no more enters no pressure is exerted on the venous trunks, and if a vein is opened there is no discharge, and under such circumstances hemorrhages at once stop.

After ordinary death, although the systemic arteries are empty, the pulmonary artery is full. That this should be the case is indicated upon our principles, for the blood cannot pass from the terminal ramifications of the pulmonary artery into the veins except by being oxidized. Respiration having ceased oxidation cannot take place, the movement is checked, and the blood remains in the artery.

In a paroxysm of asthma the lungs become obstructed with mucous secretions, and the rapidity of oxidation is therefore interfered with. Under such circumstances the passage of the blood is retarded, as is shown by the great dilatation of the jugular veins.

Whatever therefore deranges the process of oxidation deranges the flow of the blood. In violent expirations, such as in coughing, the observations of Haller show that the blood moves tardily in the lungs, and in delicate persons its retardation is so complete that it regurgitates in the great veins.

In a violent and continuous explosion of laughter, the jugular veins become excessively distended; the right cavities of the heart having no power to push the venous blood through the pulmonary capillaries, and owing to the expulsion of air from the air-cells, the blood itself fails to effect the passage with its usual speed. In this instance it must again accumulate in the veins.

The various cases here cited depend on retarded oxidation. I might now consider the reverse of this, or where oxidation goes on too rapidly, as when protoxide of nitrogen is breathed. Owing to the great solubility of this gas in serum, and its power of supporting combustion, we should expect to find it exert that control over the circulation which is well known to be one of its peculiarities. This paper is however extended to so great a length, that here I must stop, though I have made no allusion to the movements in the lymphatics or lacteals, or to the flow of sap in trees, or to the circulatory movements of the lower animals. These can all be explained upon the same principle; thus the descent of the sap follows as a necessary consequence of the decomposition of carbonic acid in the leaf. Nor have I said anything of the obvious control which certain classes of nerves have over the systemic oxidation. There are many facts which prove that the nervous system regulates this operation, and can either facilitate it or keep it in check. In this there is nothing extraordinary. A piece of amalgamated zinc exhibits no tendency to oxidize in acidulated water, but by the touch of silver or platina it is made to submit itself to the action of that medium. The act of blushing, and all local inflammations, show that changes in the relations of the nervous system control the oxidizing action of arterial blood; but to these things I propose to return on a future occasion. What is here stated is sufficient to illustrate the general principle to which I wish to draw attention, that *the chemical changes which are impressed on these circulating fluids are the true causes of their flow.*

XXXIV. *On the Existence of Finite Algebraic Solutions of the general Equations of the Fifth, Sixth, and Higher Degrees\**.  
By JAMES COCKLE, M.A., Cantab.; Special Pleader†.

7. WHEN  $y = \Lambda' x^{\lambda'} + \Lambda'' x^{\lambda''} + \dots + \Lambda^{xi} x^{\lambda^{xi}} \dots$  (n.)  
and  ${}_3Y_n = h_1^3 + h_2^3, \dots$  (o.)

$h_1$  and  $h_2$  having the forms of the quantities squared in (b.) ‡, what is the limit of  $n$ ?

8. Make  ${}_3Y_n = h_1^3 + j' \Lambda' + \mathfrak{H}'' \dots$  (p.)

then §  $j' = J_2^{(2)} + J_4^{(2)} + \&c.; \dots$  (q.)

but  $J_2^{(2)} = 0$  and  $\Lambda''$ ,  $\Lambda'''$ , disappear from  $j'$ , if ||

$y = \Lambda' x^{\lambda'} + L'' + L^{iv} + L^v + \dots + L^{xi} \dots$  (r.)

and  $L^m = \Lambda^m \left( x^{\lambda^m} - \frac{\gamma_m}{\gamma_3} x^{\lambda''' } \right) \dots$  (s.)

So,  $\mathfrak{H}'' = h_2^3 + j'' \Lambda'' + i^{iv} \dots$  (t.)

9. Let  $y_r = L' x_r^{\lambda'} + L'' \left( x_r^{\lambda''} - \frac{\gamma_2}{\gamma_3} x_r^{\lambda''' } \right) + l_r, \dots$  (u.)

$l_n = 0$ ,  $L = \Lambda + l$ , and  $l$  a constant, then ¶,

$$\left. \begin{aligned} 0 = j' &= [l_1 \cdot l_{n-1}]_1^2 \\ 0 = j'' &= [l_1 \cdot l_{n-1}]_2^2 \\ 0 = i^{iv} &= [l_1 \cdot l_{n-1}]_3^3 \end{aligned} \right\}, \dots \dots \dots (\alpha.)$$

$\therefore n - 1 > 3$ , or  $n > 4. \dots \dots \dots$  (v.)

10. Again,  $j' = 0$  is equivalent to\*\*

$$\left. \begin{aligned} \gamma_2^{iv} \Lambda^{iv} + \dots + \gamma_2^{xi} \Lambda^{xi} &= 0 \\ \gamma_3^{vi} \Lambda^{vi} + \dots + \gamma_3^{xi} \Lambda^{xi} &= 0 \\ \gamma_4^{viii} \Lambda^{viii} + \dots + \gamma_4^{xi} \Lambda^{xi} &= 0 \\ \gamma_5^x \Lambda^x + \gamma_5^{xi} \Lambda^{xi} &= 0 \end{aligned} \right\}, \dots \dots \dots (\beta.)$$

$j'' = 0$ , on eliminating  $\Lambda^{2r+1}$ , to ††

$$\left. \begin{aligned} {}^{iv}\gamma_1 \Lambda^{iv} + \dots + {}^x\gamma_1 \Lambda^x &= 0 \\ {}^{viii}\gamma_2 \Lambda^{viii} + {}^x\gamma_2 \Lambda^x &= 0 \end{aligned} \right\}, \dots \dots \dots (\gamma.)$$

$i^{iv} = 0$ , on eliminating  $\Lambda^{vi}$ ,  $\Lambda^x$ , to †††

${}^{iv}\gamma_1 \Lambda^{iv} + {}^{viii}\gamma \Lambda^{viii} = 0. \dots \dots \dots$  (δ.)

\* See my presumed solution of the equation of the fifth degree, at page 125 of the last volume of this Magazine. I there used the ratios  $z_1, z_2 \dots$  of the quantities  $\Lambda', \Lambda'', \dots$  to one of their number, but have here employed other ratios, or, more properly speaking, the quantities themselves.—J. C.

† Communicated by the Author. ‡ Phil. Mag., this vol., p. 132.

§ Ibid. S. 3. vol. xxvii. p. 126. || Ibid. p. 293 (16.).

¶ Ibid. p. 126, note ‡, and this vol., p. 132, par. 2.

\*\* Phil. Mag. S. 3. p. 126, line 9. †† Ibid. (f.) and (g.)

†† Ibid. (h.)

11. Let\*  $L^{iv} + L^{vi} + \dots = p'_0 + p'_1 x + \dots + p'_{n-1} x^{n-1}$ , (w.)  
 then †,  ${}^{iv}\Sigma x^i . L = (1+p)(p'_0 + \dots) + \dots + q_{n-2} x^{n-2}$ ; . (x.)  
 and we have, in general,  $n$  quantities  $1 + p, q_0, \dots, q^{n-2}$ , to satisfy 3 homogeneous equations  $0 = j' = j'' = {}^{iv}$ , or,  $n$  quantities  $p'_0, p'_1, \dots, p'_{n-1}$  to satisfy 3 other homogeneous conditions ( $\gamma$ .) and ( $\delta$ .), using the former  $n$  quantities to satisfy the group ( $\beta$ .), non-homogeneous with respect to them; hence

$$n > 3. \quad \dots \dots \dots (y.)$$

12. (v.) and (y.) are not inconsistent, for, if

$$A_0 = 1, \frac{\Lambda_m}{\Lambda_{m-1}} = \frac{n-r+m}{nm} \dots \dots \dots (z.)$$

and

$${}_r Y'_n = p_r - A_1 p_{r-1} p_1 + A_2 p_{r-2} p_1^2 + \dots \pm \frac{n(r-1)}{r} A_{r-1} p_r^r, (aa.)$$

then this accented value of  $Y$  is a critical value from which  $p'_0, q_0, \dots$  disappear, and, since (2.) and (3.) ‡ are respectively

$${}_3 Y'_5 = 0, \text{ and } {}_4 Y'_5 - \frac{1}{5} {}_2 Y'_5{}^{12} = 0, \dots \dots (ab.)$$

my solution would fail if  $n-1$  were  $< 4$ . Hence, for all critical functions (y.) degenerates into (v.); and, after solving critical equations, we shall have quantities  $p'_0, q_0, \dots$  left for satisfying (other) conditions whose degrees are unaffected by our previous operations.

13. However numerous might be the groups ( $\beta$ .), ( $\gamma$ .), ( $\delta$ .), or the relations forming those groups, it would seem that some of the  $\Lambda$ 's being lost at each descending step, the limit will not be proportionally elevated. We may make

$$\Sigma_r = (1+p) {}^{(r)}\Sigma_{r-1} + \dots + q_n^{(r)} x^{n-2}, \dots \dots (ac.)$$

for the  $\Lambda$ 's introduced as we ascend the groups.

14.  $\Lambda'$  and  $\Lambda''$  will introduce new values of  $1+p$  and  $q$  into the (now to be combined) equations (2.) and (3.), or (ab.), but, as it appears to me, no new difficulty. The  $\gamma$ 's may be derived from the  $\tau$ 's §, and from each other by one operation  $\theta$ , and if

$$\gamma = \Theta(\tau), \text{ then } \gamma' = \Theta(\gamma), \text{ \&c. } \dots \dots (ad.)$$

Grecian Chambers, Devereux Court,  
 January 31, 1846.

JAMES COCKLE, Jun.

\* See Sir W. R. Hamilton's "Inquiry, &c." into Mr. Jerrard's method (Sixth Report of the British Association) from [4.] p. 301 to line 9 of p. 304.

† Ibid. p. 303.

‡ Phil. Mag. S. 3. vol. xxvii. p. 125.

§ Ibid. pp. 292, 293.

XXXV. *On some New Species of Animal Concretions.*

By THOMAS TAYLOR, Surgeon.

[Continued from p. 46.]

*Resino-bezoardic Acid Calculi.*

VERY shortly after commencing the examination of the calculi in the College collection, my attention was drawn to several concretions which possessed the easy fusibility and general characters of a resin, and which were described in the MS. Catalogue as "false West Indian Bezoars," on the supposition that they were artificial compounds. The peculiar characters however of the resin of which they consisted, and their finely laminated structure, which it would be impossible to imitate, left no doubt on my mind of their being genuine bezoars, and in January 1841 I described them to the Museum Committee as consisting of a *vegetable* resin, derived most probably from the resinous juices of the plants on which the wild goats of the East had fed. In the same year a very interesting paper appeared in the *Annalen der Chemie und Pharmacie*, by M. Goebel, describing a new species of calculus which he had found in the Zoological Museum at Dorpat, and to which, on the supposition of its being a biliary concretion, he gave the name of *lithofellinic acid*. A similar calculus from the Pathological Museum at Göttingen was shortly after examined by Professor Wöhler.

The similarity in chemical characters of the concretions examined by these chemists with the resinous concretions previously examined by myself, rendered it certain that they were identical in composition; but as it was important to determine whether they were biliary calculi or simply intestinal concretions, derived from the materials of the food, I repeated at some length my experiments, but without coming to any other conclusion than that formerly expressed. The reasons which have induced me therefore to place the calculi among the intestinal calculi in the College Catalogue are as follows.

In the first place, the greater number of them contain, as the subjoined analysis will show, a small quantity of a soft viscid resin, resembling a vegetable balsam.

Secondly. They resemble all other concretions formed in the intestines, by having a foreign body, as a piece of wood or a seed, for their nucleus.

Thirdly. They frequently attain a very large size, quite inconsistent with the notion of their being biliary concretions, or having been contained in the gall-bladder. There is one specimen in the Museum which measures three inches and a half in length, and the same in its greatest breadth. This

calculus is of a rude triangular figure; it has evidently been accompanied by other calculi, as both of its extremities possess the smooth depressed surfaces found in concretions which have been in contact with others. Another specimen, of an oval figure, is four inches in length by three in breadth. Against the notion that these concretions may have been formed from the natural or the altered constituents of the bile concreting around foreign bodies in the intestines, it may be remarked that we have no other instance of a biliary calculus being so formed\*. The large biliary concretions which are sometimes passed *per anum* by the human subject, have undoubtedly received no increase in bulk while in the intestine, but have made their way into the intestine either through an ulcerated opening or through the dilated biliary duct, which is capable of undergoing dilatation to a much greater extent than is generally imagined.

The circumstance of the Oriental Bezoar being composed of a vegetable acid, as I have shown in a former paper, together with the assertion of most Oriental travellers, that the resinous concretions are found in the stomach of the animal (not a very likely spot for a biliary calculus), adds considerable weight in favour of their vegetable origin. It is however right to state that I have not been able to detect the presence of resino-bezoardic acid in several of the known resins. Our acquaintance with these substances is however so limited that it would require a very extended series of experiments to determine this question in the negative. In its chemical relations, resino-bezoardic acid closely resembles the pimelic acid of M. Laurent, which that excellent chemist has recently shown to be the natural acid of the fir. This fact, coupled with the circumstance that the calculi are not very uncommon, and that vast forests of pines abound in the regions inhabited by the goats, render it not improbable that this resin is derived from some of the fir tribe. As the term *lithofellinic acid* gives therefore an erroneous idea of the origin of these concretions, I have ventured to substitute that of *resino-bezoardic acid*, which does not differ materially from that of "résine animale bézoardique" given to them by Fourcroy. This name will also serve to identify the circumstances under which it was first discovered, should its natural source be hereafter ascertained.

\* Ambergis perhaps forms an exception to this statement. This substance is found in the intestines of the Spermaceti Whale, or floating on the sea. In the Catalogue I have placed it among intestinal concretions, although I have pointed out at the same time that it is a biliary product; its principal constituent, ambreine, bearing the same relation to the bile of the Whale as cholesteroline does to that of Man.

Resino-bezoardic acid calculi are usually of an oval figure. Their external surface is smooth and polished, and has generally a greenish yellow, green, or a light brown colour. They are made up of thin concentric layers, which are frequently of a deeper tint than the exterior. In the centre of the calculus some foreign body is invariably found which forms the nucleus. These calculi are exceedingly brittle; the fracture is conchoidal, and has a resinous lustre. They vary considerably in size, but are usually larger than the ellagic acid species. One specimen in the Museum measures nearly ten inches in circumference. They melt like resin in the flame of a candle, and when more highly heated, give off white vapours, which have an aromatic odour, catch fire, burn with a brilliant flame, and leave behind a small shining carbonaceous ash.

Resino-bezoardic acid calculi readily dissolve in alcohol, with the exception of a small quantity of flocculent matter. The alcoholic solution varies in colour in different calculi, but is usually of a red or greenish-red tint. The solution gradually deposits small crystals, which, when examined by the microscope, are seen to consist of low six-sided prisms with flattened extremities. When the alcoholic solution is mixed with water the resin is thrown down. The precipitate appears under the microscope in the form of small crystalline tufts.

Digested in solution of potass these calculi readily dissolve, the solution is of a brownish green colour, and when neutralized by an acid, a thick curdy precipitate is produced, which by agitation adheres together, and while warm may be kneaded between the fingers or drawn into threads like cobbler's-wax. The viscosity of this precipitate is owing to another resinous matter which the calculi contain; for the pure resino-bezoardic acid similarly treated forms an amorphous precipitate which cannot be made to adhere together. They dissolve in solutions of ammonia and its carbonate. In concentrated sulphuric acid they also dissolve. The solution is of a red colour, and is rendered turbid by the addition of water. The precipitate is not crystalline, like that from its solution in alcohol, but consists of minute transparent yellow particles. Nitric acid acts with energy upon these calculi, nitrous acid is evolved, and a light red solution is formed, which quickly becomes yellow.

#### *Analysis.*

About 400 grains were reduced to a fine powder, mixed with distilled water, and subjected to distillation in a glass retort until about two ounces had passed over. The distilled



liquid was quite transparent, and possessed the peculiar aromatic odour of the calculus, but no volatile oil could be detected. The powder was separated from the rest of the water by filtration, and dried at 200° Fahr. It was dissolved in twelve ounces of boiling alcohol. The solution was of a bright red colour when viewed by transmitted light, and had a greenish tinge by reflected light; with the exception of a small quantity of flocculent matter it was quite transparent.

In order to separate the insoluble matter, the liquid while still hot was filtered, and the matter on the filter washed with a fresh portion of alcohol and dried. This matter was of a dirty brown colour, with a shade of green. It was when quite dry rather soft, so as to admit of being moulded between the fingers. When heated on platina foil it did not fuse, but softened, caught fire, and burnt briskly, emitting at the same time the odour of heated Indian rubber. It was insoluble in water, either hot or cold.

That this substance was not caoutchouc, was shown by its not being dissolved or softened when acted upon by absolute æther or oil of turpentine. A solution of caustic potass extracted some of its colour, but did not appear to dissolve it. The exact nature of this matter I am unable to decide; its vegetable nature is rendered probable by the total want of any animal odour while burning. It amounted to about two per cent.

The filtered alcoholic solution became slightly turbid on cooling; after standing a short time small crystals were deposited, and a crystalline crust formed upon its surface. Some of the crystals when examined by the microscope had the form of very regular six-sided plates, and others that of six-sided prisms. When a drop of the liquid was allowed to evaporate on a glass plate, and the residue examined by the microscope, crystals were formed, whose figure was not very distinct, but appeared to be that of a six-sided prism lying on its side; occasionally a six-sided plate was also visible.

The liquid was put into a retort, and about two-thirds of its bulk distilled over. It was transparent while hot, but on cooling deposited an abundant crop of small crystals. These crystals had the form of three-sided plates; when carefully fused upon a slip of glass, they were converted into six-sided plates.

The crystals obtained at different times were purified by being repeatedly crystallized from their alcoholic solution, which removed nearly the whole of their colouring matter. They possessed all the characters of the lithofellinic acid of Professor Goebel, and constituted the bulk of the calculus.

The mother-liquor was mixed with water, when a precipitate separated which by agitation was converted into a viscid turpentine-looking substance that adhered to the sides of the glass. When a drop of the mother-liquor was evaporated on a glass plate, very few crystals could be detected, but a great number of thick, viscid oily-like drops: by heating the glass vapours arose, and a hard uncrystalline resin was left. When the alcoholic solution of the crystals was mingled with water, a crystalline precipitate was thrown down, which beneath the microscope appeared in the form of small irregularly-shaped prisms, arranged in stellate groups. This difference in the character of the two precipitates appeared to indicate that the mother-liquor contained either a volatile oil or some soft resin in addition to the crystalline resin previously described. To determine this question the whole was placed in a retort, and submitted to distillation; the spirit came over quite free from any essential oil, merely retaining the peculiar odour of the calculus: the last portions smelt much stronger, and were slightly turbid. The precipitate had melted and formed a deep red oil, which adhered to the sides of the retort; when cold it was soft and ductile between the fingers. It was readily soluble in solutions of potass and ammonia, the solutions were rendered milky by the addition of an acid, but no precipitate fell. The milky liquor when examined by the microscope gave the appearance of oily globules. The soft resin remaining in the retort was now divided into two portions; to the one solution of ammonia was added, and to the other æther.

The ammoniacal solution was perfectly clear and of a bright red tint; it was neutralized by muriatic acid, a viscid precipitate separated, which was collected together, washed, broken into fragments, and put into a glass tube together with æther. It only partially dissolved, and after standing some days six-sided prisms were found adhering to the tube, the æthereal solution was evaporated, and a resinous matter more fusible than the former was left.

That portion of the soft resin which had been digested with cold æther did not entirely dissolve, but left some crystals of resinous matter undissolved; the æthereal solution was evaporated, and the residue, which was quite similar to that which had been previously treated with ammonia, was mixed with it and both dissolved in alcohol, sp. gr. 0.840. The tincture was set aside for some weeks; only a small quantity of crystalline matter was deposited, together with a little soft resin; it was therefore distilled, and the residue again treated with absolute æther, in which, with the excep-

tion of a very small quantity of resin, it entirely dissolved. On distilling off the æther a semi-fluid viscid balsam of a dark-red colour was left, which did not solidify at the temperature of the air, and acquired a pellicle on its surface by exposure to air. When heated a portion of it was volatilized, giving off at the same time the odour of melted caoutchouc. It readily caught fire and burnt brightly; its combustion was unaccompanied by the slightest trace of the odour given out by animal matter. It readily dissolved in caustic potass, and the addition of an acid threw it down unchanged: it possessed a biting acrid taste, felt particularly about the fauces; by exposure to the air it became a hard resin.

When submitted to distillation in a small tube retort, no oil passed over until the resin had acquired a temperature at which it began to decompose, when an empyreumatic oil came over: the quantity submitted to distillation was, however, too small to render the experiment quite satisfactory.

The only conclusions that can be drawn from this analysis are, that the principal constituent of the calculus is a vegetable resin, which is characterized by crystallizing in the form of six-sided prisms; that it is accompanied by a small quantity of a soft resin, probably containing volatile oil; that in addition to these it contains some other substances, as colouring and extractive matter, the precise nature of which it is impossible to determine, but which are doubtless also of vegetable origin.

M. Goebel detected in the concretion examined by him a small quantity of the colouring matter of the bile. In no one of the concretions examined was I able to satisfy myself of the presence of that substance. It is probably therefore only an accidental constituent. Its presence is however no proof of their biliary origin, since the colouring matter and other constituents of the bile are frequently found in hair-balls and other concretions known to be formed in the intestine.

Resino-bezoardic acid, when freed from the other substances with which it is mixed in the calculus, possesses the following properties:—It slowly dissolves in cold alcohol, more rapidly in hot; according to Goebel, one part of resino-bezoardic acid requires 29·4 of alcohol to dissolve it at 68° of Fahrenheit and 6·5 of boiling alcohol; in cold æther it is very sparingly soluble, 444 parts being required, but only 47 when boiling. Its alcoholic solution has an acid reaction, and the resin is slowly deposited from it in the form of short six-sided prisms. The crystals are exceedingly small; they have generally a yellow tint, but may be obtained quite colourless by previously digesting the alcoholic solution with animal char-

coal : they are hard, brittle, and easily reduced to powder, inodorous, and have a bitter resinous taste ; their summits are generally quite flat, but are sometimes bevelled at their edges. Three-sided prisms are occasionally deposited, which apparently result from an extension of each alternate face of the six-sided prism.

The crystallized acid fuses at  $401^{\circ}$  Fahrenheit, and when not heated beyond that temperature becomes on cooling an opaque crystalline mass. If the fused acid be heated only a few degrees above  $401^{\circ}$  Fahrenheit, it forms when cold a transparent glass, without the slightest trace of crystalline structure : when alcohol is poured over the fused mass a number of minute cracks are suddenly formed, which possess considerable regularity. If a thin layer of alcohol is allowed to remain over it, the whole is quickly converted into an aggregated mass of regular crystals. The most remarkable circumstance is that the melting-point of the vitreous or amorphous resino-bezoardic acid is nearly  $180^{\circ}$  lower than that of the crystallized acid, Prof. Wöhler having determined that the crystallized acid melts at  $400^{\circ}$  Fahrenheit, while the amorphous fuses at a temperature between  $220^{\circ}$  and  $230^{\circ}$  Fahrenheit. In this respect resino-bezoardic acid resembles sugar, sulphur, amygdaline and silvic acid ; all of which bodies have two distinct fusing-points, according as they are either in a crystalline or amorphous state : this property Wöhler believes to be possessed by all dimorphous bodies. The above fact may be readily observed in the following manner, which serves as a very characteristic test of resino-bezoardic acid. Let a few grains of the powdered resin be strewed over a thin slip of glass and held over the flame of a candle until a portion only of the resin is melted : if the edges of the semi-fused portion be examined by the microscope groups of very regular six-sided plates are seen ; the perfectly fused portion is glassy and devoid of crystalline structure. This test does not always succeed with the raw calculus, owing to the foreign substances which it contains. When heated beyond its melting-point this acid gives off white vapours, which have an aromatic odour ; it finally catches fire and burns like resinous bodies in general.

Resino-bezoardic acid is insoluble in water and muriatic acid. It is thrown down from its alcoholic solution by water as a white precipitate, which under the microscope appears in the form of small prismatic crystals arranged in stellate groups. It is readily soluble in a solution of potass, soda, ammonia, and carbonate of ammonia, and is precipitated on the addition of an acid. The precipitate at first forms a dense white

coagulum, but shortly becomes pulverulent; when examined by the microscope it is not crystallized, but consists of minute transparent amorphous particles. It melts at 220° Fahrenheit, and is evidently the amorphous state of the acid.

When the potass solution is evaporated a transparent gummy mass is left, which is insoluble in solution of potass, but dissolves in pure water. When the potass solution is concentrated by boiling, the compound of the resin and alkali separates from the liquid and swims on its surface; when cold it forms a hard yellowish mass like resin, which dissolves in æther, alcohol and water. When the ammoniacal solution of this acid is evaporated, the resin separates unaltered. Nitric acid decomposes this acid; nitric oxide gas is evolved, and a beautiful red solution formed, which quickly becomes yellow.

Concentrated sulphuric acid dissolves the resino-bezoardic acid: the resin is precipitated unaltered on the addition of water in the amorphous state.

3·777 grs. of the crystallized acid, which had been rendered perfectly colourless by digestion with animal charcoal, when dried at 180° Fahr. in a current of dry air and burnt with chromate of lead, gave 3·64 water and 9·793 carbonic acid. This result agrees with the analyses of Messrs. Ettling and Will and Professor Wöhler, who found—

	Ettling and Will.		Wöhler.		T. Taylor.
Carbon	71·19	70·80	70·83	71·09	70·71
Hydrogen	10·85	10·78	10·60		10·71
Oxygen	17·96	18·42	18·57		18·58
	100·00	100·00*	100·00†		100·00

Messrs. Ettling and Will, who have analysed some of its salts, regard the formula of the crystallized acid as  $C_{42}H_{74}O_7 + HO$ , while Professor Wöhler represents it as  $C_{40}H_{70}O_7 + HO$ .

Resino-bezoardic concretions were first examined by Fourcroy and Vauquelin. Their account is very slight and imperfect, but is accompanied by a very accurate drawing of a fragment of one of them. Fourcroy states, without mentioning his authority, that they are taken from some unknown species of Asiatic or African animals, and believes them to be formed from the resinous juices of the plants on which these animals fed.

In the College Catalogue I have described them as being the true *Occidental Bezoar*. Subsequent consideration how-

\* *Ann. der Chem. und Pharm.*, xxxix. 242.

† Poggendorff's *Ann. der Phys. und Chem.*, liv. 259.

ever inclines me to believe that the true Occidental Bezoar consisted of diphosphate of lime, and that these concretions, which Kæmpfer states were termed in Persia *Lapis Bezoar Occidentalis*, on account of their similarity to the concretions brought from South America, were so called from their exterior possessing the same smooth polished exterior as the diphosphate of lime concretions. The concretions described by Kæmpfer under the name of *Coagulum Resinosum Bezoarticum*, are evidently identical with resino-bezoardic acid calculi; for he says that the Swedish ambassador, on his departure from Ispahan, purchased some specimens, which, when thrown upon burning coals, melted and gave out an aromatic odour like that of colophony or olibanum. In the work of Clusius there is a figure of the occidental bezoar which is quite characteristic of this calculus, and Monardes asserts that they were taken from the wild goats of Persia. It is not however probable that any particular species of concretion was confined exclusively to the animals of one or the other hemisphere, since the resinous and bitter juices from which the concretions are formed exist in the plants of both divisions of the globe.

[To be continued.]

XXXVI. *On the Winter Storms of the United States.*  
By Lieut.-Colonel SABINE.

*To the Editors of the Philosophical Magazine and Journal.*

GENTLEMEN,

**Y**OUR meteorological readers, and especially those who take an interest in the law of storms, will, I am sure, be glad to have their attention drawn to a second memoir by Professor Loomis of New York, on the phænomena of the great storms which are experienced in the United States during the winter months. In this memoir (Art. IV. of vol. ix. of the Transactions of the American Philosophical Society) two storms are investigated, one of which occurred about the 3rd of February, 1842, and the other about the 15th of the same month of the same year. The method of investigation is the same which Professor Loomis adopted in his account of the great storm of December 20th, 1836, viz. the assemblage in one view of the atmospherical circumstances simultaneously observed over the whole extent of the United States, both during the continuance of the storm and for one or two preceding days. It is by this path that we may confidently hope to attain to a knowledge of the causes which produce these great atmospherical derangements; and, thanks to the spirit

of co-operative labour which distinguishes the present time, we have every prospect of seeing this path successfully pursued. Although the memoir itself is not very long, the illustrations which accompany it are many, and its republication in this country without them would not convey the full amount of instruction to be derived from the text and plates conjointly. Memoirs of this description scarcely admit of an abstract; there are however certain circumstances which present themselves in so striking a manner as common to the three storms above mentioned, as to induce the belief that they may be viewed as the characteristics of a particular class of storms which occur in the United States in the winter months of every year. The circumstances alluded to may admit of a brief notice in the light of a first generalization; and it may have the additional advantage of attracting the attention of some of your readers to the original memoir in the Transactions of the American Philosophical Society.

We may picture to ourselves, in the first instance, a normal state of the atmosphere over the United States, in the departure from which we may trace the successive phases of derangement which constitute the storm. In this normal state the wind is from the west, or a few degrees south of west, in the lower as well as in the upper current, with the thermometer and barometer at or near their respective mean heights for the time and place: the whole body of the air from the surface of the earth to its upper limit, is proceeding harmoniously in the one direction, and having blown across the greater part of the continent of America before it reaches the middle states of the union, it is extremely dry, and the atmosphere perfectly clear.

The interruption to this normal state, which in the order of time appears first to present itself, is *a change in the direction of the lower stratum of the air*, which becomes southerly in the countries situated in the north of the Gulf of Mexico, and south-easterly in the south-eastern states. The change in the direction of the lower stratum of air is speedily followed, or perhaps it should rather be said is accompanied, by cloud, and by a rise of temperature, which progressively increase, the one in extent and the other in intensity, attended by a falling barometer. The cloud condenses into rain or snow, the area of which progressively extends till but a comparatively small margin of cloud remains without precipitation. The thermometer continues to rise, the barometer to fall, and the rain or snow to descend, until the instant when the abnormal winds from the south and east give place to a more violent rush of air from the west and north-west, by which

the phenomena of the storm are swept onwards, and transferred successively from the middle to the eastern states, and thence to the sea, with a velocity which in different instances has been noted to vary from about twenty to thirty-six statute miles an hour. The maximum of the thermometer and minimum of the barometer coincide generally (and with great exactness in the eastern states) with the change of the wind, the derangement of the temperature being so great as  $20^{\circ}$  and even occasionally  $30^{\circ}$  above its normal state.

The description thus given was written after reading the second memoir, and was consequently drawn principally from the phenomena of the two storms of 1842, with only a general recollection of the similarity of the circumstances of the storm of December 1836, described in Mr. Loomis's previous memoir, which I had not looked into for some months. On its reperusal since, I find a condensed view of the facts of that storm at once so graphical, and by its accordance with the description drawn from the two other storms exemplifying so well their common character, that I am induced to insert it.

“The principal characteristics were as follows:—After a cold and clear interval with barometer high, the wind commenced blowing from the south. The barometer fell rapidly, the thermometer rose, rain descended in abundance. The wind veered suddenly to the north-west, and blew with great violence: the rain is succeeded by hail or snow, which continues but a short time; the barometer rises rapidly; the thermometer sinks as rapidly. These changes are not experienced everywhere simultaneously, but progressively from west to east.”

Such then are the phenomena, and such the order of their occurrence, in a class of storms which in the winter season, and in the localities referred to, are of frequent occurrence; that which has been described as the normal state of the atmosphere, and that which has been described as the interruption to it, appearing to follow each other in repeated succession. The facts being thus before us, it is for meteorologists to consider of their explanation.

When it is remembered that the temperature of the surface water of the Gulf of Mexico which washes the southern shores of the United States is considerably higher than the ordinary temperature of the surface water of the ocean in the same parallel; and that the gulf-stream which coasts the south-eastern states conveys heated water into parallels where its relative difference from the ordinary ocean-temperature is even greater than in the Gulf of Mexico, we should be prepared to expect that the abnormal southerly and south-



easterly winds should be *extremely* humid as well as warm; whilst the normal westerly wind, which has crossed the rocky mountains as well as a wide extent of continent, must be *extremely* dry as well as cold. Now the warm and moist air being once conveyed to the previously cold and dry localities where the storm appears to originate, the subsequent order and succession of the phænomena are sufficiently intelligible. The fact of which it seems most difficult to render an explanation, and to which therefore attention may be profitably directed, is that of the apparent tendency of the south and south-easterly winds to insinuate themselves in the lower stratum of the air, and to prevail over the regular and normal west wind, whenever the latter has moderated after its temporary violence. The phænomenon is confined to the lower stratum of the air, as the direction of the upper clouds is preserved steadily from the west. Mr. Loomis suggests in explanation that the momentum which the westerly wind acquires at its period of violence causes it to overblow itself, and produces a reaction, each storm having thus, as he conceives, a direct tendency to produce its successor. Is it not possible that the elastic force of the vapour rising over the heated surface of the ocean to the south and south-east of the United States, and making its way to the dry interior of the continent, may have a tendency to impede and counteract the current of air proceeding from an opposite direction? It is not inconsistent with the notion of the independence of air and vapour when at rest, that when in motion either should affect the other. It is I believe a common opinion that air in motion carries vapour with it; the supposition here made is the counterpart of this. I remember to have heard that at Newfoundland,—where the north-west (the prevailing) wind is particularly cold and dry, and where the surface of the sea to the south-east is of unusually high temperature for the latitude, owing to the gulf-stream, the sea fog, as it is called,—frequently makes its way from seaward against the wind; and that the wind then gradually dies away and is succeeded by a gentle breeze from the opposite or sea quarter.

But whatever may be the fate of conjectures which may be hazarded before the true explanation of the phænomena shall be arrived at and generally accepted, the very clear and lucid manner in which Mr. Loomis has arranged and combined the facts which he has collected together, and the ability and true philosophical spirit in which he has discussed them, call for our grateful acknowledgements, and cannot fail to operate as a stimulus to the co-operators in the United States to persevere in their meteorological observations. Mr. Loomis has ex-

pressed an earnest wish that the co-operation should be extended towards the north into the countries occupied by the Hudson's Bay Company; and it cannot fail to be seen, on reading his memoir, how much observations in that quarter are wanted for the elucidation of questions which arise. We may hope that his wishes in this respect will not be disappointed.

Believe me, sincerely yours,

Woolwich, February 12th, 1846.

EDWARD SABINE.

XXXVII. *On the Anthracite and Bituminous Coal-Fields in China.* By RICHARD COWLING TAYLOR, F.G.S.\*

WE have seen the recent announcement of the sailing, from hence, of a vessel containing 308 tons of Pennsylvania anthracite, destined for Hong-Kong in China. Some very natural speculations have arisen from this circumstance, as to the probability of that remote country furnishing a market for American anthracite. As no details accompany the statement alluded to, we are not in possession of any material facts whereby an estimate can be formed of the probable success of the undertaking, in a commercial sense; and we are not sure but the coal may have been employed for convenience merely, as ballast.

In the East Indies various depots of European coal have been established, for the service of the British government steamers. This fuel, for the most part, it is understood, consists of the anthracitous and partially bituminous coals of South Wales, of course obtained at great expense. It appears that 5000 tons of English coal, at a freightage of about £2 per ton, are annually imported into Bombay, for the Company's steamers. Bituminous coals have been derived from much less distant sources; among which the Burdwan coal-field, in the vicinity of Calcutta, may be named. Mergui Island, also, in the Bay of Bengal, has lately furnished some steam coal to Singapore. The steam ships on the China seas, during the war with that vast country, were supplied from these various sources.

I do not propose to discuss the profitableness, or otherwise, of a Chinese market for our American anthracite. But as during the process of collecting statistical information for a proposed volume on "The Geological and Geographical Distribution of Coal and other Mineral Combustibles †," some

\* From a pamphlet communicated by the Author.

† See in *Philosophical Magazine*, vol. xxvi. p. 263, the Prospectus of this work, by Mr. R. C. Taylor, for which subscriptions are received by Messrs. Wiley and Putnam.—ED.

notes reached me, of an interesting character, which are not generally accessible to the majority of readers, with relation to the Chinese coal-fields, it has struck me that a portion of these details, in an abridged form, might be just now acceptable, particularly as the intercourse with that country is on the increase. I venture even to omit, for the present, the authorities for the facts I shall have to communicate; reserving them in detail for the volume adverted to. It must, nevertheless, be premised that to the Jesuit Fathers, the French Missionaries who were permitted to reside at Peking during the 18th and preceding centuries, we are indebted for details of the highest interest, not alone on this subject, but on many other objects of philosophical inquiry in that little-known region.

It is probable that coal was discovered, and was in common use in China, long before it was known in the western world. It is mentioned by a noble traveller of the 13th century, as abounding throughout the whole province of "Cathay," of which Peking is the capital, "where certain black stones are dug out of the mountains, which stones burn when kindled, and keep alive for a long time, and are used by many persons, notwithstanding the abundance of wood."

The good missionaries were fully capable of describing the coals which were supplied to Peking, since they there erected a furnace or stove, in which they experimented on the properties of those combustibles; particularly with reference to the ordinary domestic uses, and for the warming of apartments and the purposes of their laboratory.

Among the people of Peking three kinds are in use.

1. That employed by the blacksmiths. It yields more flame than the other qualities; is more fierce, but is subject to decrepitate in the fire; on which account, probably, the blacksmiths use it pounded in minute particles.

2. A harder and stronger coal, used for culinary purposes, giving out more flame than the other sorts so employed; it is less quickly consumed, and leaves a residuum of gray ashes. There are several gradations of these. The best are hard to break, of a fine grain, a deep black colour, soiling the hands less than the others. It sometimes is sufficiently siliceous to give fire with steel. Others have a very coarse grain, are easily broken and make a bright fire, leaving a reddish ash. Another species crackles or decrepitates when first placed on the fire, and falls down, almost entirely, in scales, which close the passage of the air, and stifle the fire.

3. A soft, feebly burning coal, giving out less heat than the 2nd class; consuming more quickly, it breaks with greater facility, and in general is of deeper black than the sorts previ-

ously mentioned. It is commonly this description which, being mixed with coal-dust and a fourth part of clay, is employed to form an artificial is economical fuel. This being moulded in the form of bricks and balls is sold in the shops of Peking. Wagon-loads of coal-dust are brought to that city for this sole purpose.

The coal merchants have also an intermediate quality between the classes 2 and 3.

We cannot in this place recite the numerous details which are furnished by these intelligent Fathers. Suffice it to add, that nearly the whole of the properties and applications are now in every-day use in the United States, and are familiar to all. They are, in fact, the natural results suggested by qualities possessed in common by the combustibles of remote parts of the same globe. Even the modern method of warming all the apartments of our dwellings, which we view as the result of superior practical and scientific investigation, was in use, with very little deviation, centuries ago by the Chinese. Many a patented artificial fuel compound both in Europe and America, has been in practical operation in China at least a thousand years.

4. ANTHRACITE.—Another description of coal abounding about thirty leagues from Peking, but which was not then in such general use there as the other kinds, is called by the Chinese Che-tan. Che means a stone, but tan is the name they give to wood-charcoal. Therefore, according to the genius of the Chinese language, this compound word signifies a substance resembling or having the common properties of stone and charcoal. There can be little difficulty here in recognising the variety of coal which in our day has been denominated anthracite, a compound word of similar meaning.

The Chinese *glance coal* forms a remarkable exception to the unfavourable conclusion prevailing against Oriental coal; and, according to more recent authority than those we before cited, deserves to rank at the head of the list, in respect of its purity as a coke, although in specific gravity it does not come up to the character of the Pennsylvania or Welsh fuel; neither has it the spongy texture which contributes much to the glowing combustion of the latter.

So late as 1840, a Russian officer has described the coal formations of the interior, as occupying the western mountain range of China, in such abundance that a space of half a league cannot be traversed without meeting with rich strata. The art of mining is yet in its infancy among the Chinese; notwithstanding which, coal is thought to be at a moderate price in the capital. Anthracite occurs in the western range of

mountains at about a day's journey, or thirty miles only from Pekin. The coal formation is largely developed, in which thick beds of coal occur. They appear to be of various qualities. Some of this coal, occurring in shale beds, is singularly decomposed, and its particles have so little cohesion, that they are almost reduced to a state of powder. Beneath these coal shales are beds of ferruginous sandstone, and below those occur another series, consisting of much richer seams of coal than the upper group.

In this range are seen also both horizontal and vertical beds of conglomerate, accompanied by seams of coal which have the conglomerate for the roof and diorite or greenstone for the floor. As might be expected, this coal very much resembles anthracite. It is shining, of compact texture, difficult to ignite, does not flame in burning, or give out any smoke. Its substance is entirely homogeneous. Every thing respecting it leads to the belief that there had been a great development of heat at the period of its formation, or subsequently. The horizontal coal beds are the most important and valuable, and are denominated large; but no greater thickness than three and a half feet is quoted. The blacksmiths and those who work in copper, prefer this coal, on account of the intense heat which it gives out.

Throughout the whole of this mountain range may be continually seen the outcrops of this combustible, where they have never, as yet, been touched by the hand of man.

In those parts of China where wood is very dear, coal is worked on a great scale for the Pekin market: but the process of mining is very little understood by those people, who excel in the preparation of charcoal.

*Coal in other parts of China.*—The Missionaries and others inform us that coal is so abundant in every province of China, that there is perhaps no country in the world in which it is so common. The quays at Nankin are stored with the finest native coal. Some of the coal which was brought down to the coast, from the Pekin country, to the Gulf of Pe-tchee-lee, was anthracite, partaking of the character of plumbago or graphite. Coal, apparently of the brown coal species, exists extensively in the direction of Canton; while all the coals seen on the Yang-tse-kiang river, south of Nankin, resembled canal coal. Nearer to Canton it possessed the comparatively modern character of the brown coal. It was abundantly offered for sale in the different cities through which Lord Amherst's embassy passed, between the lake Po-yang-how and Canton, and the boats were largely supplied with it. It is there obtained by means of pits, like wells; and we infer that, like

nearly all the brown coal deposits, the beds were horizontal, and at no great depth. A sulphurous coal, interstratified with slate, and in the vicinity of red sandstone, also prevails towards Canton.

Thus, therefore, we possess evidence, the main object which this communication was designed to exhibit, that extending over large areas in China, are beds of tertiary or brown coal, of cannel coal, a dozen varieties of bituminous coal, of anthracite, glance coal, and graphitic anthracite; all of which, for ages, have been in common use in this remarkable country, and have been there employed for every domestic purpose known to civilized nations of all times; including gas lighting, and the manufacture of iron, copper, and other metals.

*Mode of Mining Coal in China.*—It might be expected that in China, where most of the practical arts have from time immemorial been carried on with all the perseverance of that industrious people, the operations of mining coal would be conducted with some regard to science, in relation to sinking, draining, and extraction. We have, however, good authority, especially in regard to the environs of Peking, for stating that the process is still in a very imperfect state. Machinery to lighten labour is there unknown. They have not even an idea of the pumps indispensable to draw off the water. If local circumstances allow, they cut drainage galleries; if not, they abandon the work whenever the inundation has gained too far upon them. The mattock and shovel, the pick and the hammer, are the mining instruments—the only ones, in fact, which the Chinese employ in working the coal. The water of the mine is emptied by the slow process of filling small casks, which are brought up to the surface by manual labour. Vertical shafts are not used. In working horizontal coal seams, the timbering is expensive, and the materials cost about two copecs per poud, = \$8,50 per ton, English wood being sold by weight in China.

The coal, when mined, is put into baskets and drawn upon sledges, which are raised to the surface by manual strength. Each basket contains about three pouds of coal, and one man can raise about eight baskets in a day. This is equivalent to 1032 Russian pounds, or to 12 cwt. English per day. The miners' wages are at the rate of 30 copecs a basket; which is equal to 240 copecs (copper currency), or 46 cents of United States currency, per day; being \$0,76 U. S. per ton.

*Prices at Peking.*—At the pit's mouth, this coal is sold for 60 copecs per poud, = \$4,63 per ton of 20 cwt. It is then conveyed on the backs of mules, through the mountains, and thence on camels to Peking, where the price is 1½ rouble,

=  $1\frac{1}{2}$  franc, = 29 cents United states, per poud; which, if our calculation be correct, is equivalent to \$11,60 United States, or £2 8s. 3d. per ton of 2240 pounds English. We perceive, therefore, that the best of fuel is expensive at Pekin, and hence the necessity for resorting to the artificial compounds and substitutes to which we briefly alluded.

There is, however, a kind of coal sold in that city at a much lower price, particularly when it is mixed with one-half of coal-dust. This coal, in 1840, sold for one rouble per poud, which is at the rate of \$7,75, = £1 12s. 3d. per ton. It is of indifferent quality, however; giving out but little heat, and is quickly consumed.

The compound fuel, consisting of coal-dust and clay, is still prepared after the mode described by the Missionaries last century; but its use is chiefly confined to the indigent classes.

*Coal Gas Lighting in China.*—Whether, or to what extent, the Chinese artificially produce illuminating gas from bituminous coal, we are uncertain. But it is a fact that spontaneous jets of gas, derived from boring into coal-beds, have for centuries been burning, and turned to that and other economical purposes. If the Chinese are not manufacturers, they are, nevertheless, gas consumers and employers on a large scale; and have evidently been so ages before the knowledge of its application was acquired by Europeans. Beds of coal are frequently pierced by the borers for salt water; and the inflammable gas is forced up in jets twenty or thirty feet in height. From these fountains the vapour has been conveyed to the salt-works in pipes, and there used for the boiling and evaporation of the salt; other tubes convey the gas intended for lighting the streets and the larger apartments and kitchens. As there is still more gas than is required, the excess is conducted beyond the limits of the salt-works, and there forms separate chimneys or columns of flame.

One cannot but be struck with the singular counterpart to this employment of natural gas, which may be daily witnessed in the Valley of the Kanawha, in Virginia. The geological origin, the means of supply, the application to all the processes of manufacturing salt, and of the appropriation of the surplus for the purposes of illumination, are remarkably alike at such distant points as China and the United States. Those who have read, even within the present month, the account of the recent extraordinary additional supply of gas, and the services it is made to perform at the Kanawha salt-works, must be impressed with the coincidence of all the circumstances with those which are very briefly stated in the previous paragraph in relation to China. In fact the parallel is complete.

To the coals and combustible minerals of China, I cannot further advert here. But what a conviction irresistibly presses upon the mind, as to the incalculable utility of the *railroad system*, and coal-mining improvements, in such an empire! If ever there were concentrated at one point all the circumstances especially and unequivocally favourable to that system, and imperiously calling for improvements of the character suggested, it seems to be presented in the case of the city of Pekin. Here, with its enormous population of 1,500,000 souls, it is situated only at a day's journey—computed at thirty miles—from an immense region of coal, comprising several varieties. Yet its inhabitants cannot purchase the best qualities of this coal, brought from the mountains on the backs of mules and camels, under \$11,60 per ton, and the very worst for less than \$7,75 per ton.

Without making unnecessary or invidious comparisons, it might not unreasonably be suggested, that a Pekin railroad, in connection with the coal mines, would be a far more profitable enterprise in its results, than the transportation of American coals to China.

I will only add one circumstance, which had nearly escaped me. Borneo, “the largest island in the world,” which is only twenty degrees due south of Canton, has lately come into repute for the great quantity of coal which it contains, not only accessible to ships along the coast, but extensively occurring in the mountains of the interior. Much information has also been acquired from the natives; and the facts which are already elicited are regarded as of considerable importance, in respect to the facilitating the steam navigation in the China seas. Philadelphia will, of course, have her share in the enlarged commercial intercourse with China. Would it, then, be asking too much of those who are personally interested in this improving trade, to communicate any additional facts, which are either unknown to, or have been omitted by, the author of these scanty notes?

Respectfully,

Philadelphia, April 28, 1845.

RICHARD C. TAYLOR.

NOTE.—The prices and admeasurements which are quoted in the foregoing article, were reduced to the United States and English currencies and measures from the Russian, as furnished by the Engineer Kovanko; who, in like manner, converted them into the Russian from the Chinese standards. In consequence of this triple conversion of standards, additional care has been taken to avoid error in these calculations.



XXXVIII. *On the Conversion of the solid Ferrocyanide of Potassium into the Sesqui-ferrocyanide.* By C. F. SCHÄNBEIN\*.

**I**N a former notice I have shown that a solution of the yellow prussiate of potash in water, placed in contact with an atmosphere of ozone, instantaneously destroys the latter, and is converted into the red sesqui-ferrocyanide. Since that communication was made I have ascertained that even the solid yellow salt very readily absorbs ozone and is changed into the red one. If a crystal of the common prussiate is suspended in a balloon containing an atmosphere strongly charged with ozone, and kept in that state by means of phosphorus and water, it will soon assume the colour peculiar to the red cyanide, just in the same way as it would do when held in air containing chlorine. The surface of the crystal, after having remained in the ozonized air for about twelve hours, is changed into the red salt, which may be easily separated from the yellow nucleus by mechanical means. A crystal of about a cubic inch in bulk appeared after thirty-six hours' suspension in ozonized air covered with a crust of the red cyanide, at least one line thick; and in another case I saw a smaller crystal of the yellow salt entirely converted into the red one. I hardly need say that by changing the yellow compound into the sesqui-ferrocyanide, the cohesive state of the former undergoes a material alteration. The red crust surrounding the yellow nucleus is rather brittle, and consists of a heap of small crystals of the sesqui-ferrocyanide. It is worthy of remark, that under the circumstances mentioned the yellow prussiate becomes moist, and exhibits in that state a very strong alkaline reaction.

XXXIX. *On the Decomposition of the Yellow and Red Ferrocyanides of Potassium by Solar Light.* By C. F. SCHÄNBEIN\*.

**A** SOLUTION of the yellow prussiate of potash kept in the dark does not change its colour, but when exposed to the action of solar light it becomes of a deeper yellow. To render that change very perceptible, a weak, *i. e.* nearly colourless, solution must be used, in which case the liquid will assume a yellow colour after having been acted upon by strong sunlight only for a few minutes. If the bottle containing the solution be closed and not quite filled with the liquid, an odour of prussic acid is perceptible, and at the same time

\* Communicated by the Author.

a reddish yellow sediment subsides, which seems to be the peroxide of iron. The decomposition of the cyanide takes place much more rapidly when strips of filtering paper or linen are immersed in a solution of the salt and exposed to the action of solar light. In a very short time that part of the strip turned towards the sun becomes yellow, whilst the opposite side remains colourless, or nearly so. If strips of paper moistened with the solution of the common prussiate of potash are closed up in glass bottles containing air, they also turn yellow by exposure to the sun, and a strong smell of prussic acid is perceptible in the vessels after a short time. In the shade no such action takes place. A large piece of linen cloth drenched with a solution of the yellow salt, after having been exposed in the open air to the action of solar light for thirty-six hours, had turned deeply yellow, and yielded, when treated with distilled water, a deep yellow solution, which on being filtered and heated to boiling became turbid, and deposited flakes of peroxide of iron. The same solution exhibited a stronger alkaline reaction than the solution of the common prussiate does. From the facts stated, it appears that the yellow ferro-cyanide is decomposed by light into prussic acid, oxide of iron and potash, and a compound formed yielding with water a yellow solution. Is that compound carbonate of potash and peroxide of iron; and do the constituent gases of the atmosphere take part in the decomposition besides the solar light? Further experiments must answer those questions. A limpid solution of the red cyanide also becomes turbid when exposed to the action of solar light, prussic acid being evolved and peroxide of iron thrown down.

*XL. A Reference to former Contributions to the Philosophical Magazine, on Physical Optics. By Prof. POTTER, A.M., F.C.P.S., late Fellow of Queen's College, Cambridge, &c.\**

**W**ITHOUT the slightest wish to interfere in the controversies of others, I now beg to refer the readers of the *Philosophical Magazine* to my papers in the *Magazines* for January 1840 and May 1841. In the former, at page 20, I have shown Mr. Green's formula for the intensity of reflected light to fail entirely as a representation of nature; and in the latter I have shown the peculiar refraction near the optic axes of biaxial crystals not to be represented by Sir William Hamilton's analytical deductions from Fresnel's equation to the wave surface in biaxial crystals.

\* Communicated by the Author.

The anonymous correspondent Jesuiticus in the last Number, refers to those analytical researches triumphantly in favour of the undulatory theory of light. I do not write to disturb the philosophical opinions of Jesuiticus, but to remind the readers of the Magazine where they will find the discussion of the points referred to.

**XLI.** *On Differentiation as applied to Periodic Series: with a few Remarks in reply to Mr. Moon.* By J. R. YOUNG, Professor of Mathematics in Belfast College\*.

**I**F in the general expression at p. 430 of my paper on Periodic Series, in the last volume of this Journal, A be made equal to  $-1$ , we shall have the identity

$$\frac{1}{2} = \cos \theta - \cos 2 \theta + \cos 3 \theta - \&c. \pm \frac{\cos (n+1) \theta + \cos n \theta}{2(1 + \cos \theta)};$$

and if we multiply this by  $d\theta$ , and integrate, we shall further have

$$\frac{\theta}{2} = \sin \theta - \frac{1}{2} \sin 2 \theta + \frac{1}{3} \sin 3 \theta - \&c.$$

$$\pm \int \frac{\cos (n+1) \theta + \cos n \theta}{2(1 + \cos \theta)} d\theta.$$

Now it is demonstrable, from other and independent principles, that, when  $n$  is infinite, the right-hand member of this equation, omitting the integral, is the true development of  $\frac{\theta}{2}$ , for all values of  $\theta$  not exceeding  $\pi$ . Hence we may infer that, for  $n = \infty$ , this integral is necessarily zero. If we suppress it therefore, we shall commit no error in the expression for  $\frac{\theta}{2}$ ; but a very considerable error will be introduced if we

attempt to derive from that expression, thus limited to the particular case of  $n = \infty$ , a series of other equations, by the aid of differentiation, as is commonly done. If the evanescent integral be restored, we may then apply the process of differentiation as far as we please: our resulting equations will all be identical equations; holding, whatever be the value of  $n$ , and supplying the necessary corrections of those erroneous developments which, in the case of  $n = \infty$ , are so commonly met with in analysis.

I have elsewhere observed that differentiation fails to be applicable to the series

\* Communicated by the Author.

$$\frac{\theta}{2} = \sin \theta - \frac{1}{2} \sin 2 \theta + \frac{1}{3} \sin 3 \theta - \frac{1}{4} \sin 4 \theta + \dots$$

in the isolated case of  $n = \infty$ ; and it is plain that in this case  $\frac{1}{n} \sin n \theta$  is 0, though  $\sin n \theta$  is itself indeterminate; the indeterminateness is therefore rendered nugatory. But if differentiation be allowed, this indeterminateness reappears in  $\cos n \theta$ , in the right-hand member of the result, though the left-hand member remains determinate, which is absurd. Still we are not precluded from applying differentiation to the general forms above, since these are universally true; they comprehend *all* values of  $n$ , and are *identical*. It is in virtue of this identity, and of this alone, that the results of differentiation may safely be extended to  $n = \infty$ , although for this isolated value of  $n$  differentiation be inapplicable.

I have very little to say in reference to Mr. Moon's attack in the last Number of this Magazine. The papers which have called it forth,—whether justly or not, I leave others to determine,—Mr. Moon confesses that he does not understand; and humiliating as such a confession may seem, the whole tenor of his remarks shows that he is sincere.

I beg to say, that I did not write expressly for Mr. Moon, and Mr. Moon therefore cannot reasonably expect that I should attend to his demand, and define the terms I use. I have employed nothing but the recognised language of analysis, and I cannot undertake to encumber the pages of this Journal with a glossary of scientific terms for Mr. Moon's especial benefit; if he will only take the trouble to turn to the Penny Cyclopædia, Mr. De Morgan will fully instruct him in all these things.

The occasion of my mentioning Mr. Moon's name was this:—I found Mr. Moon, in the June Number of this Magazine, floundering amidst difficulties which he showed himself unable to cope with. I had long previously contemplated a paper, of which the main object was to remove those difficulties, and in drawing it up for this Journal, I could not well avoid the mention of Mr. Moon's name. But I mentioned it with the most scrupulous courtesy and respect; I was especially anxious on this point, on account of the peculiarities which Mr. Moon had so often displayed in his published communications; so anxious indeed was I to avoid offence, that—at the risk of losing all credit for discrimination—I even went the length of calling him “an able contributor to this Journal!” As I have already said, I did not certainly write

expressly for Mr. Moon, but the instruction conveyed to him through my short papers, was precisely that of which he obviously stood in need. Instead of accepting this with thanks, he ungratefully turns round and bites the hand that brings him aid; and, not content with this, he is ungenerous enough, and unjust enough, to say, that everything in those papers, which is not erroneous, has already been given by himself!

Belfast, February 9, 1846.

J. R. YOUNG.

[We omit the remainder of Mr. Young's letter, in which he animadverts upon Mr. Moon in terms which the communications of the latter seem well-calculated to provoke. The same discretion has been exercised with regard to some parts of Mr. Moon's letter in the present number, as from the character which the controversy has assumed, we are not disposed to devote any more of our space to its continuance.—EDIT.]

## XLII. Mr. MOON in Reply to Jesuiticus\*.

**A**FTER the notice which appeared in the last Number of this Journal respecting his previous papers, there would be an obvious impropriety in the writer of the following remarks attempting to force on the Editors of the Magazine any matter which would tend to produce further discussion† on the subject to which he has of late called attention, except so far as he be driven to do so in self-defence. As, however, the views to which the Editors afforded the means of publication have been openly attacked in this Journal, their author conceives he has a right to say a few words in their behalf.

An anonymous writer, who subscribes himself Jesuiticus, commences certain animadversions on my first paper on Fresnel's Theory of Double Refraction, by the remark, that "the hypothesis on which Fresnel's Theory of Double Refraction is based is the following:—'That the displacement of a molecule of the vibrating medium in a crystallized body is resisted by different elastic forces according to the different directions in which the displacement takes place.'"

He then proceeds to make some remarks on the reasonableness of this hypothesis, which it is not my present purpose to dispute; but I must beg to observe, *en passant*, that the above is *not* the hypothesis on which Fresnel's Theory of

\* Communicated by the Author.

† We omit some portions of Mr. Moon's communication, where he appears to us to have lost sight of his declared purpose of confining himself to self-defence, and has introduced matter "tending to produce further discussion."—EDIT.

Double Refraction is based. It rests on a lower level still. The true basis of the theory is, that the æthereal medium consists of particles separated by finite intervals (to use a well-known, but improper mode of expression), and acting upon each other by their mutual attractions. From this principle, the so-called fundamental hypothesis of Jesuiticus is a sufficiently easy inference: I have thought it necessary to remark upon this inaccuracy, however, as from the extraordinary want of precision of the writers on this subject, it is somewhat difficult to say what is their real starting-point; at the same time, that in order to make a proper estimate of Fresnel's theory, and of the skill and judgement with which he has worked it out, it is very desirable that that fact should be clearly ascertained.

Jesuiticus afterwards goes on to say, "It is then proved, that if any particle of the æther be suddenly displaced, *the other particles remaining quiescent*, the force of restitution developed by such disturbance will not in general be in the direction of the displacement, but only when such displacement is in the direction of the aforesaid axes of elasticity. The elegant demonstration of Mr. Smith, quoted by Mr. Moon, is by Mr. Moon's own showing fully adequate to establish the theorem as I have enunciated it, which is doubtless the sense in which Fresnel conceived it."

I admit that Mr. Smith's demonstration is fully adequate to establish the theorem as Jesuiticus has enunciated it, but I must beg to assure Jesuiticus, that unless the demonstration establishes a great deal more than the theorem so enunciated, it is not, for the purpose for which it is adduced, worth the paper it is written upon. What is the use of considering the impossible case of a single particle suddenly disturbed while all the other particles remain quiescent, and then reasoning upon what takes place in the *beginning of the motion in that case*, as if the same held good *throughout the whole motion in the actual case*, when all the particles are vibrating together, when it is perfectly certain that it does not?

Jesuiticus says, "Any one who understands the subject must at once acknowledge that any theory of light must be, to a considerable extent, imaginative; and that theory which can explain the greatest number of facts ought to claim the attention of the philosopher more than any other." Of the justice of the remark contained in the first part of the above sentence, Fresnel's theory is no doubt a remarkable confirmation; in the sentiment of the second clause of it I am disposed to concur, with the reservation that some portion of the credit due to a theory depends on its antecedent probability. But

mark what follows:—"It is to this that the undulatory theory owes its great celebrity, and of all parts of the undulatory theory, that of double refraction is the most extraordinary. It ought to be regarded as a stupendous monument of human ingenuity. It must not be forgotten how admirably the properties of uniaxal crystals follow from the general investigation of the biaxal class; but above all, how from this same investigation, conical and cylindrical refraction were discovered by Sir William Hamilton."

I would ask of Jesuiticus, what is the hypothesis upon which Fresnel professes to explain the separation of the ray? Whether it is not substantially what I have stated it to be in the early part of this paper? And if so, I appeal to the world whether I have not shown incontrovertibly in my two papers on this subject contained in the last two Numbers of the Philosophical Magazine, that Fresnel entirely fails to explain the separation of the ray on that hypothesis. It may be true that some of Fresnel's expressions for the disturbance in doubly refracted and other polarized waves may involve in them certain elements of truth (though for my own part I should be sorry to answer for any of them); but they do not on that account afford any evidence of the truth of his principles, for this plain reason, that they do not follow from them. It may happen, that from the ruins to which this great theory must soon, if it be not already reduced, may be gathered some useful fragments which may form part of a new and more durable edifice; but Jesuiticus may take my word for it, or if he do not choose to do that, he will not have long to wait for the verification of the prediction, that the time is at hand when Fresnel's theory will be considered as a "stupendous monument" of anything else but ingenuity. As to the supposed discoveries of conical and cylindrical refraction, if Jesuiticus had been aware of their very doubtful character, he would hardly have ventured to have brought them so prominently forward.

\* \* \* \*

As to the investigation which I examined in the first of my two papers, I do not doubt that Mr. Airy considered it merely as an illustration; but even in that point of view, and without adverting to the error which I pointed out in his reasoning, it would be entirely worthless, as it is obvious that the state of things he contemplates could only exist for a single moment, whereas the results he deduces are supposed to be always subsisting. His object is to show that an undulation consisting of transversal vibrations might be propagated according to a certain law, when even on his own premises it is quite obvious that if the disturbance originally communicated

were of that character, it would immediately cease to be so, or in short, that a transversal undulation (if I may be permitted the expression) would not be propagated according to any law. With a full sense of the value of Mr. Airy's contributions to other departments of science, I cannot shut my eyes to the fact, that by allowing such investigations as the one under consideration (which but for its adopted parentage would not be worth a comment) to pass not merely without censure, but with apparent sanction, he has introduced an absence of precision,—a laxity of principle (so to speak) into mathematical inquiries, which has produced the most injurious effects both in the mixed and the pure sciences.

But to come to the error which Jesuiticus imagines he has found in my reasoning. He says, “that in substituting for  $u$ ,

$$u - \frac{du}{dx}h + \frac{d^2u}{dx^2} \frac{h^2}{1.2} - \&c.$$

and for  $u'$ ,

$$u + \frac{du}{dx}h + \frac{d^2u}{dx^2} \frac{h^2}{1.2} + \&c.$$

the substitutions stopping at  $h^2$ , merely require that  $h$  should be small in comparison with the *length of a wave*, not in respect to  $u$ .”

It is true that if we suppose the initial disturbance to be represented by  $\alpha \sin \frac{2\pi}{\lambda}(vt - x)$ , the substitutions stopping at  $h^2$  are defensible on the ground suggested by Jesuiticus; but does Jesuiticus conceive that when Mr. Airy wrote out this demonstration, he ever thought about the length of the wave, or any other circumstance connected with the initial vibration? If he does, I can only say that he is a very extraordinary person. In my paper I took Mr. Airy's investigation for what it purported to be, namely a proof that a certain hypothesis as to the disposition of the particles and the nature of their mutual action, without reference to the form of the initial disturbance, leads to the conclusion that transversal undulations may be propagated; and in that point of view I have no hesitation in saying it entirely fails; and, independently of all others, on the ground I have pointed out, *i. e.* of false approximation. If Jesuiticus has any doubt as to whether Mr. Airy did or did not consider himself to have proved the proposition generally, I would recommend to his attention Art. 127 of Mr. Airy's Tract, in which he takes the general integral of the equation



$$\frac{d^2 u}{dt^2} = \frac{1}{n} \left( 1 - \frac{1}{2^{\frac{3}{2}}} \right) \frac{m}{h} \frac{d^2 u}{dt^2}$$

to wit,  $u = \phi (rt - \mu x),$

which he would not have been justified in doing if he had not proved the general proposition. \* \* \* \*

Liverpool, February 10, 1846.

XLIII. *Proceedings of Learned Societies.*

ROYAL SOCIETY.

Jan. 15, "ON the Viscous Theory of Glacier Motion." By 1846. James D. Forbes, Esq., F.R.S. &c. Part II\*. "An attempt to establish by observation the Plasticity of Glacier Ice."

The two first sections of the present memoir are occupied with a critical examination of the theory advanced by De Saussure to account for the progressive motion of glaciers, which he considered as formed of masses of rigid and inflexible ice, and with the further explanations of that theory given by Ramaud, Bischoff, Agassiz, and Studer. The author, on the other hand, regarding these masses as possessing a considerable degree of plasticity, explains on that supposition the phenomena they present; and, in the third section of the paper, he relates a series of experiments which he carried on in the Mer de Glace, near Chamouni, in the summer of 1844, with a view to determine by direct measurement the relative motion of different parts of the glacier. This he accomplished by selecting a spot on the western side of the Mer de Glace, between Trelaporte and l'Angle, where the ice was compact and free from fissures, and erecting on the surface a row of posts at short distances from one another, in a line transverse to the general direction of the moving mass. He was thus enabled to discover by trigonometrical observations the movements of different points in this line; and he ascertained that they advanced more and more rapidly in proportion as they were distant from the sides of the glacier; and that when not under the influence of neighbouring *crevasses*, these motions were gradual and uninterrupted; as was shown by the lines carried through the posts forming, after the lapse of a few days, a continuous curve, of which the convexity was turned towards the lower end of the glacier.

"An Account of the Southern Magnetic Surveying Expedition." By Lieut. H. Clerk, R.A., in a letter to Lieut.-Colonel Sabine, R.A., F.R.S. Communicated by Lieut.-Colonel Sabine.

The letter, which is dated from the Magnetical Observatory at the Cape of Good Hope, June 28, 1845, reports the return to the Cape of the Pagoda from her voyage to the high southern latitudes after the successful completion of the magnetical service on which she had been employed by direction of the Lords Commissioners of the Admiralty, at the request of the President and Council of the Royal Society.

\* An Abstract of Part I. will be found at p. 538, vol. xxvi., of this Journal.

January 22.—“On the Supra-renal, Thymus and Thyroid Bodies.” By John Goodsir, Esq. Communicated by Richard Owen, Esq., F.R.S. &c.

In this paper, the author enters on the development of the theory he advanced two years ago with regard to the origin and nature of the supra-renal, thymus and thyroid bodies, and the correctness of which, with certain modifications, he has been enabled to confirm by subsequent observation and reflection. His hypothesis was that the three organs in question are the remains of the blastoderma; the thyroid being the development of a portion of the original cellular substance of the germinal membrane grouped around the two branches of the omphalo-mesenteric vein; the supra-renal capsules, the developments of other portions grouped around the omphalo-mesenteric arteries; and the thymus, the development of the intermediate portion of the membrane arranged along the sides of the embryonic visceral cavity. He has since ascertained, however, that the thyroid body derives its origin in a portion of the included *membrana intermedia* remaining in connexion with anastomosing vessels between the first and second aortic arches, or carotid and subclavian vessels. He considers these organs as essentially similar in their structure, as well as in their origin in continuous portions of the blastoderma situated along each side of the spine, and extending from the Wolfian bodies to the base of the cranium: the development of the supra-renal capsules having relation to the omphalo-mesenteric vessels; the thymus, to the jugular and cardinal veins and ductus Cuvieri; and the thyroid gland, to the anastomosing branches of the first and second aortic arches. The functions of these organs he regards as being analogous to those of the blastoderma; with this difference, however, that as the blastoderma not only elaborates nourishment for the embryo, but absorbs it also from without, that is, from the yolk, the developed organs only elaborate the matter which has already been absorbed by the other parts, and is now circulating in the vessels of the more perfect individual.

January 29.—“On the Use of the Barometric Thermometer for the determination of Relative Heights.” By James R. Christie, Esq. Communicated by S. Hunter Christie, Esq., Sec. R.S., &c.

The objects of this communication, as stated by the author, are, first, to show the theoretical foundation of the very simple law pointed out by Professor Forbes, according to which the difference of the boiling temperature of water at two stations is connected with their difference of level; and next, to test the accuracy of this law by a comparison of results deduced from his own observations on the boiling-point of water at different stations among the Alps of Savoy, Piedmont and Switzerland, with the heights of the same stations as determined by other observers and by different means; and thus to arrive at a just conclusion with respect to the value of the barometric thermometer as an instrument for determining differences of level.

Combining DeLuc's formula reduced to English units,

$$b = \frac{99}{\cdot 899} \log 10 \beta - 60\cdot 804,$$

where  $b$  is the variable boiling-point on Fahrenheit's scale and  $\beta$  the corresponding barometric pressure, with the formula of Laplace for the determination of the difference in level of two stations from barometric observations, he obtains the formula

$$H = 547.99 (b - b') \left\{ 1 + (t - 32^\circ) \cdot 00222 \right\},$$

where  $b$  and  $b'$  are the boiling-points on Fahrenheit's scale at the two stations,  $t$  the mean temperature of the air at the stations, and  $H$  their difference of level in English feet.

The author describes the particular instrument he employed in his observations, and his mode of determining the correction which it required; and then gives, in a table, the observations he made on the boiling-point of water at thirty-eight different stations in the Alps; the heights of the corresponding stations above the sea level, deduced from these observations; and, for the purpose of comparison, the heights of the same stations deduced by other observers. The difference between these and some of the author's results are considerable; but as they are not greater than would probably arise from ordinary barometric measurements, and as there is a close accordance between his results and the determinations on which the greatest reliance can be placed, he concludes that the results are on the whole satisfactory. Considering it, however, desirable to obtain some test of the accuracy of each observation independently of the rest of the series, the author avails himself of the barometric observations made at the Observatory at Geneva and at the Convent of the Great St. Bernard; and determining from these the corresponding temperature of boiling water, deduces the difference of level between each of his stations and these two places considered as fixed points: the sum of the height above Geneva and the depression below the Great St. Bernard should in all cases be the difference of level between the two fixed stations. Although there are here again considerable discrepancies, yet in most cases, where the height of the station may be considered as well-established, the height deduced from the observations agrees with it in a very remarkable manner.

In another table, the author gives the difference of level between the Observatory at Geneva and the Great St. Bernard, deduced from the recorded observations at those places simultaneous with his own at his various stations; and then remarks that the differences of height determined by the two methods do not differ from one another, in any single case, by so large a quantity as do the greatest and least differences of height deduced from the barometric observations; while in many cases the accordance is almost perfect.

The conclusion drawn from the comparisons in these tables is, that the barometric thermometer is capable of affording highly accurate and satisfactory results, perhaps even more so than the common form of barometer, but that there is considerable uncertainty attached to its indications. This uncertainty, far from being wholly attributable to the imperfections of the instrument as a measure of the atmospheric pressure, might, the author thinks, arise from an

extreme susceptibility to rapid changes in that pressure, which remain unindicated by the more sluggish barometer.

“On the Decomposition and Analysis of the Compounds of Ammonia and Cyanogen.” By Robert Smith, Esq., Ph. D. Communicated by Captain William Henry Smyth, R.N., F.R.S.

This paper is divided into four parts; the first relates to the decomposition of ammonia and its compounds; by the compounds of chlorine, and the collection and measurement of the nitrogen gas which is disengaged, the amount of which the author considers as furnishing a ready and accurate mode of estimating the quantity of ammonia in the solution subjected to analysis. The chloride of lime was the salt usually employed for this purpose: this method is regarded by the author as being peculiarly applicable to the analysis of organic substances.

The second part treats of the decomposition and estimation of hydrocyanic acid and its compounds by means of chloride of lime, yielding nitrogen gas and carbonate of lime; a process which occupies but a few seconds. In some cases, the employment of chloride of soda is preferable to that of chloride of lime, on account of the solubility of all the compounds that are formed. The author found the same method applicable also to the analysis of the salts of cyanogen; for the cyanides of the alkalies are decomposed by it as rapidly as the pure acid itself. The ferro-cyanides are also very readily decomposed.

The author, in the third part of his paper, relates the results of his trials of the hypochlorites as agents for the decomposition of uric acid, which proved so satisfactory as to induce him to believe that these salts might be advantageously used as solvents of uric calculi in the living bladder. He also proposes the employment of chloride of lime as a ready and accurate mode of estimating the quantity of nitrogen contained in urine, from the amount of gas disengaged by its action on the nitrogenous compounds. In the last part, the apparatus used in the experiments is described.

“On a point connected with the dispute about the invention of Fluxions.” By Augustus De Morgan, Esq., M.A., F.R.A.S., &c. Communicated by Samuel Hunter Christie, Esq., Sec. R.S., &c.

An assertion made by Sir Isaac Newton in a letter to Conti, published in Raphson's History of Fluxions, that the materials of the *Commercium Epistolicum* were “collected and published by a numerous Committee of gentlemen of *different nations*, appointed by the Royal Society for that purpose,” appeared to be at variance with the list of the Committee as it was appointed on the 6th of March, 1711-12, and which only contains the names of Arbuthnot, Hill, Halley, Jones, Machin and Burnet, who were all English. But on further search of the records of the Society with the aid of Mr. Weld, the Assistant Secretary, the author ascertained that other members were subsequently added to the Committee, among whom were Bonnet, the Prussian minister, and De Moivre, both of whom were foreigners; thus showing that the imputations which might have been cast on Newton's veracity are groundless.

February 5, 1846.—“On the Secretary Apparatus and Function of the Liver.” By C. Handfield Jones, M.D. Communicated by Sir Benjamin C. Brodie, Bart., F.R.S.

The author is led by his researches into the minute structure of the liver, to results which confirm the view of Mr. Bowman, in opposition to those of Mr. Kiernan on this subject: and particularly with regard to the absence of real tubercular ducts from the interior of the lobules. He concludes that the secreting process commences in the rows of epithelial cells surrounding the central axis of the lobule, and that the fluid there secreted is transmitted to the cells forming the margin of the lobule, where it is further elaborated, and, by the bursting of these cells, is conveyed into the cavity of the surrounding duct. A few diagrams are annexed, illustrative of the descriptions of microscopic structure given in the paper.

“An Account of some Experiments on the Electro-Culture of Farm Crops.” By Mr. William Sturgeon. Communicated by S. Hunter Christie, Esq., Sec. R.S., &c.

Grass grown on a parallelogram of land, fifty-five yards long by twenty-two yards wide, enclosed by underground wires, was found to be much more abundant than in any other part of the field; especially in a plot “upwards of fifty yards long, whose breadth was within the wires, and nearly at right angles to the axis of the parallelogram.” This plot of grass was principally on the western side of the wires, and extended but a very little way on the eastern side. The axis of the wire-enclosed parallelogram was in the magnetic meridian.

“On the Comet of 1844–45.” By John Collingwood Haile, Esq. Communicated by Charles Terry, Esq., F.R.S.

The author gives a series of observations, accompanied by a diagram, made by him at Auckland, in New Zealand, on the comet of 1844–45, which there appeared on the 20th of December 1844 and disappeared on the 30th of January following, having been visible forty-two days. Its most remarkable feature was that during its greatest brilliancy, the nucleus was not surrounded by the nebulous matter, but was situated at the very extremity of the head, and at times even appeared quite detached.

---

ROYAL ASTRONOMICAL SOCIETY.

[Continued from vol. xxvii. p. 307.]

June 13, 1845.—Observations of the Solar Eclipse of 1845, May 5, and of the Transit of Mercury of 1845, May 8, in a Letter from W. Lassell, Esq.

“Starfield, Liverpool, 10th June, 1845.

“I send you such observations as I have been able to make of the late solar eclipse and transit of Mercury, for which the weather was, in some respects, but very unfavourable.

“May 5, 1845. With a very unpromising sky I prepared to ob-

serve the first contact of the solar eclipse, by placing that part of the sun's limb (then very indistinct) which the moon would first touch, between the parallel threads of the micrometer, applied to the nine-foot equatoreal, with a power of ninety-six times.

“As the time of contact approached the sky somewhat cleared, and the moon's first impression took place at  $23^{\text{h}} 14^{\text{m}} 37^{\text{s}}.8$  sidereal time, or  $20^{\text{h}} 18^{\text{m}} 24^{\text{s}}.17$  mean time at the observatory. During the greatest part of the obscuration the sky was very cloudy, but towards the end it cleared, and the last contact was well observed at  $1^{\text{h}} 33^{\text{m}} 48^{\text{s}}.9$  sidereal, or  $22^{\text{h}} 37^{\text{m}} 12^{\text{s}}.45$  mean time. No phænomena beyond what is usual occurred; nor was there, to my senses, any perceptible diminution of light on the landscape.

“May 8. For the transit of Mercury the appearances a short time before it began were still more unpromising. During the forenoon we had several showers, with a most gloomy sky; and even as late as half-past three P.M. we had a smart shower of close, small rain. A little change for the better occurred shortly before four, and I had just time to set the micrometer by the sun's limb after he became visible, and get settled at the telescope, when the first notch was cut out by the planet. The sun's limb was beautifully sharp, but occasionally obscured by passing clouds. From the time, however, of the first impression until the planet had advanced about two of its diameters upon the disc of the sun, it was generally unclouded, and the atmosphere remarkably tranquil. The first contact took place at  $7^{\text{h}} 13^{\text{m}} 36^{\text{s}}.3$  sidereal time, or  $4^{\text{h}} 8^{\text{m}} 12^{\text{s}}.36$  mean time. The internal contact, or complete immersion of the planet, took place at  $7^{\text{h}} 16^{\text{m}} 48^{\text{s}}.7$  sidereal, or  $4^{\text{h}} 11^{\text{m}} 24^{\text{s}}.24$  mean time. Both times were carefully and, I believe, accurately noted. Whilst the planet was traversing the *edge* of the sun, an apparent distortion took place, the parts of the sun's edge, or limb, in contact with the planet, appearing *rounded off*; and a moment or two before the complete immersion of the planet, an appearance analogous to *Mr. Baily's beads* took place,—the planet apparently breaking contact two or three times with the sun's limb before the final separation occurred. Mercury had also, to my eye, somewhat of a pear-like shape previously to his entering quite within the sun's disc. When he had advanced two or three of his diameters, the clouds rapidly thickened, and I saw him no more.

“I take this opportunity of stating, that a late redetermination of the longitude of my Observatory depending upon the lately determined longitude of the Liverpool Observatory, inclines me to adopt finally  $11^{\text{m}} 47^{\text{s}}.34$  as my longitude west of Greenwich, which differs scarcely a quarter of a second from that given in my paper contained in the forthcoming volume of the Society's Memoirs.

“The latitude I have also redetermined lately by transits of seven stars over the prime vertical, giving  $53^{\circ} 25' 3''.5$  as the mean result.”

Observations of the Transit of Mercury made at Aylesbury by Thomas Dell, Esq. Communicated by Dr. Lee.

“*Transit of Mercury.*—The first contact of the two limbs was in-

visible, the sun being obscured by heavy clouds; but almost immediately afterwards it broke through them, and the interior contact of the limbs was well observed at 4<sup>h</sup> 18<sup>m</sup> 33<sup>s</sup> mean time. The sun was covered by light fleecy clouds, through which it was distinctly visible, and the discs of both the sun and Mercury were most beautifully and sharply defined until 5<sup>h</sup> 12<sup>m</sup>, when the whole sky became densely overcast, and continued so until after sunset. The time of interior contact is, I believe, accurate to a second, as the error of the chronometer had been determined at the sun's transit at noon."

*Mathematical Society.*—After the conclusion of the business of the Ordinary Meeting, a Special General Meeting was held to take into consideration a subject, of which due notice had been given to the Fellows by the following circular:—

“Somerset House, June 5th, 1845.

“SIR,—I have the honour of notifying to you, that in pursuance of a Resolution of the Council, passed on Friday, the 23rd of May last, a Special General Meeting of this Society will be held at the Society's apartments on Friday, the 13th day of June instant, immediately after the business of the Ordinary Meeting to be held on that day is concluded, for the purpose of taking into consideration and deciding upon a recommendation of the Council to suspend upon that occasion the Bye-laws relative to the Election of Fellows, and to elect as Fellows of this Society the remaining Members of the Mathematical Society (now reduced to nineteen in number, of whom three are already Fellows), without payment of the usual Admission Fees and Annual Contributions (or compositions in lieu thereof), the Mathematical Society having announced its resolution to transfer its valuable Library, with its Records and Memorials, to the Royal Astronomical Society.—I have the honour to be, Sir, your most obedient servant,

“ROBERT MAIN, Secretary.”

It was then moved by Professor De Morgan, and seconded by Mr. Galloway, and resolved unanimously,—

“That the recommendation of the Council in the circular now read be approved and adopted by this Meeting; and that, on the Library, Records, and Memorials of the Mathematical Society being delivered over to this Society, the remaining Members of the Mathematical Society be admitted Fellows of the Royal Astronomical Society without payment of the admission fees or annual contributions required by the Bye-laws.”

November 14.—The President announced that the whole of the books of the late Mathematical Society had been delivered over to this Society, and had been arranged by Mr. Stratford, who would acquaint the meeting with a few of the particulars.

Mr. Stratford stated that the books received consisted of

76	volumes	folio
622	..	4to.
1442	..	8vo.
311	..	12mo.

131 books not bound or catalogued; and that 6 volumes were yet to be delivered: that the Council had this day determined to complete the deficient sets of the most valuable works, to rearrange the library, and to prepare a new catalogue, uniting the books of the two societies as early as possible.

It was then moved by the Rev. R. Sheepshanks, seconded by Mr. Drach, and resolved unanimously, that the warm thanks of the meeting be given to Mr. Stratford, for the trouble which he had taken in behalf of the Society, in carrying into effect the resolution of the last meeting with regard to the Mathematical Society.

Sir J. Herschel exhibited to the meeting a model of the surface of the moon, constructed by Frau Hofrätthinn Witte, a lady resident in Hanover, from her own observations made with an achromatic telescope by Fraunhofer, placed in a small observatory on the roof of her dwelling-house, in that city. The model is composed of a mixture of mastic and wax, forming a globe 12 inches  $8\frac{1}{2}$  lines, Paris measure, in diameter, on which the positions and general outlines of the craters, and other remarkable features of the moon's surface, were in the first instance laid down from the latitudes and longitudes given by Messrs. Baer and Mädler in their work entitled *Der Mond*, and from their chart of the moon, and the modeling performed (with the aid of a magnifying-glass) from the actual appearance of the objects as presented in the telescope above mentioned. The globe in question is the ten millionth part of the actual diameter of the moon, in which proportion, therefore, the horizontal linear dimensions of the several mountains, &c. are laid down. But, in respect of the height, a double proportion is adopted, since otherwise the relative heights would have been with difficulty distinguishable on so small a model. Sir J. Herschel having explained the nature and mode of construction of this admirable work (of which only one other exists, now in the Royal Museum of Berlin—both being originals, and attempts to multiply copies by taking plaster casts having hitherto failed), pointed out several of the principal craters, and explained the nomenclature adopted by Messrs. Baer and Mädler in their work referred to, in describing the several characteristic peculiarities of the moon's surface. The model was, on the breaking up of the meeting, submitted to the closer inspection of the members.

December 12.—On a Direct Method of determining the Distance of a Comet by Three Observations. By J. J. Waterston, Esq.

The following is the author's explanation of his method:—

“It is well known that three observations of a comet afford sufficient data for computing its distance from the earth independently of any assumption as to the orbit in which it moves. The formula is, I believe, originally due to Lambert, and appears in the works of the principal mathematicians who have given analytical solutions of the problem by the differential method. It is unfortunate that the nature of the equation does not admit of much precision in the results of the calculation, which are consequently apt to be greatly affected by small errors of observation. The disturbing power of these unavoidable inaccuracies varies much according to the condi-



tions of the problem, and it is, perhaps, impossible to recognise it in the analytical expression without a much greater effort of the attention than can be given when merely computing an orbit. With good observations the method has the advantage of revealing any obvious tendency to an ellipse or hyperbola; and, besides, it will in most cases afford a useful approximation to begin with in computing the parabolic formulæ. As to the expediency of putting the observations to this preliminary test in all cases, there would, perhaps, be little difference of opinion, if the labour of computation in doing so were available in the last part of the process, and if the conditions upon which the degree of accuracy depends could be easily distinguished.

“The object of this paper is to submit to the Astronomical Society an account of a method which has occurred to me of solving the equation by means of a constant curve, and to show how the preliminary calculation may be made available in Olbers’s parabolic method, and likewise in a differential method, without requiring the original equatoreal position in either case to be transferred to the ecliptic. The conditions of accuracy also become so apparent in using this curve, that the effect of an error of right ascension or declination may be estimated by inspection.

“The method of solution is derived from the projection of the three observations on the plane perpendicular to the direction of the motion of the comet at the middle epoch. The earth’s orbit being projected, its deflection, caused by the sun’s attraction, is brought into view, and since its apparent direction is the same as that of the sun, and the projected direction of the sun from the comet is the same as at the earth, the radii vectores of both being identical on the projection; it is clear if the differentials at the middle time are alone considered that the deflection of the orbit of the comet, as it appears on the plane of projection, coincides in direction with the projected deflection of the earth’s orbit, and that its magnitude depends on a function of the angle at the comet. We thus obtain the means of forming an equation for the angle at the comet in terms of the deflection of the earth’s orbit; and this equation, although derived from a simple geometrical construction, appears to be similar to that which is given in the analytical discussion of the problem by Laplace, Lagrange, Legendre, and Airy. It depends wholly on the effect of the sun’s centripetal force during the elapsed time as it appears on the plane of projection; and, as this, in the short differential period of a few days, bears but a small proportion to the projection of its chord, or velocity, the results are much more liable to be affected by the unavoidable errors of observation than if the equation expressed the same unknown quantity in terms of the velocity. But in the last case we have to suppose the nature of the conic section known, in the first no assumption of the kind is required, the deflecting effect of the sun’s force being necessarily the same in all orbits at the same central distance.

“The equation for the angle at the comet is solved by drawing one line on the constant curve, and the preliminary computation required

to do so affords an expression for the ratio of the distances at the first and third observations on the usual assumption that the chord is divided in the ratio of the times.

“This expression may be converted into the elegant form given by Olbers, so that it is identical with the value of  $M$  in his formulæ, and is expressed in terms that are likewise required in drawing the line on the constant curve.

“An example is given from the Trevandrum observations of the great comet of 1843. The formulæ are also applied to Göttinger’s observations of the second comet of 1813. A copy of the constant curve is given upon a separate sheet, and the lines of these examples drawn. The co-ordinates of the curve consist of the cotangent and cube of the sine. It is easily constructed by the common tables. If drawn with ordinary care, it will give the reading of the angle at the comet to greater nicety than even the best observations can afford.

“I have appended a modification of Olbers’s formulæ for the radii vectores and chord adapted to equatoreal positions, and involving the use of the angular quantities already computed for the use of the constant curve. The additional work of computation does not appear to be so great as that which is required to convert the right ascension and declination into latitude and longitude; and, besides, it is easier to compare observations with the computed elements when the latter are referred to the equator. The inclination of the orbit and position of the nodes are transferred to the ecliptic by the solution of one spherical triangle.

“In the recent improvements which Olbers has made in his method, by expanding Euler’s formula into a series and reversing, the means are afforded of constructing a small table which shortens considerably the process of finding the distance by trial and error. Another improvement consists in the new expression given for the chord being more favourable to accurate computation. I have included a form of the same kind in terms of the right ascension and declination which is almost wholly made up of angular quantities that have already been prepared and used with the constant curve.

“In the last part of the paper an expression for the angle at the comet is given to be used with the differential method which, in solving by trial and error, requires only five tabular references.”

Extract of a Letter from Sir John Herschel to the President, dated Collingwood, November 29, 1845.

“Being on the subject of the satellites of Saturn, I will mention here a singularity which, though obvious enough, has not (so far as I am aware) been noticed before, viz. that the periodic time of the first satellite (first in order of the ring) is *precisely* half that of the third, and the periodic time of the second *precisely* half that of the fourth. This is far too remarkable and close a coincidence to be merely casual, and (the second satellite being a certainty) the extension of the law to the first (a law so out of the way and unlikely) would of itself be evidence of its real existence, even had it not been (as it now certainly has been) re-observed. If such atoms perturb one another’s motions, there must be some very odd secular equations

arising from this singularity. It is not worth while to make a formal communication of such a thing to the Astronomical Society; but if you think it worth your verbal mention at the meeting, it may be interesting to those (if any) who are busy about satellitary perturbations."

On a new Double-image Micrometer, communicated in a Letter to the President by Professor Powell.

"My dear Sir,—The following suggestion for a very simple double-image micrometer occurred to me a few years ago; but not having much practical acquaintance with these matters, I should hardly have supposed it to possess novelty or prospect of utility enough to render it worthy the notice of the Astronomical Society, had you not encouraged me to communicate it.

"The optical principle is merely that of a ray of light, refracted obliquely through a plate of glass with parallel surfaces, and emerging parallel to, but not coincident with, its original direction.

"If, then, such a plate intercept half the cone of rays going from the object-glass of a telescope to its focus, there will be formed, at the focus, besides the direct, a deviated image; and the angular deviation will be dependent on the inclination, the thickness, and the refractive power of the glass, involving a constant factor to be found by observation for the particular instrument, agreeably to the following formula, which may be easily tabulated for all inclinations.

"If  $\phi$  and  $\phi^1$  be the angles of incidence and refraction, and  $t$  the thickness of the plate, a moment's consideration will shew that the oblique path of the ray within the plate =  $t \cdot \sec \phi^1$ ; and, for the angular space  $\theta$  between the direct and the deviated ray,  $c$  being the constant for the instrument, we have

$$\theta = c \cdot t \cdot \sec \phi^1 \sin (\sin \phi - \phi^1).$$

"If such a plate be placed within the tube of a telescope between the object-glass and its focus, so that a variable inclination can be given to it, and a graduated circle be read off outside; then, when the plate is perpendicular to the axis there will be no deviation; but, when it is inclined, the deviation, found as above, will give the measurement of a small angular space, as in other double-image micrometers.

"The less the thickness of the glass, the greater will be the range of the scale for a very small deviation.

"The idea has as yet been put to trial only in a very rough manner; and I offer it without at all being able to say whether serious practical difficulties may not arise, which can only be decided on a more accurate construction; or should no such objection occur, it still remains to be seen whether this suggestion may afford any useful addition to the micrometrical resources already in the hands of the observer, so as at least to be available in some cases: but these are points on which the practical astronomer alone can judge; and it is mainly in the hope that it may receive such examination that I submit this idea to the Astronomical Society.—I remain, &c.,

"BADEN POWELL,

"Oxford, December 7th, 1845."

"Savilian Professor of Geometry."

XLIV. *Intelligence and Miscellaneous Articles.*

## EXPERIMENTS ON THE SPOTS ON THE SUN.

**A**T the meeting of the American Philosophical Society, June 20, 1845, Prof. Henry of Princeton, U.S., made the following communication of a series of experiments made by himself and Prof. Alexander relative to the spots on the sun.

His attention was directed to this subject, by an article in the September number of the *Annales de Chimie*, by M. Gautier, upon the influence of the spots on the sun on terrestrial temperature. It is well known that Sir William Herschel entertained the idea, that the appearance of solar spots was connected with a more copious emission of heat, and that the seasons during which they were most abundant were most fruitful in vegetable productions; and pursuing this idea, he was led to trace an analogy between the price of corn and the number of solar spots during several successive periods. The result of this investigation, so far as it was extended, seemed to favour the views of this distinguished philosopher. A mode of investigation of this kind, however, is not susceptible of any great degree of accuracy; the price of corn is subject to so many other causes of variation besides that of solar temperature, that little reliance can be placed on it.

M. Gautier has attempted to investigate the influence of the solar spots on terrestrial temperature, by comparing the temperature of several places on the earth's surface, during the years in which the spots were most abundant, with those in which the smallest number were perceptible. From all the observations collected, it seems to be indicated, that during the years in which the spots were the greatest in number, the heat has been a trifle less; but the results are far from being sufficiently definite to settle the question: and M. Gautier remarks, that a greater number of years of observation at a greater number of stations, will be necessary to establish a permanent connexion between these phænomena.

The idea occurred to Prof. Henry, that much interesting information relative to the sun might be derived from the application of a thermo-electric apparatus to a picture of the solar disc, produced by a telescope, on a screen, in a dark room. This idea was communicated to Prof. Alexander, who readily joined in the plan for reducing it to practice. It was agreed that they should first attempt to settle the question of the relative heat of the spots as compared with the surrounding luminous portions of the sun's disc. The first experiments were made on the 4th of January 1845. Mr. Alexander had observed, a few days previous, a very large spot, more than 10,000 miles in diameter, near the middle of the disc. To produce the image of this spot, a telescope of four inches aperture, and four and a half feet focus, was placed in the window of a dark room, with a screen behind it, on which the image of the spot was received. The instrument was placed behind the screen, with the end slightly projecting through a hole made for the purpose, and a small motion of the telescope was sufficient to throw the image of the spot off or on the end

of the pile. The spot was very clearly defined, and might have been readily daguerretyped, had the telescope been furnished with an equatorial movement. The form of the penumbra of the spot, as it appeared on the screen, was that of an irregular oblong, about two inches in one direction, and an inch and a half in the other. The dark central spot within the penumbra was nearly square, of about three-fourths of an inch on the side, and a little larger than the end of the thermo-pile.

The method of observation consisted in first placing, for example, a portion of the picture of the luminous surface of the sun in connexion with the face of the pile, and after noting the indication of the needle of the galvanometer, the telescope was then slightly moved, so as to place the dark part of the spot directly on the face of the pile, the indication of the needle being again noted. In the next set of experiments the order was reversed; the picture of the spot at the beginning of the experiment was placed in connexion with the pile, and afterward a new part of the luminous portion of the disc was made to occupy the same place.

¶ The thermo-electrical apparatus used in these experiments was made by Ruhmkorff of Paris; and in order to render the galvanometer more sensitive, two bar magnets, arranged in the form of the legs of a pair of dividers, were placed with the opening downwards, in a vertical plane, above the needle, so that, by increasing or diminishing the angle, the directive power of the needle could be increased or diminished, and, consequently, the sensibility of the instrument could be varied, and the zero point changed at pleasure.

In the present experiments, in order to mark more definitely the difference in temperature, after the needle had been deflected by the heat of the sun, the magnetic bars above mentioned were so arranged as to repel it back to near the zero point, so that it might, in this position, receive the maximum effect of any variation in the electrical current.

Twelve sets of observations were made on the first day, all of which, except one, gave the same indication, namely, that *the spot emitted less heat than the surrounding parts of the luminous disc*. The following is a copy of the record made at the time of the observations. The degrees are those marked on the card of the galvanometer, and are of course arbitrary.

Spot,  $3^{\circ}\frac{1}{4}$ .Sun,  $4^{\circ}\frac{1}{2}$ .Sun,  $3^{\circ}$ .Spot,  $1^{\circ}\frac{3}{4}$ .Spot,  $2^{\circ}$ .Sun,  $3^{\circ}$ .Sun,  $2^{\circ}\frac{1}{2}$ .Spot,  $2^{\circ}$ .Spot,  $2^{\circ}$ .Sun,  $2^{\circ}\frac{1}{4}$ .Sun,  $5^{\circ}\frac{1}{4}$ .Spot,  $4^{\circ}$ .Spot,  $4^{\circ}\frac{1}{2}$ .Sun,  $5^{\circ}$ .Sun,  $4^{\circ}\frac{1}{2}$ .Spot,  $3^{\circ}\frac{3}{4}$ .Sun,  $2^{\circ}$ .Spot,  $3^{\circ}\frac{1}{4}$ \*Spot,  $0^{\circ}\frac{3}{4}$ .Sun,  $2^{\circ}\frac{1}{2}$ .

\* At this observation a slight cloud probably passed over the sun's disc.

Spot,  $4\frac{3}{4}$ .  
Sun,  $5^{\circ}$ .

Sun,  $1\frac{1}{4}$ .  
Spot,  $0^{\circ}$ .

The change in the temperature during the intervals of observation, is due to the variations in the temperature of the room differently affecting the two extremities of the pile.

In consequence of cloudy weather, another set of observations were not obtained until the 10th of January, and at this time the spot had very much changed its appearance; the penumbra, while it retained its dimensions in one direction, was much narrowed in the other, and the dark part was separated into two small ones; also the sky was not perfectly clear, and therefore the results were not as satisfactory as those of the previous observations; the indications were, however, the same as in the other sets, exhibiting a less degree of heat from the spots.

Cloudy weather prevented other observations on the heat of different parts of the sun, particularly a comparison between the temperature of the centre and the circumference of the disc, which would have an important bearing on the question of an atmosphere of the sun. The observations will be continued, and any results of interest which may be obtained will be communicated to the Society.

---

METHOD OF PURIFYING OXIDE OF URANIUM FROM NICKEL,  
COBALT AND ZINC. BY PROF. WOHLER.

When the oxide of uranium, in its preparation from the pitchblende, has been so far purified as to be dissolved in carbonate of ammonia, sulphuret of ammonia is carefully and gradually mixed with the solution as long as a black precipitate falls. In this way nickel, cobalt and zinc are entirely separated, without any uranium being thrown down.—Liebig's *Annalen*, Oct. 1845.

---

ON SOME NEW DOUBLE HALOID SALTS. BY M. POGGIALE.

*Protochloride of Antimony and Chloride of Ammonium.*—The protochloride of antimony combines with chloride of ammonium in two proportions. When protochloride of antimony is added to a solution of that salt, it dissolves readily, and only a slight turbidness, arising from the formation of some oxychloride, is perceptible. On evaporating the liquid at a gentle heat, at first beautiful rectangular prisms are obtained, and subsequently hexahedrons or hexahedral pyramids. The first are  $3\text{NH}^3 \text{HCl}, \text{SbCl}^3 + 3\text{HO}$ , and the latter  $2\text{NH}^3 \text{HCl}, \text{SbCl}^3 + 2\text{HO}$ . Both salts are colourless and transparent; they become yellow and opaque in moist air, but are very permanent in dry air, and are coloured yellow in the mother-ley when heated; they are likewise decomposed by a large quantity of water.

*Protochloride of Antimony and Chloride of Potassium.*—This salt is deliquescent, becomes yellow on exposure to the air, is decomposed by water and also by heat; it forms laminar crystals, the composition of which is  $3\text{KCl}, \text{SbCl}^3$ . The mother-ley yields, on spontaneous evaporation, hexahedral crystals of  $2\text{KCl}, \text{SbCl}^3$ .

*Protochloride of Antimony and Chloride of Sodium* forms laminar crystals, having the composition  $3\text{NaCl}, \text{SbCl}^3$ .

*Protochloride of Antimony and Chloride of Barium*.—When the solution of the chloride of barium is very dilute, the two salts separate on cooling, the chloride of barium crystallizes in tablets, while the protochloride of antimony decomposes the water. It is therefore necessary, in order to obtain this compound, to use concentrated solutions. It is obtained in minute radiately-grouped needles, the composition of which is represented by  $2\text{BaCl}, \text{SbCl}^3 + 5\text{HO}$ . The protochloride of antimony combines in the same way with chloride of strontium, chloride of calcium, and chloride of magnesium.

*Protochloride of Tin and Chloride of Ammonium* forms beautiful fascicular needles, which are permanent in the air, but are decomposed by water. The analysis of this salt, which had been previously obtained by Jacquelin, gave the formula  $2\text{NH}^3, \text{SnCl}, \text{HCl} + 3\text{HO}$ .

*Protochloride of Tin and Chloride of Potassium* is obtained by direct combination of the two salts, and crystallizes in beautiful long needles, which are isomorphous with the preceding salts. Its formula is  $2\text{KCl}, \text{SnCl} + 3\text{HO}$ .

*Protochloride of Tin and Chloride of Barium* yields, on spontaneous evaporation, beautiful prisms, the *protochloride of tin and chloride of strontium* long needles. They are represented by the formulæ  $\text{BaCl}, \text{SnCl} + 4\text{HO}$  and  $\text{SrCl}, \text{SnCl} + 4\text{HO}$ .

*Chloride of Sodium and Magnesium* consists of  $\text{NaCl}, 2\text{MgCl} + 2\text{HO}$ .

*Iodide of Silver and Ammonium*.—Iodide of ammonium dissolves iodide of silver, forming with it a deliquescent double salt. It contains  $2\text{NH}^3, \text{HI}, \text{AgI}$ .

*Iodide of Lead and Sodium* crystallizes in yellow shining laminæ. It is obtained by adding a slight excess of iodide of sodium to a hot solution of iodide of lead, and placing the liquid in a warm spot. Its formula is  $\text{NaI}, 2\text{PbI}$ .

*Iodide of Zinc and Sodium* yields, on spontaneous evaporation, prismatic radiately-grouped needles. It is colourless, readily soluble in water and deliquescent. Its formula is  $\text{NaI}, \text{ZnI}$ .

*Chloride and Iodide of Lead* is obtained by dissolving iodide of lead in a solution of chloride of ammonium. On cooling, numerous yellowish crystals separate, which assume the form of needles, and consist of  $\text{PbI}, 2\text{PbCl}$ . On evaporating the mother-ley, crystals are obtained of  $2\text{NH}^3 \text{HCl}, \text{PbI} + 2\text{HO}$ , in the form of minute, silky, ramified needles; they become yellow in the air, and are decomposed by water.

*Chloride and Acetate of Lead*.—This is formed when chloride of lead is boiled in a porcelain dish with teracetate of lead, to which subsequently a slight addition of acetic acid is made. The solution is evaporated at a gentle heat, when colourless shining crystals separate on cooling. The salt has a sweetish astringent taste, effloresces readily in the air, and melts at  $180^\circ$ ; it loses its water of crystallization at  $228^\circ$ . The salt is readily soluble in water; alcohol

decomposes it, and precipitates chloride of lead. Several analyses yielded the formula  $PbCl, 5PbO C^4 H^3 O^3 + 15HO$ .

*Iodide and Carbonate of Lead* is prepared by digesting carbonate of lead with iodide of lead until the excess of iodide of lead has dissolved. This salt is yellow and insoluble in water. Its formula is  $PbI, PbO, CO^2$ .—*Comptes Rendus*, p. 1180.

ON THE VOLATILE ACIDS OF CHEESE. BY MM. ILJENKO AND LASKOWSKI.

The authors cut fifty pounds of Limbourg cheese, which possessed a very strong odour, into small pieces, mixed them with water, and submitted the mixture to distillation in a large alembic, water being occasionally added during several days. By this operation a somewhat turbid ammoniacal liquor was obtained, which was supersaturated with sulphuric acid and again distilled. The product was afterwards saturated with barytes water; the salt obtained was evaporated to its crystallizing point; the acid was again separated and converted into a salt of silver. Analysis showed that this volatile acid was entirely valerianic acid.

The residue was afterwards saponified by means of potash, the soap decomposed by potash, and subjected to a fresh distillation, and there was thus obtained an acid liquid which was saturated with barytes and evaporated to crystallize; it yielded a mixture of several salts of barytes, which were separated by means of their different solubility in water. The rough salt was mixed with seven parts of water and heated to boiling; the caproate of barytes dissolved, and afterwards separated in crystalline tufts of considerable size, whilst the butyrate remained in solution; this was converted into a salt of silver and analysed.

The barytic salts, which were not dissolved by the seven parts of boiling water, were composed of caproate and caprylate of barytes; and they also were separated by their different solubility.

It appears then that cheese contains the following volatile acids:

Butyric acid . . . .	$C^4 H^8 O^2$
Valerianic acid. . .	$C^5 H^{10} O^2$
Caproic acid. . . .	$C^6 H^{12} O^2$
Caprylic acid . . .	$C^8 H^{16} O^2$
Capric acid . . . .	$C^{20} H^{20} O^2$

Valerianic acid occurs in the largest quantity, and its presence had been previously discovered by M. Balard in the cheese of Roquefort. All these acids, it will be observed, are homologous substances.

The authors also performed some experiments on the fused portion of cheese; they obtained by means of boiling alcohol perfectly crystalline margarine from it; it was fusible at  $127^\circ$  Fahr., and margarinic acid was obtained from it. The rough margarine was mixed with some liquid glycerine. Unaltered caseine was also present, soluble in boiling water and insoluble in alcohol. There was also pre-



sent lime, a little magnesia, soda, potash, traces of iron, phosphoric acid, chlorine and sulphuric acid.—*Journ. de Pharm. et de Ch.*, Decembre 1845.

#### ON THE DOUBLE SALTS OF THE MAGNESIAN GROUP.

M. J. Isidore Pierre has paid particular attention to the salts of this group, including those of magnesia, oxide of copper, zinc, nickel, cobalt, manganese and iron.

The author observes, that it is well known that Prof. Graham has stated, with respect to the sulphates of the above-named bases, that one of the equivalents of water cannot be eliminated, except at a much higher temperature than is required for the others; that this equivalent may be replaced by an equivalent of a salt, so that the double salt formed contains one equivalent less of water than if each of the two simple sulphates had brought all its water of crystallization into the molecule of the double salt which results from their combination.

M. Pierre states that the results which he has obtained do not confirm those of Prof. Graham; he found that sulphate of zinc contains, as generally admitted, 43.72 per cent., or seven equivalents of water; and he ascertained that by exposing it for a long time to a temperature of 230° Fahr. and a current of dry air, that it lost 43.6 per cent., or the whole of its water, which is at variance with Graham's result, who found that it required a heat of 400° Fahr. to expel the seventh equivalent of water.

*Double Sulphate of Zinc and Potash.*—This salt is readily prepared by mixing together hot solutions of equivalents of sulphate of zinc and bisulphate of potash, and allowing crystallization to take place. The crystals are beautiful small, milk-white parallelogrammic tables; this salt is soluble in two and a half times its weight of boiling water, but much less soluble in cold water, for it crystallizes abundantly on the cooling even of an unsaturated solution.

When exposed gradually to a heat of 356° to 392° Fahr., it effloresces without fusing in its water of crystallization, which it loses completely and pretty rapidly at this temperature, the amount being 27.49 per cent. The author's analysis gives as the formula of this salt,  $ZnO, SO^3; KaO, SO^3 + 7HO$ , which indicates, as he shows, 27.32 per cent. of water instead of 24.49, the experimental result.

In this case it is therefore to be remarked, that the sulphate of zinc retains the seven equivalents of water which it possessed in its simple state.

*Double Sulphate of Zinc and Magnesia.*—M. Pierre observes, that it is generally supposed that these two sulphates may combine in all proportions; having found that sulphate of zinc in the double salt which it forms retains its seven equivalents of water, the author observes that if sulphate of magnesia did the same, the double salt should contain fourteen equivalents of water.

This salt is readily obtained by mixing its equivalents and crystallizing; it forms very fine oblique rhombic prisms, which are by

pressure separated into very fine needles; when quickly heated to  $212^{\circ}$  to  $248^{\circ}$  Fahr., it loses part of its water; at  $392^{\circ}$  Fahr. it retains two equivalents of water, and these cannot be expelled at a temperature lower than  $482^{\circ}$  to  $500^{\circ}$  Fahr.

When heated slowly and progressively, this salt effloresces without fusing in its water of crystallization; it may in this mode be deprived of the whole of its water of crystallization without being fused; it merely agglutinates slightly.

The formula of this salt derived from analysis is  $ZnO, SO^3, MgO, SO^3 + 14HO$ , which indicates 47.17 per cent. water; the loss of water by experiment was 47.12; the salt heated to  $392^{\circ}$  Fahr. retained ten equivalents of water.

From the preceding and the analyses of various other salts which the author prepared, he arrives at the following among other conclusions:—

1. Sulphate of zinc containing seven equivalents of water, retains the whole of it in the compound which it forms with the alkaline or alkalino-earthly sulphates.

2. The sulphates of zinc and magnesia combine equivalent to equivalent, and the resulting compound contains a quantity of water equal to the sum of the quantities which both salts contained when separate, that is to say, fourteen equivalents, if the double salt crystallizes at common temperatures.

3. The simple sulphates of zinc, copper and nickel yield all their water at a little above  $212^{\circ}$  in a long-continued current of air, instead of retaining one equivalent, as stated by Prof. Graham, at  $400^{\circ}$  Fahr.

4. The simple sulphates of zinc, magnesia, copper and nickel combine with other sulphates, or with each other without elimination of the water.—*Ann. de Ch. et de Phys.*, Fevrier 1846.


#### PREPARATION OF HYPOPHOSPHITES.

M. A. Wurtz prepared almost the whole of these salts, which he analysed, by the double decomposition of soluble sulphates with hypophosphite of barytes; the most economical method of preparing the last-mentioned salt, is to boil a solution of sulphuret of barium with phosphorus, until gas ceases to be evolved. If the ebullition has been long continued, the sulphuret of barium is almost entirely decomposed, and a slight excess only is left, which may be separated by carbonate of lead. Sometimes, however, the excess of sulphuret is more considerable; it is then proper to separate it by cautiously adding small quantities of sulphuric acid to the hot filtered liquor, as long as sulphuretted hydrogen is disengaged. If the liquor becomes acid, it must be neutralized, before evaporation, by a little carbonate of barytes.


*Hypophosphite of Potash.*—This salt was prepared by the double decomposition of hypophosphite of barytes and sulphate of potash. The aqueous solution was evaporated to dryness, and the residue, treated with hot alcohol, deposited hypophosphite of potash on

cooling. The crystals of this salt are hexagonal tables. They are very deliquescent, very soluble in weak alcohol, less so in absolute alcohol, and insoluble in æther. They lose no water at  $212^{\circ}$  Fahr.


Adopting with M. Pelouze 400 as the atomic weight of phosphorus, M. Wurtz gives as the formula of this salt,



*Hypophosphite of Ammonia.*—This salt was prepared like the preceding. It crystallizes in large irregular hexagonal laminae; it is less deliquescent than the salt of potash, and unalterable at  $212^{\circ}$  Fahr. At about  $394^{\circ}$  Fahr., it fuses into a transparent liquid without losing water, and becomes a crystalline mass on cooling. It does not decompose under  $464^{\circ}$  Fahr., at which temperature, like other hypophosphites, it disengages a little water and spontaneously inflammable phosphuretted hydrogen. Its formula, as determined by analysis, appeared to be




*Hypophosphite of Strontia.*—This salt was prepared like that of barytes, by boiling a solution of sulphuret of strontium with phosphorus, and decomposing the excess of sulphuret by carbonate of lead, or by sulphuric acid added in sufficient quantity to render the liquid slightly acid. By evaporation the hypophosphite of strontia crystallizes in the mammillated form by the juxtaposition of small laminae round a common centre. These crystals are unalterable in the air, and lose no water at  $212^{\circ}$ . They are very soluble in water, and insoluble in alcohol. The formula of this salt is



*Hypophosphite of Magnesia.*—This salt was prepared by double decomposition with sulphate of magnesia and hypophosphite of barytes; it crystallizes, as stated by M. H. Rose, in very brilliant regular octahedrons, which effloresce in dry air. The formula of the crystallized salt is  $PH^2O^3, MgO + HO + 5Aq$ ; of the salt dried at  $212^{\circ}$ ,  $PH^2O^3 MgO + HO$ ; and lastly, the formula of the salt dried at  $360^{\circ}$  Fahr., is  $PH^2O^3, MgO$ .

*Hypophosphite of Zinc.*—This salt was obtained of two different forms. It crystallizes sometimes in very efflorescent regular octahedrons, and sometimes in small rhombic crystals unalterable in the air. When a moderately concentrated solution of this hypophosphite is submitted to spontaneous evaporation, the first-mentioned crystals are usually formed. They are so efflorescent, that they lose water during pressure between folds of paper, previous to analysis. The formula of the rhombic crystals is



*Hypophosphite of Iron.*—This salt crystallizes in large green octahedrons, which effloresce by exposure to the air and become a white powder. When exposed to the air, the moist salt absorbs oxygen from it rapidly. The formula of the crystallized salt is  $PH^2O^3, FeO + 6HO$ .

*Hypophosphite of Chromium.*—This salt was prepared by double

decomposition by mixing solutions of sulphate of chromium and hypophosphite of barytes. By evaporation there was obtained an amorphous, cracked mass of a very deep green colour. This salt loses water by drying when it has been heated to  $392^{\circ}$  Fahr.; it is not soluble either in water or dilute acids. The formula of this salt is  $2\text{PH}^2\text{O}^3, \text{Cr}^2\text{O}^3 + 4\text{HO}$ .

*Hypophosphite of Manganese.*—The crystals of this salt are of a rose colour, brilliant and unalterable in the air. They do not lose water at  $212^{\circ}$  Fahr., but at about  $302^{\circ}$  Fahr. they part with one equivalent. The formula of this salt is  $\text{PH}^2\text{O}^3, \text{MnO} + \text{HO}$ .

*Hypophosphite of Cobalt.*—This salt forms large crystals of a deep red colour, which effloresce in the air. At  $212^{\circ}$  Fahr., they lose six equivalents of water of crystallization, and become a pale rose-red powder. The formula of this salt, which agrees with that of M. H. Rose, is  $\text{PH}^2\text{O}^3, \text{CoO} + 6\text{HO}$ .

*Hypophosphite of Nickel.*—The crystals of this salt are regular octahedrons, and smaller than those of hypophosphite of cobalt. When the aqueous solution is evaporated at the temperature of  $212^{\circ}$  Fahr., it is partially reduced to metallic nickel with the disengagement of hydrogen. This reduction takes place perfectly when the crystals of this salt, broken and slightly moistened, are exposed to the air at a temperature of  $248^{\circ}$  Fahr. The formula of this salt is  $\text{PH}^2\text{O}^3, \text{NiO} + 6\text{HO}$ .

*Hypophosphite of Copper.*—The solution of this salt is readily prepared by decomposing sulphate of copper with hypophosphite of barytes. It is not a permanent salt. At about  $140^{\circ}$  Fahr. it becomes turbid, and deposits hydrate of copper. By evaporating the solution *in vacuo*, small blue crystals of this salt were once obtained. These crystals decompose quickly, and with projection of the entire mass when heated to  $149^{\circ}$  Fahr.; and phosphuret of copper is formed. The formula of this salt is  $\text{PH}^2\text{O}^3, \text{CuO}$ .—*Ann. de Ch. et de Phys.*, Fevrier 1846.

---

#### BIELA'S COMET.

The following is an abstract of a letter addressed by Prof. Challis to the editor of the *Times*:—

“As I was preparing to observe Biela's comet, on the evening of the 23rd of January, I discovered a smaller comet in its immediate neighbourhood, and ascertained by my observations that evening that the two comets had the same apparent motion. A double comet is a celestial phænomenon which, I believe, has never before been witnessed, and cannot fail to arrest the attention of astronomers. It will be a matter of very great scientific interest to determine the relative motions of these two singular bodies, and the nature of the influence they mutually exert on each other. The following relative positions I have succeeded in obtaining by means of the Northumberland telescope. They are either derived from separate determinations of the places of the comets, or from direct measurements of angles of positions and differences of North Polar distance. The smaller comet is north of the other, and precedes it.

	Mean time. h	Difference of R.A. s	Difference of N.P.D. sec.	Angle of position. d. m.	Distance. sec.
Jan. 23	7·1	5·18	122·9	327·43	145·4
... 24	7·0	5·11	126·6	328·48	148·0
... 27	6·4	5·84	144·6	328·48	169·1
... 28	6·3	5·77	151·1	330·12	174·1
... 29	7·4	5·44	154·4	332·10	174·6
Feb. 11	7·4	7·38	249·7	336·8	273·1
... 12	7·1	7·86	252·7	335·3	278·7
... 13	7·5	8·30	264·9	334·54	292·6

“The observations on the 11th and 12th of February were obtained with great difficulty on account of the faintness of the comets from the effect of moonlight. The light of the larger comet spreads over a considerably greater extent than that of the other, but is not intrinsically much brighter.”

METEOROLOGICAL OBSERVATIONS FOR JAN, 1846.

*Chiswick.*—January 1. Fine. 2, 3. Frosty: fine; overcast. 4. Rain. 5. Sharp frost: cloudy: clear and frosty. 6. Drizzly. 7. Overcast and mild throughout the day and night. 8. Cloudy and fine. 9. Uniformly overcast. 10. Overcast: drizzly rain. 11. Hazy and drizzly. 12. Cold haze. 13. Hazy: very fine. 14. Foggy: overcast and fine. 15. Fine. 16. Thick fog: rain at night. 17. Hazy: drizzly: cloudy and mild. 18. Foggy: rain at night. 19. Constant rain: boisterous, with rain at night. 20. Clear and fine. 21. Rain: densely clouded and mild: boisterous, with rain at night. 22. Boisterous, with rain: densely clouded. 23. Heavy showers. 24. Hazy and mild. 25. Rain. 26. Showery: heavy rain at night. 27. Clear: cloudy: rain at night. 28. Rain: cloudy: very high tide in the Thames: clear. 29. Rain. 30. Overcast. 31. Cloudy: windy at night.

Mean temperature of the month .....	45°·54
Mean temperature of January 1845 .....	38·69
Average mean temperature of Jan. for the last twenty years	36·46
Average amount of rain for the last twenty years .....	1·60 inch.

*Boston.*—Jan. 1. Stormy: rain last night. 2. Fine. 3. Cloudy. 4. Rain. 5. Fine. 6. Rain. 7. Cloudy. 8. Fine. 9—13. Cloudy. 14, 15. Fine. 16. Foggy. 17. Cloudy: rain A.M. and P.M. 18. Foggy. 19. Rain: rain early A.M.: rain P.M. 20. Windy: rain early A.M. 21. Cloudy: rain P.M. 22. Cloudy and stormy: rain early A.M. 23. Fine. 24. Cloudy: rain early A.M. 25. Fine: rain early A.M. 26. Cloudy: rain early A.M. 27. Fine. 28, 29. Rain. 30, 31. Cloudy.—N.B. Not so warm a January since January 1834: the average of that month was 44°·3.

*Sandwick Manse, Orkney.*—Jan. 1. Snow-showers. 2. Fine: frost: cloudy. 3. Cloudy: clear. 4. Clear: showers. 5. Bright: showers. 6. Damp: clear. 7. Cloudy: showers. 8. Showers: clear. 9. Cloudy: clear. 10. Rain: cloudy. 11. Drizzle: damp. 12. Drizzle: hazy. 13. Bright: cloudy. 14. Damp: cloudy. 15. Rain: drizzle. 16. Clear. 17. Damp. 18. Bright: cloudy. 19. Damp: showers. 20. Rain: drizzle. 21. Rain: clear. 22. Damp: rain. 23. Fine: damp. 24. Fine: frost: damp: aurora. 25. Rain: cloudy. 26. Damp. 27. Damp: rain: clear. 28. Cloudy: showers. 29. Showers. 30. Cloudy: rain. 31. Drizzle: showers.

*Applegarth Manse, Dumfries-shire.*—Jan. 1. Snow-showers. 2. Frost: clear and fine. 3. Wet all day. 4. Fine A.M.: shower P.M. 5. Frost A.M.: rain P.M. 6, 7. Showery. 8. Fair. 9, 10. Slight drizzle. 11. Slight drizzle: fog. 12. Fair and mild. 13. Fair A.M.: rain P.M. 14. Fair: one slight shower. 15. Wet A.M.: cleared: fine. 16. Frost, slight: fine. 17. Fair A.M.: slight shower P.M. 18. Fair, but cloudy. 19. Rain nearly all day. 20. Rain all day: flood. 21. Fair, but cloudy. 22. Drizzling rain. 23. Rain and fog. 24. Thick fog. 25. Heavy rain: flood. 26. Drizzling rain. 27. Rain A.M.: fair: rain P.M. 28—31. Rain.



THE  
LONDON, EDINBURGH AND DUBLIN  
PHILOSOPHICAL MAGAZINE  
AND  
JOURNAL OF SCIENCE.

---

[THIRD SERIES.]

---

APRIL 1846.

---

XLV. *On the Oscillations of the Barometer, with particular reference to the Meteorological Phænomena of November 1842\**. By WILLIAM BROWN, Jun. †

[With Six Plates.]

IN order to illustrate and confirm the views I have before advanced in this Magazine (vols. xx. and xxiii.), on the connexion between the direction of the currents of the atmosphere and the oscillations of the barometer, I have endeavoured to make a direct application of them to the explanation of the various phænomena presented by the winds in this country during a great portion of the month of November 1842, that month including an extremely unsettled and stormy period of weather. For this purpose, I have collected observations showing the state of the wind and the barometer in various parts of this kingdom, and also at Christiania in Norway, and at Paris; and exhibited the direction of the wind on diagrams, with the variations of the barometer in hundredths of an inch annexed. The data from which the diagrams are constructed, and which are given p. 262, with the exception of those for North Shields, the place of my own register, are extracted from the Shipping and Mercantile Gazette Newspaper; a newspaper containing daily reports from most of the parts of Great Britain and Ireland, of the state of the wind and weather, on which nautical men are accustomed to rely for that kind of information; and the accordance of the observations at places situated near each other is sufficiently marked to confirm their general correctness, and thus give confidence in those of more isolated localities.

The barometrical observations have been collected with

\* This essay, with the exception of some additions, although only now published, was written soon after this period.

† Communicated by the Author.

great care: those for London and the Orkneys are taken from the tables published in the Philosophical Magazine and *Athnæum*, and those for Paris from the *Annales de Chimie*; but in general I have been indebted for them to the kindness of the observers themselves. I sought those from Christiania, which are from the register kept under the superintendence of Prof. Hansteen, in order to enlarge the field of the observations; it appears however from them that their locality is too far to the west to throw light on those of this country, except in a few instances; for this reason they are placed at the foot of the columns, contrary to the general order of the positions of the observations; and when reference is made to the observations as a whole, they are never included, except when specifically mentioned.

The whole of the facts brought to light by this investigation may, I think, be resolved into this general principle; that all winds may be ultimately referred to the action of one or both of two contrary currents caused by the unequal distribution of temperature on the surface of the earth; the one arising from the flow of colder and therefore denser air towards warmer, in the lower regions of the atmosphere; and the other from the descent to the surface of the earth of an opposite current belonging to the upper regions of the atmosphere, and formed by the elasticity or total weight of the atmosphere at any elevation in the warmer regions, being greater than that at the same elevation in colder, because of the greater height of the atmospheric column in the former than in the latter; the air in both being supposed of the same pressure at the surface of the earth. Thus in fig. 1, Plate V., the outline  $HbBA$  represents the general figure of a portion of the atmosphere, of which the temperature decreases from  $A$  to  $B$  ( $A$  lying on the equatorial and  $B$  on the polar side of  $C$ , or  $A$  being south and  $B$  north); the lower current of heavy air will therefore set in from  $B$  towards  $A$ . But the pressure being equal at the surface of the earth, and greater at any equal elevation in the column  $HA$  than in  $bB$ , at some certain height above  $A$  the pressure or elasticity of the air will be so much greater than at the same height above  $B$ , that its force will there overcome the pressure of the colder air of the column  $Bb$ ; and hence above this the air will flow from  $HA$  towards  $bB$ . But these currents can only be maintained by air descending and ascending in some part of them: let the upper current descend to the earth at  $A$ , it will continue to flow towards  $C$ , as represented in the figure, by the momentum acquired in its original position. These currents are the north and south winds.



From this may be deduced the following results, which may be applied to the explanation of the general atmospheric phenomena of all latitudes of both hemispheres, though when the cardinal points are referred to, they are given in terms adapted only to the northern, and to extra-tropical latitudes; for the sake of convenient reference to each, they are placed in numbered paragraphs.

1. An equality in the pressure of the atmosphere can only be maintained by the flow of these currents in their proper, that is, in their original positions, the one above and the other below; but on the descent of the upper current, which still, by its momentum, maintains wholly, or in part, its original direction, the lower is either more or less retarded, or entirely pushed back: thus whilst the air is carried away, either from the upper parts, or whole of the atmospheric columns, the flow to the lower is prevented, and consequently a diminution of air, or decrease in the atmospheric pressure, takes place in the regions where the descending current prevails on the surface of the earth: hence the great oscillations of the barometer in high latitudes, or the region of the "variable winds," and the maintenance of the equality of the atmospheric pressure in the region of the "trade winds," the ascent and descent of the air taking place at the extremities of the latter, and consequently without interruption to the course of the currents.

2. In the extent of the descending current, and for a space in front of it, following the line of its direction, the pressure of the atmosphere upon the surface of the earth will be distributed according to the curved line *adcb*, Plate V. fig. 1; and the lines *Aa*, *Dd* and *Cc*, &c. will represent the pressure at the several points on which they are drawn, *C* being supposed the point at which the descending current terminates and meets the opposite or lower one, as shown by the arrows, whilst the southerly current is blowing above the north from *c* to *b*; the minimum pressure will be near *C*, or the place of meeting of the two currents. That this will be the case is evident from the consideration, that the descending current advances by reason of the superiority of its force to that of the lower one, which it drives back; but this superiority is constantly diminishing by the rarefaction of the air produced by its flowing from *c* to *b*, which rarefaction at last reduces the force of this current below that of the resistance of the opposite one in front of it; hence it is clear that where the descending current terminates in advancing from the place of its descent, or is overcome, as at *C*, the pressure must be at the minimum, increasing from this point in both directions,

but in the greatest degree towards B, because of the resistance given to the upper current in its flow northwards after passing this point, and the increasing density of the air in the colder current. In my first paper on this subject (vol. xx.), I gave an illustration by a figure, in which for want of due consideration a rise of the barometer was conceived to occur at the point C; but in a subsequent paper on the "Storms of the Tropics" (vol. xxiii.), it is assumed that this is, as here expressed, the point of the greatest barometric depression, though I omitted to notice the discrepancy. As this part of the subject is of great importance, it may be proper to give a further explanation of it. The upper current is supposed to descend to the surface of the earth between A and C, on which it will flow to a certain distance dependent on its power to overcome the opposite one from B, and C is the point at which it meets this current and advances upon it; here therefore there will be an influx of air from both sides at the surface, whilst that above is carried away in a continuous current from *c g* to *b*. Now it will at once be evident that in this position of the currents, any change produced in the atmospheric pressure will depend upon their relative velocities; if that of the upper one is so much the greatest, that notwithstanding the check given it by the opposite force, more air is carried off from the higher parts of the atmosphere above C towards *b* than is brought to that point in the lower, the pressure at C must diminish. But the diminution of pressure thus begun by the force of the descending current, will go on until this force is reduced, by the loss of pressure sustained, to an equality with that of the opposite one, and then its momentum being destroyed it will cease to advance, and the latter will begin to advance upon it, to restore the equilibrium of the atmosphere; hence the point of its furthest advance and first cessation must be near that where the diminution of the atmospheric pressure is greatest, or the point C; and at this point, in great storms, there will be a comparative calm throughout a certain extent of the atmosphere. That the conclusion resulting from this reasoning is in consonance with observed fact, may be seen from the observations of P. J. Espy, who has shown that the space between the opposite sides of a storm is in reality the place of minimum pressure in those storms of America which he has investigated\*.

\* This, according to this observer, is the position of the fall of rain which occurs during storms, the fact upon which he has founded his theory; but it will be seen that in this case the conditions are precisely such as, according to the general opinion of meteorologists, are requisite to produce rain,—these are, the meeting of two currents in precisely opposite conditions

3. It is not meant however in the foregoing paragraph that the greatest depression of the barometer throughout the course

with regard to temperature and the quantity of aqueous vapour they contain; and hence the true nature of the connexion between the occurrence of rain and a falling barometer, both consequences of one common cause.

The phænomena attending the fall of rain are extremely dependent on geographical position, and are by no means sufficiently known to enable us fully to carry out these principles to the explanation of them, although I am firmly persuaded that when better known we shall be able to do so. I will however just refer to a few cases similar to that mentioned, of which they do give a sufficient explanation, and which are notorious weather laws:—the occurrence of rain;—just before a change of the wind, or at the time of the change, whether it be from north to south, or from south to north, though the most conspicuous in the former case; during a north-east wind with a falling barometer (§ 5), and with a south or south-east wind (occasioned by the junction of a north-east and south wind, see § 11). The last of these is the most conspicuous in the portions of storms to which § 16 refers, which in those parts where the wind is from south to south-east are always accompanied by abundance of rain. It may be thought that our dry winds from north-west (also formed by a north and south wind (§ 11)) are an exception to these results; but it is by no means necessary that rain should always occur at the meeting of these currents, for if the lower current greatly predominates in dryness or in quantity, then it is evident that there need be no precipitation of vapour in the form of rain. But there is another reason why these winds should be in general free from rain. The occurrence of rain in showers with squalls of wind, when the other portions of the day are fine, is a case to which the principle before us strikingly applies, for these squalls almost always blow in a direction somewhat different from that of the wind in the intervals between their occurrence; thus showing that they arise from an immediate onset of one or other of the opposite currents: now it is very easy to conceive that two bodies of air may meet so as to produce rain, although their relative temperatures and quantities of vapour may be so adjusted, that the resulting temperature is sufficient to maintain the same quantity of vapour; for if the collision be sudden, by the law of the diffusion of gases and vapours the vapour of the warm air will rush at once into the cold air, not waiting for the mixture to take place; and hence, being subjected to its temperature, it is immediately condensed and the rain is produced. Now (§ 11) the north-west wind is one of the most constant winds, hence one of the most favourably disposed for the gradual mixture of the opposite currents. This also explains the occurrence of fine weather with a steady barometer, for stability in the pressure of the atmosphere can only be produced by the stability of the currents. The formation and disappearance of clouds without rain may be explained in the same manner,—the precipitated vapour not being sufficiently dense to form rain is again aëriated when the cold air acquires the temperature of the mixture. [For a full description of the differences and relations of the distinct atmospheres of air and vapour by which the globe is surrounded, and on which this reasoning is based, I need scarcely refer the reader to the ‘Meteorological Essays’ of the late Professor Daniell, where they are set forth with great perspicuity and precision.]

But perhaps the fact most remarkably in accordance with this application of the principle set forth in this essay is that general one, established by W. Snow Harris by induction from a great number of particular instances, that thunder-storms result from the collision of opposite currents; for the

of a wind or storm\* is at once attained at the point C, which is there supposed to be that of the furthest extent of the descending current from the *first* place of its descent, and the place of the barometric minimum in the parts *then* occupied by the wind. The pressure continues to decrease for some time by the progressive motion of the storm. It is evident that air having begun to descend at A and to flow forward with considerable velocity in the same direction as before its descent, the lower air on the south of A must at once begin to flow towards A to supply its place; but as this air is either at rest or in a state of motion in another direction, it cannot at once begin to flow with sufficient velocity to supply the deficiency; therefore the rarefaction thus produced will cause the upper current to descend into it, and thus the space upon which it flows will gradually extend itself backward from A, or southward. But at the same time that this is going on behind A, the advanced portion of the descending current has begun to retreat from C; for its force there, at first superior to that of the opposite one, is at last overcome by it, and the heavy current of cold air then advances upon the receding wind, flowing with a force in some degree proportioned to the degree of the rarefaction, and restores the air to its ordinary pressure †, according to one general law of storms, that when the wind changes from south to north the barometer begins to rise. Thus the point of minimum pressure, C, or the furthest advanced portion of the storm, and the point of its first occurrence or descent, A, both move in one direction, from north to south, but not equally; for it is obvious that the portions of air which descend after the first have an advantage over the latter in this, that the opposition in front being partially removed, by the first portions and the diminution of pressure begun, they flow towards a rarefaction; and thus the force of the storm and the diminution of pressure at C are increased, and the motion of this point southward is retarded, whilst the wind is progressing on the south from A. But this disproportionate motion of the two extremities cannot remain, for with the increase of the rarefaction there is an increase of the force of the

torrents of rain which frequently fall during their occurrence seem to manifest that the only difference between this and the fore-mentioned cases is in intensity.

\* The only distinction here inferred between wind and storm is that of force; in many instances I use the term storm, because of the phænomena being sufficiently striking only when the wind has great force.

† In the hurricanes of the tropics, the returning current is a second storm, and it is sometimes so in those of high latitudes; but in the latter the rarefaction of the atmosphere is so extended, that the restoration of the pressure is frequently very gradual and produced by moderate winds.

north current, which will after a time carry backward the point C with increasing velocity, and gradually put a period to the storm\*.

This is more conspicuous in storms of temperate regions than in those of the tropics, because the former consist, as it were, of one deep wide depression of the atmosphere, the progressive motion of the storm being, so far as regards the movement from north to south, apparently in great measure an enlargement of this depression southward, whilst the latter occasion a much smaller one.

4. As a direct consequence of the foregoing, and as also proved by P. J. Espy, the greatest reduction of atmospheric pressure in storms is not where the wind is most violent, but where its velocity is reduced by the resistance in front of it; and the depression of the barometer at any given place depends on its position with regard to the place of the minimum, or the point C, fig. 1, as well as on the violence of the storm.

5. A further consequence of the same result is, that a considerable diminution of the pressure of the atmosphere, and consequently, fall of the barometer, takes place on the locality where a north-east wind is blowing, when this wind is immediately on the north of the northern range of the south wind which occasions the fall (as at the point E in the current from B towards C), though, as explained in § 2, to a less extent than in the localities occupied by the south wind. This result explains a fact of very frequent occurrence, the falling of the barometer during a north-east wind.

6. As the impetus of the south wind may have reduced the elasticity or pressure of the air, in the column Cc at any elevation of the atmospheric columns, below that of the air of the same elevation on the north; a very rapid increase of the pressure of the atmospheric columns on the north may give so great a check to the upper current at C, as to cause the air to accumulate so rapidly, that the increase of pressure or rise of the barometer extends to a great distance on the south; but it is yet evident, from the state of the atmospheric columns shown by the figure, that the south wind will continue to flow from A because of the greater pressure there; but the barometer will rise on account of the accumulation of air taking place at C and extending in a diminishing degree towards A; hence a frequent phænomenon, the rise of the barometer during the continuance of the south wind; as also

\* The deflection from south by the rotation of the earth is for the present left out of consideration, see § 15.

the beginning of the rise of the barometer, frequently some time previous to the setting in of the northerly current.

7. But an atmospheric pressure above the mean will also result from the opposition of these currents, but an opposition differing from that of § 2 in this respect; that whereas in that case the force of the descending current, originally much the greatest, is reduced at the place of its termination to an equality with that of the opposite one by the diminution of the pressure of the atmospheric columns composing it; in this, where a rise of the barometer takes place, the force of the lower, or current of gravity, is equal or superior to that of the descending one when at its full pressure, and the former is advancing upon the latter. Let then the two currents so circumstanced meet, as in fig. 1, at C, there will be at this station either simply a condensation of the air produced by the pressure of the two currents, or the air of the lower one will ascend, carrying with it an impetus which would tend to carry it on still in its first direction. In the case of § 2, air brought to the place of meeting is carried off in the upper current by the force of that current, but in this, if the air of the lower current does in this way ascend, it will simply check that flowing above in the contrary direction, and cause an accumulation of air to take place exactly similar to that of water occasioned by partially damming a stream. But even when the force of the former is in some degree inferior to that of the latter, it may yet be sufficient to retain so much air of the upper current by its opposition as to accomplish the same effect though in a less degree: thus the pressure will be represented by the dotted line *agb* in fig. 1, and an elevation of the barometer will ensue in the localities where the south wind is blowing, as well as in those which have the north; hence great elevations of the barometer occur with south winds as well as north.

8. The foregoing paragraph is intended to explain the great atmospheric pressure sometimes produced by strong north-east winds, and calms or very gentle breezes, as one or other of these is produced (§ 12) at places situated near the collision of the two currents, especially when the effect is increased, as shown in a previous essay, by reduction of temperature; but increase of pressure will arise from other causes; as in some locality sufficiently far to the north of a strong southerly wind to be out of reach of its influence in depressing the barometer, and upon which the air from the depression flows; though probably in this case the barometer will not rise in a great degree, on account of the air, whose removal causes a deficit of pressure in one locality, being so extensively

spread by the flowing of the upper current on others; the reason why elevations of a degree corresponding to depressions at any given place have never been found by comparative observations\*.

9. An elevation of the barometer may also be the consequence of a previous reduction; for let the pressure be reduced, as at C, fig. 1, the returning air or northerly current which sets in after the cessation of the southerly one, will occupy a portion at least of the higher regions of the atmosphere where it is not wont to flow, as is evident from the figure. On the restoration therefore of the usual pressure, the north current will be blowing not only in its own proper region, but also in part of that which properly belongs to the southerly one, and will continue there some time by reason of its acquired velocity after the original impulse has ceased to act; and thus the upper current, not resuming at once the whole of its action, and consequently the air not being allowed to flow from the upper parts of the atmospheric columns as rapidly as it is brought to the lower, will accumulate. In like manner, the elevation of the barometer may be the cause of giving to the upper current a great velocity, for an elevation being anywhere produced, the force by which the lower current causing it was urged on, must sooner or later be overcome by the increase in the pressure of the air towards which it flowed. But this current being overcome, the overplus of pressure will then increase the velocity of the upper one, and probably determine the flow of the air at the surface of the earth in the same direction; which indeed is frequently the way in which an elevation of the barometer subsides; and in the observations given in this essay, in which are included two periods of stormy weather, both began with the occurrence of a southerly storm after a high barometer.

10. The direction of the wind when one current alone prevails, is determined by the relative situations of the warm air and cold, and the deflection of the current thus produced, by the rotation of the earth, as the "trade winds" and monsoons of the tropics, and the north-east wind of high latitudes; but when the opposite currents come into collision, the direction is the resultant of their forces, and thus in the latter regions we have winds from every point of the compass, as has been pointed out by Prof. Kæmtz.

11. The action of these currents meeting together and producing the various winds may be considered as follows:—The south-west and north-east winds blowing from two stations, A and B, and meeting together at a station between them, C,

\* Daniell's Meteorological Essays.

may cause a wind in any direction, according to their force and their inclination to one another; if it be on the eastern side from any point of the compass between a north-eastern point and some point between south and west, it will be very liable to change; for winds from these points being produced by the direct collision of the descending current with that from north-east, when the latter is blowing on the surface of the earth from some more northern station, require the continuance and stability of a current whose direction, inasmuch as it is from east, depends upon its actual velocity, and which, in some parts of it at least, must be more or less interfered with by the flowing of the resulting wind; hence the winds from due south, or south of east, are the most inconstant and the least frequent of all the winds. But the case is very different with the winds on the contrary or western side of the compass; for as the direction of the southerly current is formed in the upper regions of the atmosphere, and consequently is not interfered with by that of the resulting wind below, and the opposition of the air which ought to form the northerly current being simply that of a pressure from north when not actually flowing towards the south, and only in some degree affected by the rotation of the earth when the wind is north-west, the conditions which are necessary to produce the westerly winds are much more capable of giving them stability and duration. When these winds, instead of meeting with their forces directed more or less obliquely to each other, meet in direct opposition with nearly equal strength, a calm or very light wind is the consequence.

12. But a rarefaction of the air being anywhere produced, the direction of the wind may be further modified by the situation of this rarefaction, with regard to the atmospheric columns adjoining it, where the rarefaction does not exist; thus as storms in high latitudes move towards east (§ 15), the returning current in some parts of the storm is deflected from west.

13. The cause of these currents being the difference of temperature of adjacent portions of air, their force will depend upon the amount of this difference; hence it is much greater in winter than in summer, when the great length of day in high latitudes lessens very greatly this difference\*; hence also as the degree to which the reduction of the atmospheric pressure can be carried by the flowing of the upper current (§ 2) depends on the force of that current, and the degree in which air can be accumulated (§ 7) likewise depends on that of

\* See Phil. Mag. S. 3. vol. xx. p. 467, "On the Oscillations of the Barometer."



the lower one, both the depressions and elevations of the barometer are the greatest in winter.

14. The forces which urge on these currents are accelerating forces; but the lower current being exposed to the friction occasioned by its flowing along the surface of the earth\*, as also to checks given by inequalities of temperature, its force is unequal to that of the upper one; hence storms are by far the most frequent and violent from south.

15. The progressive motion of a south-west wind or storm has been in part previously considered (§ 3), but requires further notice; it will be according to figure 2 of Plate V. Let the upper current descend upon a station A (the top of the page in this figure and those of § 16 and 17 being supposed the north) with more or less force; as shown in § 3 and the essay on "The Storms of Tropics," the storm moves or recedes from A, in the direction from B to A; but blowing from south, it is carried by the rotation of the earth towards east, or as if impelled in the direction CA; hence its actual path is the resultant of these motions, or that shown by the arrow at A. In this direction, therefore, the storm will arrive at the places which it visits, or as the line AD moving in a direction perpendicular to its length; hence also HA will be a section of the two currents at their place of meeting, and consequently the line or parallel of the line of the minimum atmospheric pressure (§ 3), extending in the same direction, or that shown by the arrow. The progressive motion however will not be in the same direction throughout; for let A be the place at which the current descends on the arrival of the storm, it will advance a certain distance along the surface. In the progressive motion just considered, there is a comparatively rapid motion from west, because although this motion is opposed by the air in front of it, yet it is principally on the east and by air at rest, for as the storm recedes and one portion of air descends behind the previous one, the opposition on the north is in part removed by the first descending air from that which descends after it; but when the air, as in this part of the storm, advances from south to north, this opposition is felt at every degree of its progress; hence the path taken by the wind in this case is simply the resultant of the directions of the two forces, but that from south-west being much the strongest, it is from a point much nearer to this than to that from which the opposite force is directed. Now

\* See Phil. Mag., October 1843, "On the Storms of the Tropics," p. 277. I may here correct an error in the note on that page: it stated, that omitting the two months in which the change of the monsoons occurs, the difference of the atmospheric pressure of the two seasons at Canton is nearly one-third of an inch: it ought to have been nearly half an inch, or 0.44.

agreeably with this, we find that the direction of the wind in such cases is generally at first from S.S.E., but as the storm continues, it changes to S. or S.W.; according to the explanation given of the lateral motion of the receding wind (one portion removing the opposition for that behind). Therefore let us suppose the direction of the advancing air to be from due south or along the line  $AB$ ; then  $a, b, c$  will be stations at which it arrives in its progress; but at the time it reaches any of them the storm will have moved on a certain distance from  $A$  in the direction of the arrow; therefore suppose  $A$  to be moved back along the line  $HA$ , then the wind will arrive on each point of the line  $EA$  from a point immediately south of it on the line  $HA$ ; and if the intervals of time which have elapsed on its arrival at stations equidistant with  $a, b, c$ , from the line  $HA$  in a direction due north, or parallel to  $AB$ , be severally represented by the lines  $af, be,$  and  $cd$ , parallel to the arrow, the wind will arrive at the stations  $a, b, c$ , or every point on the line  $AB$ , as  $f, e, d$ , or the corresponding points of the line  $EA$ , supposing it to move together with the line  $DA$ . Thus as fresh portions of air will advance as the point  $A$  moves forwards, the storm may be represented by a moving body of air, within which the wind is S.W., S. or S.E., and whose progressing front has the shape  $DAE$ ; and we may name that part of it represented by  $A\bar{E}$  the advancing portion\*, and that by  $DA$  the receding portion, according to the nature of the motion.

16. In storms of the tropical regions, and in those of high latitudes commencing with the usual atmospheric pressure, the former of these is apparently insignificant; but in some cases this portion of the storm extends over so great a space, and the phænomena presented by it are so peculiar, that it will require a distinct consideration. It is evident that its extent, or the distance to which the wind advances, will depend not only on its force, but also on the greater or less resistance of the air, or in other words, the greater or less pressure of the atmosphere in front of it (the resistance from the difference of temperature being supposed the same in all cases); hence we find that the cases in which it is traced in the following observations to a considerable distance, and to which this paragraph is intended principally to apply, are those in which the height of the barometer has been much reduced by

\* This must not be confounded with the second period of storms, in which the north wind advances upon the receding south. For a description of the winds in *both* the parts,  $AC$  and  $CB$ , fig. 1, of a progressing body of air, see this Magazine, vol. xxiii. p. 214; if the several directions there given be reversed, it will apply to this case. It is to tropical storms that we must look for an exhibition of these phænomena in their greatest simplicity.

a previous storm, and not restored on the arrival of the advancing storm, which approaches from atmospheric columns on the south but little affected by the preceding one; which difference indeed is apparently the cause of the storm; it occurs however after the returning or northerly current has set in in the north, and a little raised the height of the barometer there; and on its termination, the height of the barometer in the south is reduced as low as that in the north, in some parts of which it is depressed to a greater degree than before, the fall beginning in the south and advancing towards north as the wind itself. But not only this, the minimum height to which the barometer is reduced, and the setting in of the north wind (though always from north-west), occur first in the south, but on the western side (by the south being now meant the parts along the line *H A* of fig. 2, Plate V., where the upper current is supposed, with regard to localities north of them, first to descend); so that, in the south, the north wind is blowing and the barometer rising, whilst the south wind is blowing and the barometer falling in the north.

The advance of the south wind northwards after its cessation on the south, admits of very easy solution on the supposition of the storm being carried forwards by means of portions of air descending from an upper current flowing in its proper position, and which the reduction of the height of the atmospheric columns towards which it is moving, allows to flow with its velocity little checked; thus the south wind continues below, not by the force of the original impulse received at its outset from the line *H A*, but by that of successive impulses received during its course: moreover, the rapid increase of the heights of the atmospheric columns in the south by the influx of air from north-west, still maintains the upper current of the atmosphere.

The setting in of the northerly current from north-west in the south, and the consequent rising of the barometer before these changes occur in the north, may be explained by a consideration of the form of the space, on which the depression of the barometer previous to the occurrence of the second one produced by the advancing storm, exists. Suppose the greatest depression of the barometer to be produced along the line *A B*, fig. 3, Plate V. (*H A* of fig. 2), according to § 15, and the full effect of the first storm in depressing the barometer to have taken place; as that storm moved from *A* to *B*, the pressure will be in some degree restored on the parts of this line towards *A*; hence the depression will decrease from *B* to *A*; but it also decreases towards *C* (§ 2); hence the form of this space on the west will be somewhat like that bounded by

the line  $FcAb$ , the pressure increasing from  $B$  towards all parts of that line. Under these circumstances, then, the advancing storm occasioned by the greater height of the atmospheric columns about  $C$  than of those about  $B$ , occurs, and reduces the barometer at  $C$ , so as to make it the place of the greatest depression, the line  $cC$  now representing the line  $HA$  of fig. 2. But whilst the south current is thus setting in from  $C$ , the north-east is blowing from  $b$ , and the two meet somewhere between these points; and there, as at  $D$ , fig. 3, the wind is south-east (§ 11), though south-west in the localities south of it; but as the direction of the current which carries off the air so as to produce the depression is from south-west, the depression will be produced in that direction also, or along the line  $CB$ . Now if we consider the storm to have advanced from any line drawn from  $C$  towards the circumference of the depression as  $Cc$ , it is evident that whatever part of this we suppose a particular portion of the current to set out from, its condition will be the same, that of having at the time when the pressure of the atmosphere is reduced to the lowest point, a greater atmospheric pressure on the north-west side, or along lines drawn from it to the circumference  $cAb$ , as  $Ca$  and  $CA$ , because of that side being always adjacent to atmospheric columns less and previously affected by the storm; hence on the line  $Cc$ , the occurrence of the minimum in order of time will be in the direction from  $c$  to  $C$ , or that of the receding storm (§ 15). Let then the barometer be reduced to its minimum height at  $C$  (supposed the place where the storm is most intense), or to the limit of its equality with the resistance of the air in the direction from  $A$  to  $C$ , and let the air at  $d$  be reduced to the same degree of rarefaction, or to one (as afterwards to be noticed) not quite so great. These stations are subject to the influence of the pressure increasing along the line  $CA$ ; hence from any station  $E$  on that line, the air will tend to flow towards them, as indicated by the arrows at  $E$ , the effect of which in the case of  $C$  is to oppose the southerly current, but in that of  $d$  rather to assist it; hence whilst at  $C$  it is overcome and the wind sets in from north-west, it continues its course at  $d$ . The commencement of the restoration of the pressure of the atmosphere advancing thus from  $C$  towards  $D$ , whilst its advance from west is, as before stated, along the line  $cC$  or  $AC$ , it is evident that every point of the line  $CD B$  in succession from  $C$  will be related to a line parallel to  $AC$ , precisely as the point  $C$  is related to the line  $AC$  at the time of the setting in of the north wind; and the restoration of the pressure will commence as the line  $AC$  ad-

vances along the line A B, except that as it approaches B, from the great extension of the minimum depression in the north, the point A may move more slowly than the point C; near this time, however, the north wind begins to set in directly in front of the storm from north-east, and then the restoration of the atmospheric pressure proceeds throughout.

But not only does the minimum of the barometer occur first at the point C, its depression is also sometimes the greatest at this point; thus at 9 A.M. on the 11th, its height at Cork was 28.93, at Belfast 29.10, and at Plymouth (to the south-east) 29.11. Thus it appears that the barometer in this instance was about one-tenth of an inch lower at a station similar to the point C than at the point  $d^*$ . In § 2 it is stated merely that the minimum height of the barometer is near the point of meeting of the opposite currents, in order to simplify the reasoning of that paragraph; but it is evident that it will be rather to the south of it, for at this point the resistance of the air in front of the south wind at the surface of the earth, is either equal to its force, in which case it advances no further, or if not equal to it, it is yielding to it and retreating. In either case the opposition will cause more or less condensation where it is immediately felt, but the opposition decreases towards the higher regions of the atmosphere; hence (§ 2) the upper current, whose force is only exhausted by the destruction of its momentum, carries off the air from the point  $d$  (which, for the sake of illustration, let us suppose the limit of the south wind), and the positions south of it, so as still to carry on the reduction of pressure; but in the greatest degree a little south of  $d$  (or at C), on account of the condensation decreasing from  $d$  towards C. Now in storms in which the advancing portion is insignificant, the minimum pressure will be very near the point C of fig. 1; but in such as those now in consideration, it is apparently at a great distance, though the difference of the pressure at the point of the meeting of the currents, and that of the minimum pressure, is very slight. The distance, however, is in a great degree only apparent, being occasioned by the nature of the advance of the storm.

17. In high latitudes, the warmer regions, except when they

\* This appears to me a strong confirmation of the belief, that the origin of the south wind is that upon the supposition of which this theory is founded, for how upon any other than that of air descending from a current which flows by an acquired velocity, could a current flow from one station, C (fig. 3), to another,  $d$ , of a colder temperature and greater pressure? In this paragraph, as also in those which follow it, it will be observed that I have been obliged to depart from the form of the reasoning in the others, that of simple deduction from the principles stated at the outset, or from the results of previous paragraphs.

are influenced by geographical situation, have their position, with regard to colder, constantly on the equatorial side of them, both in winter and summer; but in tropical regions these positions in summer are reversed by the change in the point over which the sun is vertical; hence the currents also are either permanently reversed during this season, as in the monsoons of the Indian ocean; or, as in the "trade winds," their position is altered and their constancy interrupted, and sometimes at least their direction reversed\*; thus the tropical hurricanes of the Atlantic, which occur only during summer, and the storms of the Indian ocean, which occur during the same season, that is when the south-west monsoon is blowing, have their relative parts precisely the reverse of those of high latitudes, the descending current being from north, from which quarter the storm commences, and the returning one from south, whilst the progressive motion is towards north-west.

These storms visit only the western parts of the Atlantic, a fact which, on the supposition of their origin being the descent of the upper current, is readily explained by a glance at the position of the continents of Africa and America; the western part of the Atlantic having the latter stretching out on the north of it and radiating the heat of the summer's sun, whilst the former extends on the south of the eastern parts, having only the waters of the ocean on the north. But these storms near the boundary of the tropics change their direction† and pass along the eastern coast of North America and parts adjacent, and present phænomena different from those of storms which take their rise in extra-tropical latitudes, but to which the explanation given in the foregoing paragraph of the phænomena of the advancing portion of storms in the longitudes of Europe, may with some modification be applied. In my paper on the Storms of the Tropics, I have referred to two storms of this kind, which advanced along the coast of the United States in a direction from S.S.W. to N.N.E., of which the data collected by W. C. Redfield are given in Col. Reid's work on Storms. As a general explanation only was before given of their phænomena, it may now be proper to give one

\* For a particular explanation of the phænomena of these storms I must refer the reader to my essay in this Magazine, vol. xxiii. I may observe here, however, that the identity of the phænomena of the storms of the Atlantic and Indian ocean is sufficient evidence of the same condition of the currents in the former as in the latter at the time of the occurrence of the hurricane; but in the western part of the Atlantic, which is the locality of the hurricanes, the south is the prevalent wind in summer. (See an Essay on the Climate of Barbadoes, by Robert Lawson, in the Edinburgh Phil. Journal for July 1845.)

† On the Storms of the Tropics, Phil. Mag. vol. xxiii. p. 206.

more explicit. I have therefore quoted below the principal data of the second of these storms (Law of Storms, p. 18), and from them constructed a chart (Plate IV.), showing the direction of the wind at the different localities at the *onset* of the storm, and the time occupied by the first period of it.

“At Charleston (S. C.), on the 16th, the gale was from the S.E. and E. till 4 P.M., then N.E. and round to N.W.

“At Wilmington (N. C.) the storm was from the E., and veered subsequently to the W.

“In the vicinity of Cape Hatteras, at sea, the storm was very heavy from S.E., and shifted to N.W.

“Early on the morning of the 17th, the gale was felt severely at Norfolk, and also in Chesapeake Bay from the N.E.

“Off the Capes of Virginia, on the 17th, in lat.  $36^{\circ} 20'$ , long.  $74^{\circ} 2'$ , a ‘perfect hurricane’ from S. to S.S.E. from 5 A.M. to 2 P.M., then shifted to N.W.

“Off Chincoteague (M.d.), precise distance from the coast unknown, the gale was severe between S.S.E. and N.N.E.

“Off the coast of Delaware, in lat.  $38^{\circ}$ , long.  $72^{\circ}$ , ‘tremendous gale,’ commencing at S.E. at 1 P.M. on the 17th, and blowing six hours, then changed to N.W.

“At Cape May (N. J.) the gale was N.E. off Cape May, in lat.  $39^{\circ}$ , long.  $74^{\circ} 15'$ ; heavy gale from E.N.E. on the afternoon of the 17th of August.

“Near Egg Harbour, coast of New Jersey, the gale was heavy at N.E. on the same afternoon.

“Off the same coast, in lat.  $39^{\circ}$ , long.  $73^{\circ}$ , the gale was at E.N.E.

“In the same latitude, long.  $70^{\circ} 30'$ , ‘tremendous gale,’ commencing at S.S.E. and veering to N.

“At New York and on Long Island Sound, the gale was at N.N.E. and N.E. on the afternoon and evening of the 17th.

“Off Nantucket Shoals, at 8 p.m., the gale commenced severe at N.E. by E.

“In the Gulf-stream, off Nantucket, in lat.  $38^{\circ} 15'$ , long.  $67^{\circ} 30'$ , on the night of the 17th, ‘tremendous hurricane,’ commencing at S., and veering with increasing severity to S.W., W., and N.W.

“At Elizabeth Island, Chatham, and Cape Cod (Mass.), the gale was severe, at N.E., on the night between the 17th and 18th.

“On the 18th, heavy gale from N.E. at Salem and Newbury Port (Mass.).

“Early on the 18th, in lat.  $39^{\circ} 51'$ , long.  $69^{\circ}$ , severe gale from S.E., suddenly shifting to N.

“In lat.  $41^{\circ} 20'$ , long.  $66^{\circ} 25'$ , ‘tremendous hurricane’ from N.N.E. on the 18th.”

I cannot refer to the charts given by Col. Reid, because the directions of the wind marked out in them do not indicate the direction in any regular order with regard to the two periods of the storm.

Now the general order of the phænomena of the storms (for it is the same in both, the second being selected merely because of the information concerning it being more full than in the first one, apparently on account of its position being more westerly, and therefore including a larger portion of the United States) are as follows:—The storm commences at S.E. or N.E., but in both cases terminates at N.W., excepting in a few instances—principally near the limit of the storm in the north—where the direction of the wind at both the onset and termination of the storm is from N.E. The veering of the wind is sometimes from S.E. to N.E. and then to N.W., but more generally at once from S.E. to N.W.; and when the onset is from N.E., sometimes from N.E. to S.E., and afterwards to N.W., but more frequently directly from N.E. to N.W. The differences between the storms now under consideration, and those to which § 16 specially applies, are,—1st, that in the former the rarefaction of the atmosphere is much more suddenly produced, on account of the much greater force of the wind, and hence the extent and duration of each continuous portion of the storm is much less than in the latter; thus instead of one wind prevailing at once over a large extent of surface, as from the extreme south to the north of England, and for a long period, an American storm, over the same length of tract, consists of many alternate portions, in which the direction of the wind varies, as shown in the account given of its changes, and continues only for a few hours; and, 2ndly, in British advancing storms the collision of the north and south current takes place from the flow of the opposite currents towards a rarefaction produced by a previous storm; but in American storms the rarefaction in the first instance is occasioned by the recession of the storms from localities where the directions of the atmospheric currents are the reverse of those which are now concerned in it. Now from the distribution of the arrows in the chart, we perceive that the localities where the wind is S.E. have a constant position with regard to those where it is N.E.; and if we select any three or four positions in the track of the storm, at the time when the wind at the most southerly one has changed to N.W., the wind at the same instant of time will be blowing according to the directions shown by the ar-



rows at A B C E, Plate V. fig. 4\*. Now the upper current by which these storms begin within the tropic is from north-east, but after this change in their course the upper current is from south-west and the lower one north-east; let us then suppose A the station at which the storm first arrives after the change in the direction of the atmospheric currents, the upper one being now S.W., the rarefaction of the atmosphere is first produced here by the flowing of the air from it towards south, according to the recession of tropical storms, and into this rarefaction the south current descends, but instead of restoring the atmospheric pressure, it still further increases its diminution by its momentum, and extends northwards and eastwards to B. But it is evident that the rarefaction will extend to a much greater distance in front of the storm than on the western side of it †, hence the pressure on the line A C, fig. 3 (but now less inclined from C B), overcomes at length the force of the south wind at A where the depression of the barometer is the greatest and the wind sets in from N.W., as shown in the figure, and restores the pressure, so as to maintain by again raising the height of the atmospheric columns at A, the velocity of the upper current now flowing to more northerly localities.

But whilst this is going on, the north-east wind is blowing at C and E, being produced there by two causes, its recession from A, and the flow of air produced by the rarefaction at the limit of the south wind, as at E, fig. 1, Plate V.; and as the direction of the upper current is from south-west, the diminution of the atmospheric pressure is of course carried on in that direction; and hence a rarefaction is maintained, into which the south current flows from columns at a point eastward of A, as B, which as yet is not subject in so great a degree as A, to the opposition of the air on the line A C of fig. 3. But the south wind meeting the north, the wind is S.E. But as the line A C advances, its pressure prevails both at B and C, and

\* I would just remark in passing, how well these positions would accord with the hypothesis of the wind moving in a whirl, could the fourth quarter wanting, when the wind should be south-west, be found; but scarcely one observation of the wind from south-west occurs in the direct path of the storm, for when the wind is stated as blowing from south-west, it is either previous to the change in the progressive motion, or to the west of the "hurricane tract of the storm." The occurrence of the southerly current as a south-east wind in front of the north-east, as exhibited by the diagrams, is a grand illustration of § 11.

† This is evident from the direction of the progressive motion, but an observation given in the data of the storm of 1821 (Law of Storms, p. 16) shows how abruptly the storm terminates on the west, for "at Wilmington there was no gale," but "a severe gale was experienced thirty miles outside of the American coast off Wilmington (N. Carolina)."

changes the wind at both places to north-west, whilst the phænomena previously in existence at those stations are removed to others further north. Now it is obvious that whilst at a station C, fig. 4, where the wind is north-east, it may change directly to north-west, at another, E, more to the east, it will change first to south-east if the south wind has the greatest force; but if the north-east become the strongest, which at any particular spot may be the case, the wind in the first part of the storm from south-east may change to north-east before changing to north-west, as shown by some of the data;—a particular instance may be given.

From the data of the hurricane of 1821 (*Law of Storms*, p. 17), “At Cape Henlopen, Delaware, the hurricane commenced at  $11\frac{1}{2}$  A.M. from E.S.E.; shifted in twenty minutes to E.N.E. and blew very heavy for nearly an hour. A calm of half an hour succeeded, and the wind then shifted to the W.N.W. and blew, if possible, with still greater violence.” “At Cape May, New Jersey” (a little to the north-east of the previous locality), “commenced at N.E. at 2 P.M. and veered to S.E.” Thus it appears that at two stations situated with regard to each other as A and E, the phænomena were as follows:—At A the storm arrived at  $11\frac{1}{2}$  A.M., and blew as an E.S.E. wind, but about 12 P.M. the north current had increased in force and the wind changed to E.N.E., from which point it blew for an hour, or till 1 P.M. All this time it appears there was no storm at E., but the N.E. wind had receded to it at 2 P.M., and began to blow at that time, but the S. wind soon arrived with greater strength and the wind changed to S.E. But if this be the true explanation of the action of these storms, then according to that given of the mode of progression of receding storms, they ought to increase on the western side towards north-west by the recession of the north-east current; and on the south-west towards south-east by the recession of the south wind. Now with regard to the increase on the west side; as the direction of the north wind is opposed to that of the upper current, it is evident that by extending itself to the west, it cannot extend the rarefaction; and this being produced in a direction from south-west to north-east by the flow of the upper current, the resistance on the west side would soon overcome the advance from east: moreover, the north-east wind is not caused simply by its recession from south, but by the production or increase of the atmospheric rarefaction, as at E, fig. 1. § 2. None of these causes however operate as obstacles to its increase towards south-east, and hence we find that the storm actually does increase towards east, and that throughout its whole extent a

south wind progresses towards south-east. This is shown by the data, together with the report of the ship *Blanche*, whose log is given; for on the 17th (A.M.) she was in lat.  $31^{\circ} 42'$ , long.  $76^{\circ} 59'$ , with "fresh breezes and squally" from south by west, but at this time the hurricane was "off the Capes of Virginia in lat.  $36^{\circ} 20'$ , long.  $74^{\circ} 2'$ ;" again, "off Nantucket Shoals (lat.  $41^{\circ} 5'$ , long.  $70^{\circ}$ ) the gale commenced at 8 P.M. of the 17th," and "off Nantucket, in lat.  $38^{\circ} 15'$ , long.  $67^{\circ} 30'$ , on the night of the 17th; also early on the 18th, in lat.  $39^{\circ} 51'$ , long.  $69^{\circ}$ ."

18. It is obvious that the foregoing results, if correct, ought to enable us to explain the mode of veering of the wind, and so in great measure they will; and when they are defective, the want arises from our ignorance of the circumstances immediately contingent on the descent of the upper current. If air simply descends upon the north-east current, or meets it from a position on the south, it is evident that whether it changes towards west or towards south of east, will depend on the degree of easterly deflection the north wind has attained; hence, if the north wind be blowing briskly, the change would probably be towards south of east; and if feebly, towards west. Also, if a station upon which the north-east wind is blowing receive a south wind approaching it as the line *A D* (fig. 2), it is obvious that it could not change to north-west and then to south-west, for the air sweeping along the surface in the direction of *D A* would gradually draw the air adjacent to it into its own direction; consequently the wind would change first to south-east. Now this change does generally occur, but not always; for on the 7th and 8th of the month chosen for these observations, the wind changed from north-east to north-west, and blowing in that direction some time, afterwards changed to south-west on the arrival of a storm moving like that of fig. 2. From this therefore we may infer, that portions of the upper current were already descending when it arrived in full force as a south-west storm. The change of the wind from south to north, however, is not so much dependent on circumstances. The position of the line *A C*, fig. 3, which must always exist with more or less inclination to the direction of the storm or *D A*, fig. 2, and *C D B*, fig. 3 (its peculiar effect in the case of § 16 being occasioned by the peculiarity in the distribution of the pressure of the atmosphere on the line *C D*), determines it in this case; thus we find that in the southern and western portions of the locality of a south wind, the south wind first changes to north-west; but as the north-east wind advances from the northern verge of the storm or wind, it changes again to

north-east,—a change from south-west to south-east in this case, in the southern localities, being almost impossible whilst the wind continues to blow on the south-east side of the locality which the storm has left. Now it is matter of general observation that the wind very seldom changes in this direction, or from south-west to south-east, being indeed termed by nautical men “backing”; there is however a particular case in which the wind sometimes changes from S.W. to S.S.E. in the northern or central portions of a space occupied by a storm; and that is in the occurrence of a storm as that of § 16, the circumstances of which fully explain the exception, for the change is the consequence of the collision of the currents by which the S.S.E. wind is produced, and hence it takes place at what for the moment is on the northern verge of the south wind, where of course a change produced by a north-east wind meeting it may make it south-east\*.

Having carried out thus far the results of the principles stated at the beginning of this paper, I may now proceed to give the observations which I have collected, and first those of the wind, extracted from the Shipping Gazette.

*Scotland.—Orkney, Longhope.*—“November 1. N., moderate. 2. S.E., fresh breeze. 5. N.W., moderate. 8. S.W., blowing hard; rain. 11. E., strong breeze; rain. 13. N.E., squally. 15. N.N.E., fresh. 16. N.E., frosty. 19. E., strong breeze; rain. 20, 21. N. to N.E., frost and sudden squalls. 22. E. to S.E. 23. E., fresh breeze.”

*Pentland Frith.*—“Nov. 7. N.W., moderate: night, S.W.; very strong throughout the night. 8. S.W. 18. W., moderate. 19. S.E., moderate; rain: 6 P.M., N.E., moderate.”

*Thurzo.*—“Nov. 2. S.E., moderate. 8. S.W., fresh breeze. 10. N.W., heavy gale. 11. N.E. 17. S.W. 18. N.E. 19. S.E., moderate weather. 23. S.E.”

*Peterhead.*—“Nov. 1. N.W., light. 3. S.E. to E.S.E., light breezes. 4. E., moderate. 5. N.E., fresh breeze. 8. S.W., fresh breeze. 9. S.W., strong gale. 19. S.E. to N.E., rainy. 20. N.E., strong breeze. 23. E.N.E., fresh breeze. 24. E., strong.”

*Inverness.*—“Nov. 19. N.E., calm and raining. 26. N.E., calm; rain.”

*Aberdeen.*—“Nov. 17. E.N.E. 18. Variable, E.N.E.”

*Mull—Tobermorey.*—“Nov. 3. S.E. 4. S.E., moderate breeze. 5. S.E. to E., light breeze. 7. Variable; light airs and heavy showers of rain. 8. S.W., strong gales. 9. S.W., fresh breeze, with rain at intervals. 10. E.N.E., fresh breeze. 11. E., strong breeze. 12. W. to S., light airs; variable. 13. E.N.E., fresh breeze.

\* In these paragraphs I have omitted any mention of the differences of the mean pressure on different latitudes of the surface of the earth, not because of its unimportance, but because it would merely be a transcript of my essay on that subject in this Magazine, vol. xx. p. 469.

14. E.N.E., moderate breeze. 15. E.N.E., fresh breeze. 16. E. to S.E., fresh and squally. 17. S.S.W., moderate breeze. 22. S.S.E., strong gales; heavy rain. 23. S., moderate breezes. 24. N.N.E., strong breezes."

*Islay*.—*Bowmore*.—" Nov. 2. N.E. 10. S.W., blowing hard. 15. N.E. 16. S.E., a gale. 21. S.S.E. 23. N.W. 25. N.E., fresh."

*Bute*.—*Rothsay*.—" Nov. 1. W., blowing strong. 9. S.W., blowing strong. 20. S.W., blowing strong."

*Greenock*.—" Nov. 1. W., fine. 7. S.W., fine. 11. N.N.W. and S.E., moderate. 12. S., light airs; calm. 14. W., moderate. 18. S.E., moderate. 19. S., light airs; rain. 22. S.E., snow and sleet. 23. Variable; light airs and calm. 24. E., fresh breezes with showers."

*Glasgow*.—" Nov. 3. E., moderate. 6. N.E., fine. 8. 6 P.M. wind veered round to S.W., and blowing a gale outside. 10. Shortly after we had posted our letters (on the 8th) it began to blow a heavy gale, which towards night became a hurricane with rain, which continued all day (the 9th). Today (10th) wind N.E., fair. 14. 6-30' P.M., wind since Saturday last (12th) chiefly from N. to N.E., light. 17. N.E., light."

*Leith Roads*.—" Nov. 14. N.N.E., fine. 17. Variable; fair."

*Ireland*.—*Donegal*.—" On the night of the 8th, about 8 o'clock, there was a heavy gale of wind from W. to S.W. 26. W.N.W."

*Strangford*.—" Nov. 1. W. 2. E.S.E., fine. 3 and 4. E.S.E., fresh. 5. E. by N., fine. 14, 15, 16. E., strong breeze. 17. E.S.E., strong. 18. W.S.W., heavy breeze with rain. 19. W. 21. E. 22. W. by N. 23. W.S.W. 25. E.S.E., strong; rain. 26. N.W., rain."

*Arklow*.—" Nov. 6. E.N.E., fresh breezes. 9. A.M. S.W., strong gales with rain: P.M. strong gales. 10. At day-break, E., strong gales with rain: P.M. S.S.W., blowing hard. 11. A.M. S.W., strong gales. 12. A.M. S.W., strong gales and rain: P.M. W., fresh breeze. 13. A.M. W.S.W., fresh gales with rain: P.M. S.W., moderate. 14. A.M. N.E., strong gales: P.M. E.N.E. to E.S.E., a gale after sunset with heavy incessant rain. 15. E. by S., a heavy gale with torrents of rain. 16. E., heavy gale. 17. E., strong gales: P.M. E.N.E., fresh gales. 18. A.M. S.W., strong gales: P.M. S.W., strong gales. 19. S.W., strong gales. 20. A.M. E.N.E., fresh gales: P.M. S., hard gales and rain. 21. E.N.E., A.M. moderate: P.M. freshened to a gale. 22. W.N.W., fresh gale. 23. A.M. W., moderate: P.M. S., squally. 24. W.S.W., fresh gales: P.M. W., more moderate. 25. A.M. W., moderate: P.M. W.N.W., moderate. 26. W. by N., moderate."

*Waterford*.—" Nov. 2. W.S.W., S.E., and S.S.E. 3. S.E. to S.S.E. 4. E. to S.E., E., and N.E. 14. W. by N. to W.S.W., S.S.W., and W."

*Youghall*.—" Nov. 11. N.E., light breeze; hazy with rain. 12. W.S.W., fresh breeze; rain. 20. E. by S., fresh breeze, with rain."

*Galway*.—" Nov. 3. E.S.E. 5. E., fine. 6. E.S.E., moderate. 21.

Hard gales from E.N.E. for some days past. 22. Much rain last night; wind strong from E.N.E.; today more moderate; wind W.S.W. 26. N.N.W., light airs."

*Limerick*.—"Nov. 10. S.S.E., moderate. 11. S., moderate. 14. S.S.E., moderate. 15. S.S.E., blowing hard, with rain."

*Cove of Cork*.—"Nov. 2. Strong gales with rain. 3. S.E., stormy, with rain. 4. S.E., strong breezes. 5. N.E., a gale. 6. N.N.E., fresh breeze. 7. N.N.W., moderate. 9. S.W., stormy, with rain. 10. S.E., moderate; rain. 11. S.S.W., showery. 12. W.N.W., clear. 13. S.S.W. to W.N.W., variable; squally; heavy showers. 17. E., strong breeze. 18. S.W., stormy, with rain. 19. S.W., moderate. 20. E., strong breeze; rain. 21. S.E., moderate. 22. N.N.W., strong breeze; fair. 23. S.E., strong breeze. 24. W.S.W., fresh; heavy showers. 25. N.N.W. 26. N.W., stormy and showers."

*England, West Coast.—Holyhead*.—"Nov. 1. N. to N.E., moderate. 2. Variable and fine. 4 and 5. E. to E.N.E., strong; squally. 6. E. by N., fresh breeze. 7. N.E., fresh breeze. 8. W.S.W., fine breeze. 9. S.W., strong gale; hazy; wet. 10. A.M. E.S.E., fine breeze; rainy: P.M. S. to S.S.W., strong breeze and squally. 11. S., moderate. 12. W.N.W., strong breeze. 13. Variable; moderate; rain. 14. E., fine breeze. 15. E., strong gale; showery: 9 P.M. continued. 16. E., blowing excessively hard. 17. S.E., fine breeze: 8 P.M. S., moderate. 18. S.W., fresh breeze. 19. W.S.W., strong; rainy; night variable. 20. E.S.E., fresh breeze: 8 P.M. E., fresh. 21. E., moderate. 22. N.W., strong gale. 23. Last night N.W., strong gale; today veered to W.S.W., fine breeze: 8 P.M. S.S.E., fine breeze. 24. Variable from S.E. to S.S.W., fresh breeze; showery. 25. W.S.W. to S.W., fresh; showery. 26. W.S.W. to W., fresh; rain."

*Baumaris*.—"Nov. 11. S.W., much rain. 12. S.W., fresh; rain. 13. S.W., much rain. 14. E., fresh. 17. E., fine. 18. S.S.W., fresh; rain. 19. S.S.W., fresh; rain. 20. S.S.E. 22. W., fine. 23. S.E., fresh. 25. S.W."

*Bristol*.—"Nov. 4. E., strong. 8. E., moderate. 10. 4 P.M. S.S.W., very strong. 11. S.W., showery. 13. It has continued squally from S.W. to W. since my last, and a great quantity of rain has fallen. 14. Variable; moderate. 14. (second report) E., strong. 16. E., strong; constant rain. 19. S.W., a gale. 20. E., fresh. 21. N.E., fresh. 22. The wind this morning blew fresh from S.S.E. until 10 o'clock, when it gradually veered round to N.N.W., and afterwards it was very strong from that point; a quantity of rain fell last night. 23. N.W., fresh. 24. W.S.W., strong; showery."

*Scilly Islands.—St. Mary's*.—"Nov. 1. S.E., fresh breeze. 2. S.E., strong wind and rain. 3. S.E., strong gales, with rain. 4. E.S.E., strong gales; rain. 5 and 6. E., strong breeze. 7. E.N.E., strong, showers. 8. E., fresh breeze. 9 and 10. S.S.W., strong gales; rain. 11. W., strong; rain. 12. W. 13. W.S.W. 14 and 15. S.W. 16. S.W. to S.E., strong, with rain for the

last four days. 17. E., fresh breeze. 19. S.W., strong; rain. 20. W.N.W., rain. 21. E.S.E., strong; rain. 25. W.N.W., strong gales and rain. 26. W.S.W., fresh breezes and rain."

*South Coast.—Falmouth.*—"Nov. 1. N.E., light breeze. 2. E.S.E. to S.E. 3. E. by S. 4 and 5. E.N.E., strong. 6. E.N.E. 7. N.E., light. 8. N., light airs. 9. S. 10. S., a gale; rain throughout. 13. 8 A.M. S., a gale; raining; in the evening wind changed to W. 14. Weather moderated. 15. S., squally; rain; afterwards W. 16. Morning S.W. and raining; evening N.E., strong gale. 17. 1 A.M. N.E., moderate; afterwards E. 18. S. 19. S.S.W., blowing hard. 20. W., light airs and rain: 10 P.M. veered to S.E. 21. E., showery. 22. N.N.W. to N.W., showery. 23. S.W., showery. 24. W. by S., heavy gale, with rain. 26. N.W., showers."

*Plymouth.*—"Nov. 4. N.E., moderate. 5. N.E., fresh breeze. 6. N.N.E. to E.N.E. 8. N.E., light airs. 9. S.S.W., strong, with rain. 10. S.W., strong; thick rain. 11. S.S.W., strong; rain. 12. W. by N. 13. It has blown very heavy through the night from S.S.W. with rain, which (at noon) still continues. 14. Calm; hazy. 15. W.S.W., strong. 18. S.E., fresh breeze. 19. S.S.W., dirty. 21. Easterly, moderate. 22. N.W., fresh breeze. 23. W.N.W., fresh breeze. 24. W.S.W., strong; squally. 25. W.S.W., strong breeze, with showers. 26. W., moderate."

*Penzance.*—"Nov. 26. W.N.W., strong gales."

*Portsmouth.*—"Nov. 3. E.S.E. 4. E., fresh breezes. 6. N. to N.N.E., foggy; slight showers of rain. 9. S.W., blowing fresh. 11. S.W., rain. 14. W. by S.: P.M. fresh breeze. 15. S.W., light breeze; rain all day. 17. N. by E., fresh breeze. 18. S. by E., fresh breezes. 21. N.E., hazy. 23. P.M. W.S.W., fresh breezes; rain. 24. W.S.W., rain. 25. W.S.W., rain and squalls."

*Isle of Wight.—Ryde.*—"Nov. 12. W.S.W. to W. 13. S.W. to W.S.W., with rain; blowing hard. 14. W.N.W., fine. 15. S.W. 16. E.S.E., fresh. 17. N.E. to E.N.E., fresh. 18. S., fine. 21. E., fine. 24. W.S.W., strong, with rain."

*Deal.*—"Nov. 1. W.N.W. and N. by E., moderate and fine. 2. E.N.E. to E.S.E. 3. N.E., moderate and fine. 4. S.E., light airs and rain: P.M. N.E., fresh and squally. 5 and 6. E.N.E., blowing fresh and squally. 7. E.N.E., fresh breeze; squally. 10. S.W., fresh breeze; squally. 11. S.W., blowing very strong, with rain. 12. It has blown very hard all day from W.S.W., with squalls of rain. 14. N.N.W., light airs. 16. E., fresh. 17. A.M. E., blowing hard: 6-30' P.M. E.S.E. 18. S.W., moderate. 20. N.N.W. and N.E., moderate. 21. N.E. 22. A.M. S.S.W., blowing strong; rain: P.M. S.E., moderate; rain. 23. W.S.W. and W.N.W., light airs. 24. W.S.W., blowing fresh. 25. It blew a gale of wind last night and nearly the whole of this day from S.W. 26. It blew very strong during last night from the S.W. or W.S.W., with squalls of rain."

*Dover.*—“ Nov. 3. 7 A.M. S.S.W., light: noon and 7 P.M. S.E., fresh. 4. N.E., light, with rain. 5. E.N.E., strong; cloudy. 6. A.M. E., fresh: P.M. E.S.E., fresh; rain. 8. N.N.W., light. 9. S.W., fresh: 7 P.M. W.S.W., strong. 10. 7 A.M. W.S.W., fresh: noon, S.S.W., strong: 7 P.M. S.W., fresh. 15. E.S.E., strong. 16. 7 A.M. E., light wind; rain: noon and 7 P.M. E., strong. 17. E., strong. 21. N.E., fresh. 24. 7 A.M. S.S.W., strong, with rain: 7 P.M. W., strong; rain. 25. S.W., strong.”

*East Coast.—The Downs.*—“ Nov. 1. N.W., light breeze. 2. Easterly, moderate. 8. N.W., very light. 9. S.W., blowing fresh. 10. S.W., blowing fresh. 14. W.N.W., light. 15. E. to S.E., fresh. 16. E.N.E., moderate breezes. 18. S.W., very light. 19. S., blowing fresh. 20. N., moderate. 21. N.E., light breezes. 23. W., light. 25. W.S.W., blowing very fresh. 26. W.S.W., blowing fresh.”

*North Foreland.*—“ Nov. 1. N.W. 2. E. to S.E., fresh. 3. Variable. 4. E., squally. 6. Blowing fresh. 12. S.W.: noon, W.N.W.: P.M. W., squally. 14. N.W. by W. to N., light. 15. E. to E.S.E., blowing strong. 16. E.S.E., blowing fresh. 17. E.S.E., fresh. 18. S.E. to S.S.W., light. 20. N. to E., moderate. 21. N.E. 22. S.E., moderate. 23. W.S.W., light. 24. S. to S.W., blowing hard; rain. 25. S.W., squally, with rain. 26. S.S.W. to S., blowing strong.”

*Yarmouth (Norfolk).*—“ Nov. 1. N.W., fine. 2. N.N.E. 9. S.S.W., blowing fresh. 10. S.S.W. to S.W., fresh breeze. 11. S.S.W. to S.W., blowing strong, with rain. 12. 6 P.M. N.W., strong. 14. N. to W.N.W., fine. 15. S.S.E. 16. Last night and all this day the wind has blown fresh from E.N.E. to N.E. 17. E. 25. S. to S.S.W., blowing strong. 26. S.S.W., blowing fresh.”

*Lowestoft.*—“ Nov. 1. W. 2. E.N.E., light breezes. 18. N.W., light. 21. N.N.E., light breezes.”

*Flamborough Head.*—“ Nov. 1. W. to N.W., light breezes. 2. E.S.E., light breezes. 3. E. to E.S.E., light breeze. 4. E.N.E., strong breezes. 5. E.N.E., strong breezes and squally. 6. E.N.E., strong breezes. 7. N.E., strong breezes and squally. 8. W.N.W., light breeze. 9. S.W., strong gales. 10. S.W., light breezes. 11. S. by E., strong breezes, with rain. 12. S.W. to W., fresh breezes. 13. S.W., light breezes: evening, E., light. 14. N. to N.E., moderate breezes. 15. E., moderate breezes. 16. E., strong breezes. 17. E.N.E., blowing fresh. 18. W., light breezes. 20. N.N.E., strong breezes. 21. N.E., strong breeze. 22. S.S.W. to S., strong breeze. 23. W., moderate breeze. 24. S.E., strong gale; rain. 25. S., strong breeze. 26. S.S.W. to S.W., strong breezes.”

*North Shields.*—“ Nov. 1. N.W., brisk in the morning. 2. S.S.W. morning and evening; light and S.S.E. in the middle of the day; brisk. 3. S.E. and E.S.E., moderate: evening, E. 4. E.N.E., strong. 5. N.E., strong, but moderate towards evening. 6. E.N.E., moderate. 7. 9 A.M. N.N.E., light: 9 P.M. W.N.W., fresh; light rain. 8. 9 A.M. W.N.W., light: 2 P.M. W.S.W.,



light: 9 P.M. S.W., almost calm. 9. Storm from S.W., which began early in the morning and continued till 5 P.M., at which time the wind had sunk down to a calm with rain; the barometer being then at its minimum, 29.276. The calm continued through the night, with a hoar frost. 10. 9 A.M. W.N.W., very light; sky clear, but soon afterwards overcast: 2 P.M., wind extremely light and variable; rain falling: 9 P.M. wind strong from S.E., which continued, with heavy rain, all night; and until afternoon of the 11th, on which day the air became again calm in the evening. 12. 9 A.M. W.N.W., very light wind: 9 P.M. strong from N.W. 13. 9 A.M. and 2 P.M. wind extremely light from W.S.W. and S.W.; in the evening strong from N.E., with rain. 14. N.E., brisk. 15. N.E., brisk, with showers during the day, and strong at night. 16. N.E. and E.N.E., strong. 17. 9 A.M. E., very light; calm during the remainder of the day. 18. W.S.W., very light during most of the day, but strong at night. 19. Morning, W.S.W., brisk; evening, light from W.N.W., slight rain. 20. N.E., rather brisk; showers in the evening and during the night. 21. N. to N.W., rather brisk in the middle of the day; showers. 22. Morning, strong from S.S.W.; changed to S.W. in the afternoon, and sunk down to a calm in the evening; rain and snow. 23. N.W., rather brisk. 24. A storm, with rain from S.S.E., which abated towards evening. 25. S.S.E., strong till evening, when the air became calm; fine showers. 26. S.S.W. and S.W., light; rain in the evening."

The Tables which follow contain the indications of the barometer from the 1st to the 26th; each day is divided into three columns; the second and third contain the observations for morning and evening, and the middle one those of the middle of the day. The hours of observation are given after the names of the places, which are placed in the order from north to south, but the western before the eastern. In a column previous to those containing the daily indications are given the mean heights of the barometer for each place, in order that the daily heights may be compared with each other\*, in which comparison the difference of pressure due to the latitude ought to be borne in mind. With the exception of the observations at Paris, Christiania, and North Shields, which are reduced to the temperature of 32° Fahr., the numbers are those which are read off from the barometer; as,

\* It is not meant by this to be understood that these means are given as representing by their differences the differences of height of the several barometers as read off from each scale, supposing them placed in juxtaposition, but only to serve in the following observations for standards for the comparison of their variations with each other in the absence of any other method of doing it; but the unusual equality of the mean pressure of the atmosphere of this month over so large a space, renders the means chosen sufficiently accurate for this purpose.

however, one degree of Fahrenheit affects the barometer at 30 inches only .003 inch, and there is seldom a greater difference than 3° between the temperatures of consecutive observations, it will not affect any of the conclusions which may be deducible from them. The numbers representing the heights at Christiania in Paris lines, and at Paris in millimetres, have been reduced to English inches.

Names of places.	Hours of observation.	Mean of the month.	1.			2.		
Orkneys ...	9½ a.m. & 8½ p.m.	29·70	30·10		30·22	30·24		30·22
Glasgow*...	9 a.m. & p.m.	29·53	30·07		30·08	30·07		30·01
Belfast.....	9 a.m. & 3 p.m.	29·74	30·35	30·33		30·24	30·18	
Armagh†...	10 a.m. & p.m.	29·44			29·97	29·91		29·82
Shields.....	9 a.m. & 2 & 9½ p.m.	29·68	30·14	30·17	30·21	30·21	30·19	30·16
Cork.....	9 a.m. & 3 p.m.	29·74	30·22	30·15		29·98	29·92	
Bristol.....	9½ a.m. & p.m.	29·70	30·22		30·14	30·09		30·01
Plymouth...	9 a.m. & p.m.	29·71	30·27		30·16	30·09		30·00
London ...	9 a.m. & 3 p.m.	29·72	30·23	30·17		30·14	30·10	
Paris .....	9 a.m. & 3 & 9 p.m.	29·61	30·08		30·00	29·90		29·83
Christiania.	9 a.m. & 10 p.m.	29·71	29·61		29·96	30·03		30·20

	3.			4.			5.		
Orkneys ...	30·20		30·28	30·44		30·54	30·53		30·51
Glasgow ...	30·00		30·08	30·21		30·34	30·37		30·28
Belfast.....	30·16	30·18		30·38	30·45		30·57	30·55	
Armagh ...	29·86		29·97	30·12		30·23	30·30		30·23
Shields.....	30·10	30·10	30·15	30·25	30·34	30·40	30·39	30·33	30·34
Cork.....	29·83	29·83		30·16	30·16		30·33	30·32	
Bristol.....	29·94		29·98	30·07		30·21	30·23		30·16
Plymouth...	29·88		29·95	30·09		30·21	30·26		30·19
London ...	30·00	29·96		30·06	30·13		30·19	30·12	
Paris .....	29·76		29·69	29·75		29·86	29·75		29·75
Christiania.	30·27		30·36	30·42		30·32	30·28		30·27

\* These observations give a mean much below the general one, which is probably the defect of the scale. I have compared them, however, with those given in the tables of the Philosophical Magazine for Applegarth Manse in Dumfries-shire, a locality not far distant, with which their variations agree, with the exception of being a little in advance in movements coming from north, for which reason I have given them the preference here. I may add also, that the observations at Belfast were received with the information that they might be rather defective, which however is of little consequence, as those at Armagh are given. I have inserted the former, because of those at 3 P.M.

† Height of the barometer 211 feet above the level of the sea.

	6.			7.			8.		
Orkneys ...	30.49		30.43	30.30		30.00	29.83		29.42
Glasgow ...	30.24		30.27	30.23		30.07	29.87		29.58
Belfast .....	30.51			30.48	30.43		30.19	30.04	
Armagh ...	30.26		30.24	30.20		30.08	29.88		29.39
Shields.....	30.34	30.35	30.33	30.30	30.27	30.23	30.05	29.97	29.77
Cork.....	30.30	30.30		30.32	30.33		30.13	30.01	
Bristol .....				30.18		30.18	30.14		29.99
Plymouth...	30.23		30.24	30.24		30.24	30.21		30.05
London ...	30.20	30.16		30.13	30.13		30.15	30.08	
Paris .....	29.85	29.83	29.87	29.84		29.94	29.93		29.88
Christiania.	30.27		30.16	30.09		29.95	29.75		29.58

	9.			10.			11.		
Orkneys ...	28.70		28.90	29.29		29.50	29.37		29.12
Glasgow ...	29.04		29.21	29.45		29.33	28.95		28.77
Belfast .....	29.36	29.41		29.65	29.57		29.10	29.02	
Armagh ...	29.11		29.30	29.34		28.97	28.74		28.68
Shields.....	29.37	29.28	29.37	29.62	29.58	29.43	29.14	28.99	28.89
Cork.....	29.42	29.42		29.34	29.20		28.93	28.91	
Bristol .....	29.66		29.55	29.57		29.35	29.05		29.02
Plymouth...	29.78		29.66	29.61		29.34	29.11		29.13
London ...	29.81	29.70		29.68	29.64		29.18	29.00	
Paris .....	29.82	29.76	29.78	29.73		29.53	29.33		29.17
Christiania.	29.46		29.28	29.14		29.34	29.45	29.48	29.45

	12.			13.			14.		
Orkneys ...	29.05		29.14	29.26		29.44	29.67		29.86
Glasgow ...	28.74		29.01	29.08		29.13	29.65		29.72
Belfast .....	29.12	29.21		29.32	29.27		29.83	29.91	
Armagh ...	28.87		29.01	29.01		29.21	29.61		29.56
Shields.....	28.94		29.20	29.27	29.24	29.25	29.71	29.82	29.84
Cork.....	29.30	29.31		29.10	29.40		29.66	29.60	
Bristol .....	29.15		29.48				29.70		29.60
Plymouth...	29.30		29.63	29.44		29.49	29.75		29.61
London ...	29.12	29.33		29.51	29.26		29.72	29.80	
Paris .....	29.30		29.57	29.66		29.39	29.65	29.67	29.63
Christiania.	29.30		29.09	28.96	28.94	28.97	29.16		29.54

	15.			16.			17.		
Orkneys ...	29.97		30.05	30.15		30.30	30.38		30.32
Glasgow ...	29.73		29.71	29.85		30.08	30.28		30.30
Belfast .....	29.85	29.82		29.96	30.06		30.47	30.51	
Armagh ...	29.53		29.48	29.66		29.97	30.20		30.25
Shields.....	29.84	29.83	29.86	29.98	30.03	30.15	30.40		30.51
Cork.....	29.40	29.37		29.49	29.70		30.30	30.31	
Bristol .....	29.60		29.61	29.68		29.88	30.25		30.46
Plymouth...	29.61		29.67	29.63		29.78	30.25		30.47
London ...	29.69	29.62		29.76	29.79		30.23	30.36	
Paris .....	29.57	29.55	29.58	29.53	29.50	29.57	29.79		30.19
Christiania.	29.69		29.72	29.80		29.92	29.98	29.94	29.95

	18.			19.			20.		
Orkneys ...	30-16		30-21	29-90		29-93	29-97		29-95
Glasgow ...	30-25		30-11	29-68		29-66	29-78		29-80
Belfast .....	30-48	30-37		29-93	29-86		29-94	29-91	
Armagh ...	30-18		29-82	29-68		29-64	29-64		29-64
Shields.....	30-46	30-42	30-34	29-88	29-77	29-73	29-85	29-85	29-86
Cork.....	30-29	30-18		29-98	29-92		29-76	29-80	
Bristol .....	30-49		30-35	30-13		29-82			
Plymouth...	30-51		30-43	30-28		30-01	29-83		29-63
London ...	30-58	30-53		30-23	30-06		29-79	29-77	
Paris .....	30-38	30-38	30-41	30-30		30-04	29-60		29-50
Christiania .	30-08		30-14	30-00		29-82	29-64		29-56

	21.			22.			23.		
Orkneys ...	29-95		29-77	29-47		29-39	29-36		29-30
Glasgow ...	29-82		29-62	29-22		29-12	29-17		29-01
Belfast .....	30-01	29-95		29-42	29-42		29-46	29-32	
Armagh ...	29-74		29-43	29-21		29-21	29-19		28-70
Shields.....	29-90	29-89	29-78	29-42	29-30	29-27	29-31	29-27	29-19
Cork.....	29-84	29-83		29-58	29-58		29-34	29-10	
Bristol .....	29-83		29-76	29-30		29-49	29-48		29-04
Plymouth...	29-78		29-79	29-45		29-62	29-59		29-02
London ...	29-80	29-82		29-42	29-28		29-50	29-59	
Paris .....	29-51		29-63	29-31	29-13	29-30	29-50		29-33
Christiania .	29-56	29-54	29-56	29-58	29-55	29-56	29-61		29-63.

	24.			25.			26.		
Orkneys ...	29-11		29-08	29-04		29-10	29-10		29-10
Glasgow ...	28-71		28-58	28-63		28-70	28-76		28-90
Belfast .....	28-83	28-79		28-74	28-82		28-97	29-04	
Armagh ...	28-47		28-40	28-47		28-61	28-73		28-88
Shields.....	28-89	28-78	28-75	28-80	28-82	28-83	28-91		29-07
Cork.....	28-54	28-54		28-79	28-80		29-00	29-04	
Bristol .....	28-83		28-75	28-75		28-94	29-04		29-24
Plymouth...	28-95		28-87	28-83		29-03	29-13		29-28
London ...	28-91	28-92		29-90	28-88		29-09	29-17	
Paris .....	28-98		29-07	28-95		29-08	29-07		29-28
Christiania .	29-65		29-67	29-66		29-58	29-58	29-57	29-60

In the columns which follow, interposed with the text, I have given the amount of the variations in 100ths of an inch between the consecutive observations given in the preceding tables, with the exception of those in which there are three daily observations; for the 3rd column always contains the difference of the extreme observations of each day, so that in casting the eye down the columns the variations may always belong to the same periods, excepting when the time of the observations varies a little from 9 o'clock; care however must be taken with regard to the first column to observe those which include a period beginning at 3 P.M. on the day previous which is the case when there is no evening observation.

For the sake of simplifying the diagrams as much as pos-

sible, I have in the first place given a chart with the names and localities marked upon it (Plate IV.), and the subsequent ones merely contain the localities marked by dots, and set off in their several places from the chart, so that by reference to it they may easily be found. The arrow representing the direction of the wind at Christiania, placed in the north-east corner, is of course out of its relative position. I have endeavoured to represent the force of the wind, as given in nautical language, by figures placed at the feet of the arrows:—0 being a calm, 1 and 2 light airs or winds, 3 moderate, 4 brisk breeze, 5 strong breeze, 6 a gale or stormy, 7 hard gale; *v* is variable: the veering of the wind backwards and forwards between points is represented by two arrows from these points, and a change of the wind in the latter part of the day, or night, by a line crossing the foot of the arrow showing its direction. A change in the force also is marked in the same way, by a line underneath the figure. The direction of the wind at Paris and Christiania is taken from the meteorological registers of those places; and I have also occasionally introduced arrows showing its direction as noted in the registers of other places, but not without caution, as I am more disposed to rely on the mercantile reports, than on observations simply made by noting the direction of one particular vane generally only once in the day. It may also be remarked, that though the variations given in the first column of each day are those which have taken place during the previous night, they may yet be generally considered as caused by the wind indicated by the diagrams for that day, because the wind in the morning is usually that of the preceding night, and when a change occurs during the day it is marked as before stated\*.

Names of Places.	1.		2.			3.		
Orkneys .....		+·12	+·02		—·02	—·02		+·08
Glasgow .....		+·01	—·01		—·06	—·01		+·08
Belfast .....	—·02		—·09	—·06		—·02	+·02	
Armagh .....			—·06		—·09	+·04		+·11
Shields.....	+·03	+·07	·00	—·02	—·05	—·06	·00	+·05
Cork.....	—·07		—·17	—·06		—·09	·00	
Bristol .....		—·08	—·05		—·08	—·07		+·04
Plymouth .....		—·11	—·07		—·09	—·12		+·07
London .....	—·06		—·03	—·04		—·10	—·04	
Paris .....		—·08	—·10		—·07	—·07		—·07
Christiania .....		+·35	+·07		+·17	+·07		+·09

\* The shipping reports however very frequently do not state the time of the day the report refers to, when it is extremely probable the wind did not continue in the same direction the whole of the day; so as to occasion in some cases apparent discrepancies in the direction of the wind; but they may generally be removed by reference to some other report from a neighbouring locality where a change in the direction of the wind is noted.

Names of Places.	4.			5.			6.		
Orkneys .....	+·16		+·10	-·01		-·02	-·02		-·06
Glasgow .....	+·13		+·13	+·03		-·09			
Belfast .....	+·20	+·07		+·12	-·02		-·04		
Armagh .....	+·15		+·11	+·07		-·07	+·06		-·02
Shields .....	+·10	+·09	+·15	-·01	-·06	-·05	+·00	+·02	+·00
Cork .....	+·33	+·00		+·17	-·01		-·02	+·00	
Bristol .....	+·09		+·14	+·02		-·07			
Plymouth .....	+·14		+·12	+·05		-·07	+·04		+·01
London .....	+·10	+·07		+·06	-·07		+·08	-·04	
Paris .....	+·06		+·11	-·11		+·00	+·10	-·02	+·02
Christiania .....	+·06		-·10	-·04		-·01	+·00		-·11

The first very conspicuous atmospheric phenomenon is an elevation of the barometer, which at Orkney reached the highest point at the P.M. observation of the 4th, and advanced towards south. Its maximum was in the north and north-west, affording an excellent illustration of § 7; for on the 4th, the day on which the principal rise took place, we find the collision of the currents on the western and south-western side of the chart: thus at the Irish ports and the Scilly islands the wind was south-east, whilst in England and Scotland and at Paris it was north-east. Hence at the Orkneys the barometer stood at 30·54 and at Belfast 30·57, these places being situated in the line of the meeting currents; whilst at London, which, so far as the observations go, was in the line of the north-east current moving freely, the barometer reached an elevation of only 30·19; the advance of the north wind and of the elevation of the barometer likewise accorded with each other, both being from north to south.

The observations of the 1st are particularly illustrative of § 5 and § 8; in them the opposition of the southern current is seen only in the extreme south, where it is blowing with some strength, causing a fall of the barometer in the south, where the north wind is blowing immediately in front of the south wind (§ 5), and a slight rise in the north by the arrival of the air removed from the south (§ 8). It might be thought that the setting out of the currents in somewhat different directions from the western part of England might give rise to this fall, but it is evidently not so, because it decreases towards the locality where the current takes the easterly deflection. The deflection of the southern current is also worthy of remark. At the Scilly islands the south wind meeting a north-east blows from the south-east, but at Paris meeting a north-west it is south-west or west-south-west (§ 11). The cause of the north wind being north-west in the north and eastern parts, appears to be owing to a deficit of pressure in the east (the barometer at Christiania being ·39 inch below that at the

Orkneys), which on this day was rapidly removed by the air flowing towards it, for at Christiania the height of the barometer in the morning was 29.61 and in the evening 29.96. The strength of the south wind after this increases, and on the 2nd and 3rd prevailed almost throughout, occasioning a slight general fall of the barometer until 9 A.M. on the 3rd; we still however have evidence of the north-east wind blowing on the east, as on the 2nd, on the south-east coast of England, and on the 3rd in the same part, as at the North Foreland, where it was variable (the one and the other of the currents alternately prevailing), and at Deal, where it was north-east, these being almost the most easterly parts of England. After this, however, the north-east wind advanced, becoming general on the eastern side on the 4th, and raising the barometer as before described, the south wind only appearing on the west, blowing there, according to § 11, from south-east. On the 5th and 6th the north-east prevailed throughout, and the barometer was slightly depressed in all the northern stations, beginning at the Orkneys; the cause of which might be, either the subsidence of an elevation above adjacent localities on the south, by the flowing of the air towards them, or it might be the beginning of the descent of the upper current on the north, which manifested itself at the Orkneys on the 6th by changing the wind to north-west.

Names of Places.	7.		8.			9.		
Orkneys .....	-13		-30	-17		-41	-72	+20
Glasgow .....	-13		-16	-20		-29	-54	+17
Belfast .....		-05		-24	-15		-68	+05
Armagh .....	-04		-12	-20		-48	-28	+19
Shields .....	-03	-03	-07	-18	-08	-28	-40	-09
Cork .....	+02	+01		-20	-12		-59	00
Bristol .....			00	-04		-15	-33	-11
Plymouth .....	00		00	-03		-16	-27	-12
London .....	-03	00	+02	-07		-27	-11	
Paris .....	-03		+10	-01		-05	-06	-04
Christiania .....	-18		-14	-20		-17	-12	-18

Names of Places.	10.		11.			
Orkneys .....	+39		+21	-13		-25
Glasgow .....	+24		-12	-38		-18
Belfast .....	+24	-08		-47	-08	
Armagh .....	+04		-37	-23		-06
Shields .....	+25	-03	-19	-29	-15	-25
Cork .....	-08	-14		-27	-02	
Bristol .....	+02		-22	-30		-03
Plymouth .....	-05		-27	-23		+02
London .....	-02		-04	-46		-18
Paris .....	-05		-20	-20		-16
Christiania .....	-14		+20	+11		00

The elevation of the barometer indicated by the foregoing observations, though gradually decreasing from the 4th, did not subside till the approach of a storm from south-west (§15); which began at the Orkney islands on the night of the 7th, was blowing at the island of Mull on the 8th, and arrived at Glasgow at 6 P.M. of the same day; in the north of Ireland (Donegal) at 8 P.M.; and at North Shields (a little to the north of Donegal in latitude, but on account of its more easterly position later in receiving the storm) at a very early hour on the morning of the 9th. The change of the wind as the storm progressed is well-marked by the diagrams. On the 7th, the wind remained north-east throughout the day in almost the whole of England, but at the Orkneys had changed to north-west (§ 18), and was variable at Mull island in the evening: at North Shields it changed to north-west after mid-day, and at night the storm began at the Orkneys from south-west. On the 8th, the south-west wind is blowing in the greater part of Scotland, whilst to the southward the wind is still north, but in some cases north-west. In the evening the changes before noticed take place, the wind being yet northerly in the south. On the 9th the storm became prevalent throughout, on which day the barometer attained its minimum in the north (§ 3), its height in the south being very little reduced (§ 2), although it appears that on that day the wind was blowing as strongly in the south as in the north (§ 4).

The approach of the storm from north is seen also by the falling of the barometer, as indicated by the observations. At North Shields the barometer attained its minimum (there 29.276) at 5 P.M., and at 9 P.M. it had risen 0.10, though reckoning simply from the extreme observation, it had not risen at all; whilst at Orkney it had risen 0.20, and in the south it was yet falling (§ 3). On the morning of the 10th the barometer at the Orkneys had risen 0.39, with a strong gale from north-west (§ 3 and 12); but at Shields, from 5 P.M. on the previous day to about the same hour on this, although the barometer rose, the air was almost calm. The diagram, however, together with the barometric heights, fully explains this; for we see that its position was that of the meeting of the two currents, the north current blowing on the north and the south one on the south (§ 11), the latter continued by the state of the barometer; the barometer rising however by reason of the strength of the north wind setting in in the north (§ 6). The north wind in the middle and southern parts of Scotland appears (on account of the low state of the barometer at Orkney) to arise from the impetus which it has received in blowing in the



extreme north, together probably with a higher barometer on the east (shown by its easterly deflection): thus we find the wind decreases in strength as it advances, being at Mull island only brisk. The equality in the force of the two currents is not however of long continuance, for that of the south wind, evidently because of the great depression of the barometer in the north below its height in the south (Orkney, barometer 28·70, London 29·81), on the 9th, greatly increases and advances as a storm towards the north. But this difference of the atmospheric pressure requiring, on account of the great distance of the localities of the extremes of pressure, and the resistance of the opposite current blowing in the north, a long interval of time to produce its effect, does not arrive at North Shields till the latter part of the 10th as a S.E. or rather S.S.E. storm (being deflected by its collision with the contrary current) (§ 11 and 16), and continues until about the same time of the 11th.

In the phænomena now before us we have a good example of an advancing portion of a storm (§ 16). The fall of the barometer, which continues in the south whilst the rise is going on in the north, increases on the 10th, and advances, together with the wind, towards the north, where, excepting in the extreme north, it falls to a greater degree than before, and, as noticed in § 16, the greatest depression is in the south. The progressive motion of the storm may however be traced in both the directions of the figure of § 15, but the south-east movement is in the south: thus at Cork and Plymouth the minimum depression occurred on the evening of the 10th, at Bristol about noon of the 11th; for out of four observations that at noon was the lowest (29·99); and at Paris (south-east) and at North Shields (north-east) it happened simultaneously about 9 P.M. Hence if we suppose Cork to represent the point C, fig. 3, then the line  $cC$  prolonged would extend to Paris, the storm however diminishing in intensity, and the line  $CB$  from Cork to North Shields. The limit of this storm may be observed in the north; for at Orkney the wind continued north-east, and at Mull island east, but yet the barometer at Orkney falls, though during a north-east wind, but not to so great a degree as the next station (Glasgow) (§ 5), for on the next day (the 12th) the minimum depression, or the point C, was in the south of Scotland or north of England; hence Orkney would represent the point E. of fig. 1.

The diagrams of the 10th and 11th (Plate VII.) are also of interest as regards the deflection of the currents produced by their meeting: on the north the wind is N.E., in the south

of England and in Ireland it is S.W., and at the localities between the two it is S., S.S.E., or S.E.

Names of Places.	12.			13.			14.		
Orkneys .....	-07		+09	+12		+18	+23		+19
Glasgow .....	-03		+27	+07		+05	+52		+07
Belfast .....	+10	+09		+11	-05		+56	+08	
Armagh .....	+19		+14	00		+20	+40		-05
Shields .....	+05		+26	+07	-03	-02	+46	+11	+13
Cork .....	+39	+01		-21	+30		+26	-06	
Bristol .....	+13		+33						-10
Plymouth .....	+17		+33	-19		+05	+26		-14
London .....	+12	+21		+18	-25		+46	+08	
Paris .....	+13		+27	+09		-27	+26	+02	-02
Christiania .....	-15		-21	-13		+01	+19		+38

Names of Places.	15.			16.			17.		
Orkneys .....	+11		+08	+10		+15	+08		-06
Glasgow .....	+01		-02	+14		+23	+20		+02
Belfast .....	-06	-03		+14	+10		+41	+04	
Armagh .....	-03		-05	+18		+31	+23		+05
Shields .....	00	-01	+02	+12	+06	+17	+25		+11
Cork .....	-20	-03		+12	+21		+60	+01	
Bristol .....	00		+01	+07		+21	+37		+21
Plymouth .....	00		+06	-04		+15	+47		+22
London .....	-11	-07		+14	+03		+44	+13	
Paris .....	-06	-02	+01	-05	-03	+04	+22		+40
Christiania .....	+15		+03	+08		+12	-06		-03

The phenomena on the 12th commence a period during which the restoration of the atmosphere to its usual pressure and the rise of the barometer above its mean elevation take place; but their chief interest is in their being those ensuing on the cessation of an advancing storm. The occurrence of the minimum of the atmospheric pressure in the south before its taking place in the north has already been noticed, and is also very apparent from the whole of the observations. Thus at Cork, which seems to represent the point C, § 16, as being in the line of the greatest intensity of the storm, the barometer at 9 A.M. has risen 0.39 inch, and the rise lessens in both directions towards south-east, or along the line *c* C of fig. 3 prolonged, and towards north on the line C B; and in the latter direction at Orkney the barometer still continues to fall, though very slightly. In accordance with this state of the barometer (referring to the same paragraph), the wind is blowing strongly from north-west in the south-west and middle portions, though still opposed by the south wind on the extreme south. In the north, we have clear evidence of the extensive low state of the barometer on the west, for at Christiania, about 10° to the east of the Orkney islands, the height

of the barometer at 9 A.M. was 29.30 inches, and at Orkney the wind is from east, hence at the same distance on the west of these islands the barometer is probably below this. If we now suppose, whilst the wind is blowing as shown by the diagram, the point C of fig. 3, by the motion of the line A C (§ 16), to move considerably northwards, and to be a little to the north of Holyhead, where the wind is strong from north-west, the phænomena will be simply a particular case of the general result of the paragraph, for the wind is strong from north-west on the localities in the direction which would be that of the line C A, and variable between west and north-west at Flamborough-head, and west-north-west and very light at North Shields, places, which with regard to the figure would be nearly on a horizontal line with the point C; and in the north, but not extending to Orkney (where the barometer is just beginning to rise), the wind is variable between south and west, and extremely light; but at the locality nearest to Orkney, Greenock, it appears to change to north-east in the latter part of the day. This however is the extreme portion of the storm, and accordingly, soon after this, the north wind sets in briskly from north-east instead of from north-west.

It may be observed that in the south the rise of the barometer ceases at Cork whilst it continues at Plymouth; the phænomena of the next day (13th) however, explain this, for by the setting in of the current from north-west on the south-west, which opposing the south wind blowing in the more southern parts, causes a very rapid rise of the barometer in the south, but more especially in the south-west, a disproportionate pressure there is again produced, and the consequences are in some degree the same as before; for on the 13th, although the barometer rose rapidly at the Orkney islands, and the north-east wind fully set in in Scotland, the south wind blowing previously only in the extreme south, increases in strength and becomes prevalent in all the southern part, causing a considerable fall of the barometer in the south, which, as before, occurs first at Cork, confirming the view that the rarefaction of the atmosphere is greatest in the north-west. At North Shields, where on the night previous the wind was strong from the west, it changed to south-west, though extremely light, and caused a slight fall of the barometer between 9 A.M. and 2 P.M.

The phænomena presented by the diagrams of this day, together with the variations of the barometer given in the first column of the 13th, though of the same kind as in two previous instances, afford so striking an example of the case

of § 2, because of the blowing of both currents being so fully pointed out, that I cannot pass it over. We see that on the south the south wind is prevalent and strong, and the north is blowing as a fresh breeze in the north, the two currents meeting and balancing one another a little to the north of the centre of the field of observation, and yet in this place the barometer falls, the fall increasing towards south on account of the greater height of the barometer there at its commencement; but at the same time a rise takes place in the north. Now it is certain that if these winds were simply the flow of exactly similar currents, the one flowing from north and the other from south to a space between them, on this space the barometer would rise. What then becomes of the air brought to the place of meeting if the southern current does not carry it off in the upper regions of the atmosphere, as shown in fig. 1 by the upper arrows between *c* and *b*? That it does not arise from any atmospheric change, originating in a central portion, such as a change in the elasticity of the atmospheric columns, causing a portion of air to roll off from their upper parts and a current to set in towards their bases, is very evident, because the diminution of pressure begins and is greatest at the most remote parts of the south wind flowing towards it; but if we admit the explanation given by § 2, the phænomena presented by the barometer are perfectly consistent with the action of the two contrary currents, which appear to have met so directly that a calm, or a state of the air nearly approaching to it, is produced\*.

On the afternoon of the 13th the north wind becomes the most prevalent, and the barometer rises rapidly throughout,

\* As the north and south winds are deflected, the one from east and the other from west, the relative position of England and Scotland might at first sight give rise to the opinion, that when the north was blowing in Scotland and the south in England, they do not blow in opposition to each other, but in parallel bands; in the cases of the 13th and 15th, however, as well as others in which this opposition has been remarked, the observations in Ireland and the extreme west of England remove all doubt as to the actual collision of the currents, for we see by these the two currents blowing directly towards each other in the more remote parts, and variable winds, calms, or the deflections of the south current from east, near the place of meeting.

It is evident, however, that when the north-east wind prevails in a much greater degree on the eastern parts than on the western, as appears to be the case in some days of this period, the opposite currents may blow in parallel bands for some distance; but on the parts immediately adjacent to the north wind, the south wind will be south-east; hence a northern locality may have a south-east wind when a north-east blows on a southern one more to the east, a case frequently occurring in this country, as on the 2nd and 3rd.

although in the south during the blowing of a south wind (§ 6). On the remaining days of this series the north wind continues on the whole to gain in predominance over the south, and to cause the barometer to rise, not however without a check in the south, where the south wind again increases in strength for a time, and causes a slight depression of the barometer on the 14th and 15th, which has its limit northward in the north of England or south of Scotland, where the force of the wind is balanced by the opposite current, repeating the phænomena of the 13th, though with this difference, that on the 15th the north wind prevails to a greater degree than on the 13th, so that at North Shields the north-east wind itself is blowing; still however there is a slight fall of the barometer (§ 5). On the 16th the south wind has greatly decreased, and serves only to produce the great comparative rise of the barometer at Orkney (§ 7), where it attains a height considerably above the mean. On the 17th, the day on which the barometer attains its maximum elevation, we have very little indication of the south wind blowing on the south; it is however blowing on the west, and accordingly we find the elevation beginning in the north-west and extending itself towards the south-east; now as this is the direction in which the point of C of fig. 1, whether the point of depression or elevation moves, and this elevation differs from that of the 4th, during which also the opposition of the south wind was on the west, in extending eastward and south; and also as in many previous instances the south wind merely retreated, and did not altogether disappear when the north wind advanced; we may infer that in the present case it is yet blowing in localities southward of the latitudes included in the diagram. If not, the phænomena yet admit of easy explanation, on the supposition that the north-east wind, which is blowing with great strength, occupies (§ 9) a greater proportion of the height of the atmosphere than the equal flow of the upper current admits of.

Names of Places.	18.		19.			20.		
Orkneys .....	-·16		+·05	-·31		+·03	+·04	-·02
Glasgow .....	-·05		-·14	-·43		-·02	+·12	+·02
Belfast .....	-·03	-·11		-·44	-·07		+·08	-·03
Armagh .....	-·07		-·36	-·14		-·04	·00	·00
Shields .....	-·05	-·04	-·12	-·46	-·11	-·15	+·12	·00
Cork .....	-·02	-·11		-·20	-·06		-·16	+·04
Bristol .....	+·03		-·14	-·22		-·31	·00	
Plymouth .....	+·04		-·08	-·15		-·27	-·18	-·20
London .....	+·22	-·05		-·30	-·17		-·27	-·02
Paris .....	+·19	·00	+·03	-·11		-·26	-·44	-·70
Christiania .....	-·13		+·06	-·14		-·18	-·18	-·08

Names of Places.	21.			22.			23.		
Orkneys .....	·00		-·18	-·30		-·08	-·03		-·06
Glasgow .....	+·02		-·20	-·40		-·10	+·05		-·16
Belfast .....	+·10	-·06		-·53	·00		+·04	-·14	
Armagh .....	+·10		-·31	-·23		·00	-·02		-·49
Shields .....	+·04	-·01	-·12	-·36	-·12	-·15	+·04	-·04	-·12
Cork .....	+·04	-·01		-·25	·00		-·24	-·24	
Bristol .....			-·07	-·46		+·19	-·01		-·44
Plymouth .....	+·15		+·01	-·34		+·17	-·03		-·57
London .....	+·03	+·02		-·40	-·14		+·22	+·09	
Paris .....	+·01		+·12	-·32	-·18	-·01	+·20		-·17
Christiania .....	·00		·00	+·02		-·02	+·05		+·02

Names of Places.	24.			25.			26.		
Orkneys .....	-·19		-·03	-·04		+·06	·00		·00
Glasgow .....	-·30		-·13	+·05		+·07	+·06		+·14
Belfast .....	-·49	-·04		-·01	+·08		+·15	+·07	
Armagh .....	-·23		-·07	+·07		+·14	+·12		+·15
Shields .....	-·30	-·01	-·14	+·05	+·02	+·03	+·08		+·16
Cork .....	-·56	·00		+·25	+·01		+·20	+·04	
Bristol .....	-·21		-·08	·00		+·19	+·10		+·20
Plymouth .....	-·07		-·08	-·04		+·20	+·10		+·15
London .....	-·68	+·01		-·01	-·02		+·21	+·08	
Paris .....	-·35		+·09	-·12		+·13	-·01		+·21
Christiania .....	+·02		+·02	-·01		-·08	·00		+·02

Having now given a detailed explanation of the several phenomena of the preceding observations, it will be sufficient to give a very general account of those of the succeeding ones; they are however of great interest. The first in occurrence, whose approach is indicated by the p.m. observation of the barometer at the Orkneys on the 17th, is a depression of the barometer by a south wind of just sufficient strength to be called a storm, and its subsequent rise; both progressing from north-west to south-east. Before the restoration of the usual pressure of the atmosphere a second storm occurs, prevailing in Ireland on the night of the 21st, and in England on the 22nd; and whilst the storm was blowing in the latter country the returning current set in strong from north-west in the former, raising the barometer there to a height considerably above that in England (§ 16); its height however on the following day was rapidly reduced by the setting in again of the south wind, as we have before seen in cases of a disproportionate elevation of the barometer in the south, and continued the following day (the 24th). Both these storms, but more particularly the latter, approached this island from about the north of Ireland, as appears from the fall of the barometer occurring first at Armagh. It must be recollected however that the observation there is registered at 10 p.m., thus one hour and

a half later than that at the Orkneys; and as the observation at 3 P.M. at Belfast shows that it occurred between these hours, the difference in time accounts for part of the difference in the variation of the barometer; but in accordance with this view of the direction of its motion, we observe that the greatest depression of the barometer, as shown by the observations, is at Glasgow, and therefore, if not precisely there, it must be somewhere between this and the Orkney islands, and hence it is that the south wind during these storms very seldom extends to the north of Scotland, and that the north wind is generally prevalent there; and when it is not, the south wind is very inconstant, as there is always a northerly direction given on the same locality, excepting when the wind is simply stated east, when there can be little doubt it was from the north of that point; and if not so, it at least shows that the south wind had little strength. On the 23rd and 24th, however, the days on which the principal depression of this period occurred, the north wind in the north is well-marked, whilst the south wind is blowing in the south. The latter of these storms offers an example of the case of § 16, but one in which there is a more equable division between both portions of the storms of § 15.

The two remaining days of the period represent by similarity, the remaining portion of the month not included in the observations given; the weather continued stormy, and the barometer fluctuating according to the prevalence of the north or south wind, the north however being on the whole predominant, so that the barometer attained its mean height on the 1st of December.

As a very remarkable depression of the barometer occurred on the 13th of January, 1843, I included the first fifteen days of this month in my collection of observations, but with the exception of the storm which occasioned that depression, they do not offer anything sufficiently worthy of notice, after what has already been given, to make it necessary to insert them here. That storm however I notice particularly, because it presents an additional illustration of the action of the storms of § 16, its phenomena confirming the account given of storms, derived from a consideration of those of the 10th and 11th of November 1842. It advanced to the north of England from a line running somewhat in the direction from Cork to Plymouth; thus at Plymouth it began at 11 P.M. of the 12th, and at North Shields about half an hour after 5 A.M. of the 13th, six and a half hours later; but in the south its progress towards south-east, or its recession along the line *c C* of fig. 3, is well-marked, for it began at Cork at 7 P.M. of the 12th,

four hours sooner than at Plymouth, and at Portsmouth at 2 A.M. of the 13th, three hours later than at Plymouth. At London and Bristol it began at the same hour as at Portsmouth, 2 A.M.\*, so that its progress in both directions is clearly seen.

In this case also, as in the previous one, the storm succeeded a great depression of the barometer in the north. The order of time in which the minimum height of the barometer was attained, coincides exactly with that of the beginning of the storm, and is very conspicuously marked in both directions; thus in that of the receding portion, or along the line *c C* of fig. 3, the time of the minimum at Cork was early in the morning, or during the night; at Falmouth 9 A.M.; at Plymouth 10 A.M., and at Paris about noon; again in the direction of the advancing portion, or along the line *C B*, it was two hours later at Bristol than at Falmouth, and at Shields five hours later than at Bristol. I have not thought it necessary to give a diagram showing the directions of the wind, because they may be so easily described. The wind was south-west in the south of Ireland and of England, east and north-east in the north of Scotland†, and south-east or south-south-east (§ 11) between the two extremes, as in the north of England and south of Scotland.

This storm affords also an illustration of § 4; for though, as on the 11th and 12th of November, 1842, the height of the barometer at its cessation was not very far from equal throughout, the reduction was greater in the north of England than in the extreme south, yet in the former region it was a storm of short continuance and no extraordinary violence, whilst in the latter it is described as a perfect hurricane.

*Height of Barometer, January 1843.*

Names of Places.	11.		12.		
Orkneys .....	28·76		28·79	28·84	28·82
Glasgow .....	28·75		28·65	28·74	28·70
Belfast .....	29·10	28·99		29·02	29·05
Armagh .....	28·85		28·57	28·78	28·55
Shields .....	28·80		28·81	28·84	28·90
Bristol .....	29·14		28·78	28·84	29·13
Cork .....	29·08	28·88		29·20	29·20
Falmouth .....	29·24		28·83	28·80	29·22
Plymouth .....	29·22	29·00	28·79	28·71	29·17
London .....	29·11	29·02		28·74	28·97
Paris .....	29·12	29·03	28·79	28·67	28·77
Christiania .....	28·55		28·67	28·83	28·98

\* Shipping Gazette Newspaper, January 1843.

† At Tobermorey (Mull) a hard gale from E.N.E. (§ 2), at North Shields S.S.E., and in Leith Roads a gale from E.S.E.



Names of Places.	13.		Minimum on 13th.	Time of minimum.	14.			
Orkneys .....	28·57	28·51			28·52	28·51		
Glasgow .....	27·90	28·12			28·56	28·61		
Belfast .....	28·10	28·22			28·90	28·84		
Armagh .....	27·83	28·48			28·61	28·64		
Shields .....	28·18	28·05	28·15	28·032	4 P.M.	28·62	28·77	
Bristol .....	28·18		28·75	27·975	11 A.M.	29·03	28·85	
Cork .....	28·58	28·81			*	28·70	28·75	
Falmouth .....	28·45	28·78	29·03	28·45	9 A.M.	29·08	29·04	
Plymouth .....	28·44	28·70	29·01	28·436	10 A.M.	29·17	28·72	29·01
London .....	28·35	28·32				29·05	28·74	
Paris .....	28·79	28·74	28·93	28·74	about noon	29·19	28·68	
Christiania .....	28·96		28·91			28·71	28·56	

Variations of Barometer.

Names of Places.	11.		12.		13.		14.		
Orkneys ...	+·28	+·03	+·05	-·02	-·25	-·06	+·01	-·01	
Glasgow ..	+·19	-·10	+·09	-·04	-·80	+·22	+·44	+·05	
Belfast ...	+·06	-·11	+·03	+·03	-·95	+·12	+·68	-·06	
Armagh ...	-·03	-·28	+·20	-·23	-·72	+·65	+·13	+·03	
Shields ...	+·01	+·01	+·03	+·09	+·09	-·75	-·13	-·03	+·47
Bristol ...	-·01	-·36	+·06	+·29	-·95	+·57	+·28	-·18	
Cork .....		-·20	+·32	·00	-·62	+·23	-·11	+·05	
Falmouth .	-·10	-·41	-·03	+·42	+·43	-·78	+·33	+·58	+·03
Plymouth		-·22	-·43	-·08	+·46	+·51	-·78	+·26	+·57
London ...	+·06	-·09		-·28	-·23	+·44	-·62	-·03	+·73
Paris .....	-·08		-·33	-·12		-·32	+·14	+·26	-·51
Christiania		+·12	+·16		+·15	-·02	-·05	-·20	-·15

An extraordinary depression of the barometer (height 28·612) for that latitude occurred about the same time at Paris† as that in England; it however occurred at 6 A.M. of the day previous to that in this country, for which reason I have noticed it here, as at first sight it might seem opposed to the general order of the phænomena presented by the foregoing observations; but it did not occur during “the hurricane,” for the wind in the Paris meteorological register for the 12th is quoted merely “strong,” but on the following day, when the fall corresponding to the great depression in England took place, the wind is quoted “very strong,” and the tables show also a fall of the barometer in England on the 12th corresponding with that at Paris.

I may now conclude this part of the subject by remarking that observations for other periods than those chosen might exhibit difficulties which do not appear in these, although I have made no selection in publishing them, but merely given those I collected on account of the period included in them

\* Time of minimum not noted but during the night of 12-13.

† *Annales de Chimie et de Physique.*

being more than usually stormy. Whatever difficulties do arise, however, it must be borne in mind that the situation of this island is peculiar, having in summer a temperature below that of parts of Europe on the north, and in winter above that of part of the continent on the south; hence a very considerable complexity in the directions of the currents must occasionally occur.

The results given in the first portion of this paper are deduced from the simple fact of the descent of the upper current of the atmosphere, and are altogether independent of the immediate cause of its descent; that it must always be descending in some portion or other of its course, to supply the place of the air flowing in the surface current towards the equator, is very evident; but the indications of the barometer show that the acting cause of its descent at any particular time is not—always at least—a deficiency or rarefaction of the air, such as would be occasioned by the flowing of the lower current, were its effect uncompensated by the arrival of air from above, for the south wind often sets in when the barometer is high. I have alluded to this subject before (*Phil. Mag.* Oct. 1843, p. 280), noticing the effect of the difference of the opposite currents with respect to the quantity of aqueous vapour in each; but though there appears no reason to doubt that effect being, in a greater or less degree, as there supposed, I stated a difficulty which it is probable does not exist, that of the descending air being warmer than the air previously in its place—an opinion derived simply from the fact of the upper current having, from its origin, generally the higher temperature. Now it may very often be observed, though not always, that the temperature does actually become colder immediately at the change of the wind from north to south, though it rises again on the continuance of the wind; for when once it has found its way to the surface, then of course, whatever its temperature subsequently, it will continue there until its force be overcome, one portion making way for the next following. The cases in which the temperature rises immediately on the change of the wind may be those in which the change either takes place, not from an immediate descent of the current, but simply from its advance (§ 16) from southern localities, where it has previously descended, or perhaps been blowing for some time, and advanced by reason of an increase of force; or by its recession (§ 15) from north\*.

\* This is a distinction which must be carefully borne in mind during the reading of these remarks, and during all consideration of this subject, in order to guard us from drawing inferences in any one particular case of a change of wind, when its nature in this respect is not known.

Now the south wind descends or blows in the greatest degree during the winter half of the year \*, and then most at night; that is, when the temperature of the earth and the atmosphere is falling by radiation; whilst in spring and summer—the temperature increasing—the lower current, or north-east wind, prevails more than at any other time of the year. Now it is obvious that were the cooling of the atmosphere unaffected by that of the earth, its upper strata, notwithstanding the facility with which heat passes through it, would be first and most cooled by radiation; but we know that within small heights—to the extent of about 100 feet—above the surface of the earth, in which such observations can be easily conducted, the lower strata are much the coldest in nights when radiation is vigorous, by reason of the cooling of the earth. It is obvious however that it does not at all necessarily follow from this that this increase of temperature in ascending, or rather as it would be after a certain height, increase compared with the general progressive decrease of the temperature, should go on towards the higher regions of the atmosphere; for it is evident, from the very great rapidity in which the temperature decreases towards the surface, that it is very much, if not almost entirely, owing to contact; but on the contrary there will be a certain elevation, perhaps not very great, at which the cooling of the atmosphere by the greater radiation of heat into space in the strata above than in those below it, is so much greater than the cooling in the air below, occasioned by radiation to the earth, that the previous relation of the temperatures of the air of the two currents may become at any time, when this difference is not very great, reversed, and the upper one, being then comparatively the colder, descends. The cooling may also be materially affected by the presence of clouds, and by the heights above the surface of the earth at which they are formed; as these would present a comparatively dense radiating body, and by cooling the particles of air in contact with them, would cause currents of cold air to

\* The blowing of the wind from west, whether north or south, may of course (except in the case (§ 12) where it is caused by the flowing of air to restore the atmospheric pressure as in storms) be taken as evidence of the descent of the upper current; thus in the cold months of the year, although the north wind is strongly urged on towards warmer regions by the greater difference of temperature between adjacent latitudes, due to the season, and frequently prevails,—it is generally from north-west.

The phænomena of tropical regions correspond to those of high latitudes; thus the time of descent, or of hurricanes, is after the sun has attained its greatest northern declination, consequently when solar radiation is decreasing; the temperature however is very little fallen, and is in somewhat of the variable state of high latitudes, the blowing of the trade-wind being then nearly suspended.

descend; and also prevent radiation from the earth and atmospheric strata beneath them. Admitting such a cause of descent of cold air, we shall have an explanation of those sudden colds often experienced, which cannot be accounted for on the supposition of the arrival of air from colder regions.

But in the summer half of the year the time of descent of the upper current is not in the night more especially than during the day.

Being desirous of confirming the opinion that the south wind prevailed most at night, I selected from my own register of the wind, made three times a day, all the changes of wind from north to south and from south to north for one year, noting the time (whether night or day) at which they occurred, and met with the unexpected result, that whilst with regard to the winter months the opinion was amply confirmed, in those of summer the result was rather the opposite to it, the change on the whole taking place most frequently in the daytime; and in order to confirm this conclusion, the changes were selected from two other years, with the same result. I have presented in a table the average number of changes for the three years, merely remarking that with scarcely an exception the same month of each of the three years gave a result in accordance with the mean one. The table also contains the changes from south to north, which are in general opposite to the contrary ones.

*Number of Changes of Wind.*

Month.	Night.		Day.		Month.	Night.		Day.	
	From N. to S.	From S. to N.	From N. to S.	From S. to N.		From N. to S.	From S. to N.	From N. to S.	From S. to N.
Jan.	7·3	1·7	1·7	6·0	April	2·3	5·3	6·3	2·7
Feb.	3·7	2·0	1·7	3·7	May	4·0	4·0	3·3	3·7
March	6·3	4·3	2·3	4·3	June	3·3	2·7	1·3	1·3
Oct.	4·3	3·3	2·7	3·3	July	3·3	5·3	5·3	4·3
Nov.	5·3	1·3	1·7	6·0	Aug.	3·7	5·0	5·0	3·7
Dec.	5·0	4·0	2·3	3·3	Sept.	3·7	3·0	2·3	3·3
Total	31·9	16·6	12·4	26·6	Total	20·31	25·32	23·6	19·0

The want of exact consistency which appears in the results of the summer months is easily accounted for by the lightness of the wind and its arising often from merely local circumstances; the total result however is sufficiently decided to show the difference of character of the winter months. Now in summer the temperature of the lower strata of the atmosphere with respect to those above, from the heat derived from the ground, is comparatively much greater than in winter,

the difference of course being greatest in the daytime; thus then the daytime is more favourable for the descent of air from the upper strata than in winter.

Hence then I think it may be concluded that the air of the upper current becomes relatively colder than the lower strata of the atmosphere by loss of its heat by its own radiation, and that when the cold has arrived at a certain degree, it descends, if other conditions which influence its descent are favourable; these conditions being the state of the temperature and pressure of the air of adjacent latitudes, by which the force urging forward the surface current of the atmosphere is affected; and the state of the opposite currents with respect to aqueous vapour.

XLVI. On the Derivation of the Word Theodolite.

By Professor DE MORGAN\*.

THE word *theodolite* has puzzled all who have tried to trace it to its origin. Some have connected it with the roots of *θεάομαι* and *δολιχός*, and made it a *seer of lengths*, though the instrument neither does, nor ever did, see anything but angles. In a modern dictionary of good reputation, it is connected with *θεάομαι* and *δόλος*, and made a *seer of stratagems*, which might apply to a telescope: but unfortunately the use of the term *theodolite* was prior to the invention of the telescope.

The word is exclusively English, never having obtained any mention from foreigners till comparatively recent times. The *Encyclopédie Méthodique* (1789) does indeed give the word without allusion to its origin; but Savérien's dictionary (1753) says that the *theodotile* (as it is spelt) is an instrument used by the English, much resembling the *graphomètre*.

I find that the use of the word runs back to the "Geometrical practise named Pantometria," begun by Leonard Digges, and finished by Thomas Digges his son (published London 1571, quarto, reprinted in 1591). But it seems as if the name was not then new. Chapter 27 is on "the composition of the instrument called Theodelitus," and it is plain from various modes of speaking that the word is here an adjective or participle. This "circle called Theodelitus," or "planisphere called Theodelitus," is nothing but a graduated circle with a revolving diameter furnished with sights, and placed horizontally. Held vertically, it would have been the astrolabe of the period, and nothing else. In Leybourn's 'Compleat Surveyor,' 1657, we learn that the altitude circle was *sometimes*

\* Communicated by the Author.

added; and in Stone's Mathematical Dictionary (1726) that it was *sometimes* furnished with a telescope.

A ruler with sights, travelling upon a graduated circle, was a constituent part of various astronomical instruments imported into Europe from the East, and was accompanied by the Arabic term *alhidada* to express it. The word *alidade* or *alhidade* (for it is spelt both ways) is completely naturalized in France, and appears in the common dictionaries. It was also used by the English writers of the sixteenth century, and among others by Digges himself. The original *theodolite* being nothing but a graduated circle with an *alidade*, some connexion between the terms might be suspected by those to whose notice they are brought. But so different do the words appear, that I, for one, should never have been reminded of the first by the second, if I had not happened to find, in a writer contemporary with Digges, an intermediate formation, which brings the two words nearer together. William Bourne's 'Treasure for Travailers' was published in 1578; he does not use the word theodolite, but calls the instrument the "horizontall or flatte sphere." He begins by spelling the word *alhidada* thus, *alydeday*, but soon changes it, and keeps very steadily to *athelida*, which is the only technical term introduced in his description of what Digges calls theodolitus. From these premises, I cannot help inferring that the *theodelited* circle of Digges, and the *athelidated* circle of Bourne, which are certainly the same things, are but described by different corruptions of the Arabic word whose earliest European form is *alhidada*.

In our day such a transformation might not be easy; but when the works above-mentioned were written, nothing was more common than to spell the same word in two different ways in the course of one sentence. Bourne himself, though he sometimes spells the name of Digges's work correctly, *Pantometria*, yet in the first place in which it occurs, he makes *Pantometay* of it, possibly a misprint for *Pantometry*.

The fact seems to have been thus in this and many other instances. In the sixteenth century, before the language was well-settled, an author more accustomed to Latin than English, would try to anglicize some technical terms; and, not finding his results please his own fancy, would then fall back upon the Latin. Bourne has done this with both *athelida* and *pantometria*; and, were it worth while, I could show abundance of similar instances in other writers.

Nor is it against the connexion of the words that Digges uses them both. Instances are not wanting in which two different spellings of the same word are used by the same writers

for different things. For example, the original English sense of the word *square* applies to an angle, not a figure; a right angle is a square corner; and to this day the carpenter's right angle is called a square. But I could name half-a-dozen writers of the end of the sixteenth century who use the two spellings *square* and *squire*, the former in the modern sense, the latter for the carpenter's instrument.

---

XLVII. *Reply to the Observations of M. Pierre, on the Proportion of Water in the Magnesian Sulphates and Double Sulphates.* By THOMAS GRAHAM, Esq., F.R.S.\*

**I**N a late number of the *Annales de Chimie*, a paper by M. Isidore Pierre appears, On the Double Salts formed by the Oxides of the Magnesian Group, of which an abstract is also given in the March Number of the Philosophical Magazine, containing statements which demand some remark from myself. It presents new analyses of the sulphate of magnesia and potash, and other double sulphates of the same type, from which the author infers that these well-known double salts possess seven atoms of water crystallization, and not six atoms, as resulted from my own analyses and the analyses of all other chemists who have of late years examined these salts. The double salts in question are thus made by M. Pierre to have the same proportion of water as sulphate of magnesia itself; while the latter salt, also, is not found to retain its seventh atom of water more strongly than the other six, but to become anhydrous at  $212^{\circ}$ , or a few degrees above that temperature, in a current of dry air. The author then infers that his results are subversive of the theory which was originally published by myself, of the constitution of the magnesian sulphates, and to which I still adhere, namely that they contain an atom of water strongly attached and not easily expelled by heat, but readily replaced by an alkaline sulphate, with formation of a double salt.

Although confident of the accuracy of the analyses thus impugned, I considered it due to M. Pierre, who, although a young chemist, has afforded every evidence of habitual care and accuracy in another experimental inquiry of importance, to repeat my experiments.

Of the double sulphate of zinc and potash, 31.46 grains by drying at  $212^{\circ}$  for several days, lost 7.75 grains of water; and by fusion at a heat verging on redness, 0.08 grain of water additional, making the whole loss 7.83 grains. Hence the composition of the salt with reference to water is as follows:

\* Communicated by the Author.

	Experiment.	Theory of 6HO.	Theory of 7HO (Pierre).
Water . . . . .	24.89	24.03	27.32
Sulphate of zinc and potash	75.11	75.97	72.68
	100.	100.	100.

The experiment obviously indicates six and not seven equivalents of water. The slight excess of 0.86 per cent. of water is not more than is usually found in crystallized salts, arising from the difficulty of divesting them entirely of water mechanically interposed between the plates of the crystals. The peculiarly high disposition of this particular class of salts to retain mechanical water, has been noted by Mitscherlich, myself, and almost every one else who has made them the subject of investigation. It has probably been the cause of the error into which M. Pierre has fallen, in over-estimating their proportion of water.

Although it is scarcely necessary to extend the inquiry to the other double salts of the class, which being isomorphous with the last have necessarily the same proportion of water, still I may be allowed to avail myself of a series of five analyses of the double sulphate of copper and potash lately executed in the laboratory of my friend Prof. Fownes, and which he has kindly communicated to me.

	Expt. 1.	2.	3.	4.	5.	Theory of 6HO.	Theory of 7HO (Pierre).
Water.....	25.20	24.00	25.00	25.2	24.4	24.44	27.40
Sulph. of copper and potash ...	74.80	76.00	75.00	74.8	75.6	75.56	72.60
	100.	100.	100.	100.	100.	100.	100.

These experiments all concur in proving that six equivalents is the proportion of water in the double sulphate of copper and potash, and not seven equivalents.

Although M. Pierre gives seven atoms of water to the double sulphate of magnesia and potash, he adds, near the end of his paper, as if to qualify the statement, that when he communicated his results to M. Balard, that chemist informed him that the double sulphate of magnesia and potash contained no more than six equivalents of water, and was therefore consistent with the views of Mr. Graham.

With reference to the single atom of water strongly retained by the magnesian sulphates, an experiment was made on sulphate of zinc. The crystallized salt dried for several days at 212°, in the same circumstances as those in which the double sulphate of zinc and potash became anhydrous, still retained water. The heat being continued for three or four days after



the salt ceased to lose weight, it was thereafter found to consist of—

	Experiment.		With one equivalent of water.
Sulphate of zinc . . . . .	20·42	89·20	90·03
Water . . . . .	2·46	10·75	9·97
	<u>22·88</u>	<u>100·</u>	<u>100·</u>

It is sufficiently evident, therefore, that sulphate of zinc, which is admitted by M. Pierre to contain seven equivalents of water, retains one equivalent of water by a stronger affinity than the other six, contrary to his observation; while, moreover, this strongly retained atom of water is absent in the double sulphate of zinc and potash, the last containing only six atoms of water—the experimental data on which the view of the constitution of these salts controverted by that chemist is founded.

XLVIII. *On the Cohesion of Liquids and their Adhesion to Solid Bodies.* By M. F. DONNY, *Agrégé à l'Université de Gand, Préparateur du Cours de Chimie.*

*To the Editors of the Philosophical Magazine and Journal.*

GENTLEMEN,

THE twenty-sixth volume of your valuable Magazine (p. 541) contains an account of two communications made by Prof. Henry to the American Philosophical Society, on the 5th of April and 17th of May 1844, both relative to the cohesion of liquids.

I have been investigating the same subject from the beginning of 1841 to the end of 1843, when I gave a full description of my experiments on cohesion and adhesion in a written communication addressed to the Académie Royale de Bruxelles. The reception of this memoir is recorded in the *Bulletin de la Séance du 2 Décembre 1843* (tome x. p. 457), and the memoir itself is printed in the *Mémoires Couronnées et des Savants Etrangers*, tome xvii. I beg leave to direct your attention to the contents of this communication.

Having discovered, in 1841, as Prof. Henry did in 1844, that the cohesion of liquids is a powerful attraction, entirely misrepresented in the works on natural philosophy, I endeavoured to find out the cause of this misrepresentation. With this object I constructed a very simple instrument, which enabled me to observe accurately how the separation of water from water is effected, in the well-known experiment of a

plate suspended from a scale-beam over a vessel of water. The use of this new instrument convinced me immediately that there is no similitude whatever between the rupture of a solid body and this mode of separating water from water. I perceived plainly that such a separation was the final result of a series of successive transformations undergone by that portion of the liquid which is lifted up during the ascension of the plate; which transformations ultimately reduce the thinnest part of that ascending liquid to so small a diameter, that it gives way, even without any further exterior exertion. The first experimenters, not being aware of this mode of acting, considered the separation of water from water as if it were similar to the rupture of a solid body; they made their calculations accordingly, and so doing, reduced to the lowest proportions that very strong molecular attraction which fixed my attention in Europe and Prof. Henry's in America.

The learned Professor has proved the magnitude of this molecular attraction by observations on soap-bubbles. I followed quite a different course, and arrived at more extensive results.

I constantly employed liquids placed in glass tubes, whose interior diameter measured from eight to ten millimetres (from three-tenths to four-tenths of an English inch). In similar circumstances, two distinct molecular forces are acting,—the attraction of water for water, or *cohesion*; and the attraction of water for glass, or *adhesion*. In my experiments, both cohesion and adhesion appeared very weak when the liquid was not deprived of that portion of air which it usually contains; and, on the contrary, proved very powerful when air was excluded.

In order to exhibit this power of attraction in airless liquids\*, I have made use of two different disjunctive forces; that of mechanical traction in my first two experiments, and that of repulsive caloric in the other.

My first experiment was made on sulphuric acid deprived of air by means of a very powerful air-pump†. In that case molecular attraction proved to be superior to the weight of a column of acid, whose height was 1250 millimetres (more than 4 English feet).

\* By *airless liquids*, I mean liquids deprived of air by one of the peculiar processes described in my memoir. In this sense, distilled water, although containing less air than common water, is far from being an airless liquid.

† This pump, constructed on a new plan, without either cock or valve, was described in 1841. It is recorded in the *Rapport du Jury et Documents de l'Exposition de l'Industrie Belge en 1841*, p. 161.

The second experiment was made on airless water, and the molecular force exceeded the weight of one atmosphere.

The third experiment proved the molecular attraction of airless water to be superior to the weight of three atmospheres, and exhibited very curious phænomena. The liquid had been placed in such circumstances as to be free from any pressure whatever; its temperature was carried to  $+135^{\circ}$  Centigrade (about  $+275$  Fahr.); and, nevertheless, it did not exhibit the least symptom of ebullition, but by still increasing heat, a part of it was suddenly vaporized with a kind of explosion.

A fourth experiment was tried by placing distilled water (not deprived of air) in a tube similar to that used in the third experiment; an external pressure equal to three atmospheres was applied to the liquid, which was then carried to the above-mentioned temperature of  $+135^{\circ}$  Centigrade: a calm, ordinary ebullition ensued, without any symptom of explosion.

In a fifth experiment, airless water was placed in a situation comparable to that of water in a steam-boiler working under low pressure. Continually increasing heat could not bring the airless liquid to ordinary ebullition; but the molecular attraction gave way from time to time by distinct explosions, becoming successively more and more violent, till a final one, blowing up the liquid mass and fracturing the instrument, put an end to the experiment.

My sixth experiment exhibited the molecular force in a still more striking form. A tube quite open at one end, half-filled with airless water, was heated over a lamp: no ebullition ensued, but a violent explosion took place, the water being at the same time suddenly projected out of the tube and converted into a cloud of vapour.

After a complete description of the experiments, a new theory of the ebullition of liquids is proposed as a consequence of the above-mentioned results, and of some peculiar considerations fully expounded in my memoir, and whereof it will be sufficient to mention here two of the most striking.

1. The molecules composing the surfaces of volatile bodies are very much inclined to assume a gaseous form, even whilst the internal molecules are kept together by a strong attraction.

2. Ordinary ebullition does not take place at once in the whole mass of a boiling liquid, the ebullitive motion being generated from some points of that portion of the boiler's internal surface which is near the source of heat; which points evolve a succession of large bubbles of vapour, tumultuously ascending through the liquid to its uppermost surface.

According to this new theory, *ebullition is a peculiar kind*

*of very rapid evaporation generated on those internal liquid surfaces which surround one or more bubbles of a gaseous fluid.*

I am, Gentlemen,

Your most humble Servant,

Ghent, March 2, 1846.

F. DONNY.

XLIX. *Experimental Researches in Electricity*.—*Nineteenth Series*. By MICHAEL FARADAY, Esq., D.C.L., F.R.S., Fullerian Prof. Chem. Royal Institution, Foreign Associate of the Acad. Sciences, Paris, Cor. Memb. Royal and Imp. Acad. of Sciences, Petersburg, Florence, Copenhagen, Berlin, Göttingen, Modena, Stockholm, &c. &c.\*

§ 26. *On the magnetization of light and the illumination of magnetic lines of force*†.

¶ i. *Action of magnets on light*.

2146. I HAVE long held an opinion, almost amounting to conviction, in common I believe with many other lovers of natural knowledge, that the various forms under which the forces of matter are made manifest have one common origin; or, in other words, are so directly related and mutually dependent, that they are convertible, as it were, one into another, and possess equivalents of power in their ac-

\* From the Philosophical Transactions for 1846, Part I., having been read November 20, 1845.

† The title of this paper has, I understand, led many to a misapprehension of its contents, and I therefore take the liberty of appending this explanatory note. Neither accepting nor rejecting the hypothesis of an æther, or the corpuscular, or any other view that may be entertained of the nature of light; and, as far as I can see, nothing being really known of a ray of light more than of a line of magnetic or electric force, or even of a line of gravitating force, except as it and they are manifest in and by substances; I believe that, in the experiments I describe in the paper, light has been magnetically affected, *i. e.* that that which is magnetic in the forces of matter has been affected, and in turn has affected that which is truly magnetic in the force of light: by the term magnetic I include here either of the peculiar exertions of the power of a magnet, whether it be that which is manifest in the magnetic or the diamagnetic class of bodies. The phrase "illumination of the lines of magnetic force" has been understood to imply that I had rendered them luminous. This was not within my thought. I intended to express that the line of magnetic force was illuminated as the earth is illuminated by the sun, or the spider's web illuminated by the astronomer's lamp. Employing a ray of light, we can tell, *by the eye*, the direction of the magnetic lines through a body; and by the alteration of the ray and its optical effect on the eye, can see the course of the lines just as we can see the course of a thread of glass, or any other transparent substance, rendered visible by the light: and this was what I meant by *illumination*, as the paper fully explains.—December 15, 1845 M. F.

tion\*. In modern times the proofs of their convertibility have been accumulated to a very considerable extent, and a commencement made of the determination of their equivalent forces.

2147. This strong persuasion extended to the powers of light, and led, on a former occasion, to many exertions, having for their object the discovery of the direct relation of light and electricity, and their mutual action in bodies subject jointly to their power†; but the results were negative and were afterwards confirmed, in that respect, by Wartmann‡.

2148. These ineffectual exertions, and many others which were never published, could not remove my strong persuasion derived from philosophical considerations; and, therefore, I recently resumed the inquiry by experiment in a most strict and searching manner, and have at last succeeded in *magnetizing and electrifying a ray of light, and in illuminating a magnetic line of force*. These results, without entering into the detail of many unproductive experiments, I will describe as briefly and clearly as I can.

2149. But before I proceed to them, I will define the meaning I connect with certain terms which I shall have occasion to use:—thus, by *line of magnetic force*, or *magnetic line of force*, or *magnetic curve*, I mean that exercise of magnetic force which is exerted in the lines usually called magnetic curves, and which equally exist as passing from or to magnetic poles, or forming concentric circles round an electric current. By *line of electric force*, I mean the force exerted in the lines joining two bodies, acting on each other according to the principles of static electric induction (1161, &c.), which may also be either in curved or straight lines. By a *diamagnetic*, I mean a body through which lines of magnetic force are passing, and which does not by their action assume the usual magnetic state of iron or loadstone.

2150. A ray of light issuing from an Argand lamp, was polarized in a horizontal plane by reflexion from a surface of glass, and the polarized ray passed through a Nichol's eye-piece revolving on a horizontal axis, so as to be easily examined by the latter. Between the polarizing mirror and the eye-piece, two powerful electro-magnetic poles were arranged, being either the poles of a horse-shoe magnet, or the contrary poles of two cylinder magnets; they were separated from each other about two inches in the direction of the line of the ray,

\* Experimental Researches, 57, 366, 376, 877, 961, 2071.

† Philosophical Transactions, 1834. Experimental Researches, 951-955.

‡ *Archives de l'Electricité*, ii, pp. 596-600.

and so placed, that, if on the same side of the polarized ray, it might pass near them; or, if on contrary sides, it might go between them, its direction being always parallel, or nearly so, to the magnetic lines of force (2149.). After that, any transparent substance placed between the two poles, would have passing through it, both the polarized ray and the magnetic lines of force at the same time and in the same direction.

2151. Sixteen years ago I published certain experiments made upon optical glass\*, and described the formation and general characters of one variety of heavy glass, which, from its materials, was called silicated borate of lead. It was this glass which first gave me the discovery of the relation between light and magnetism, and it has power to illustrate it in a degree beyond that of any other body; for the sake of perspicuity I will first describe the phænomena as presented by this substance.

2152. A piece of this glass, about two inches square and 0.5 of an inch thick, having flat and polished edges, was placed as a *diamagnetic* (2149.) between the poles (not as yet magnetized by the electric current), so that the polarized ray should pass through its length; the glass acted as air, water, or any other indifferent substance would do; and if the eye-piece were previously turned into such a position that the polarized ray was extinguished, or rather the image produced by it rendered invisible, then the introduction of this glass made no alteration in that respect. In this state of circumstances the force of the electro-magnet was developed, by sending an electric current through its coils, and immediately the image of the lamp-flame became visible, and continued so as long as the arrangement continued magnetic. On stopping the electric current, and so causing the magnetic force to cease, the light instantly disappeared; these phænomena could be renewed at pleasure, at any instant of time, and upon any occasion, showing a perfect dependence of cause and effect.

2153. The voltaic current which I used upon this occasion, was that of five pair of Grove's construction, and the electro-magnets were of such power that the poles would singly sustain a weight of from twenty-eight to fifty-six, or more, pounds.

\* Philosophical Transactions, 1830, p. 1. I cannot resist the occasion which is thus offered to me of mentioning the name of Mr. Anderson, who came to me as an assistant in the glass experiments, and has remained ever since in the Laboratory of the Royal Institution. He has assisted me in all the researches into which I have entered since that time, and to his care, steadiness, exactitude, and faithfulness in the performance of all that has been committed to his charge, I am much indebted.—M. F.

A person looking for the phænomenon for the first time would not be able to see it with a weak magnet.

2154. The character of the force thus impressed upon the diamagnetic is that of *rotation*; for when the image of the lamp-flame has thus been rendered visible, revolution of the eye-piece to the right or left, more or less, will cause its extinction; and the further motion of the eye-piece to the one side or other of this position will produce the reappearance of the light, and that with complementary tints, according as this further motion is to the right- or left-hand.

2155. When the pole nearest to the observer was a marked pole, *i. e.* the same as the north end of a magnetic needle, and the further pole was unmarked, the rotation of the ray was right-handed; for the eye-piece had to be turned to the right-hand, or clock fashion, to overtake the ray and restore the image to its first condition. When the poles were reversed, which was instantly done by changing the direction of the electric current, the rotation was changed also and became left-handed, the alteration being to an equal degree in extent as before. The direction was always the same for the same *line of magnetic force* (2149.).


2156. When the diamagnetic was placed in the numerous other positions, which can easily be conceived, about the magnetic poles, results were obtained more or less marked in extent, and very definite in character, but of which the phænomena just described may be considered as the chief example: they will be referred to, as far as is necessary, hereafter.

2157. The same phænomena were produced in the silicated borate of lead (2151.) by the action of a good ordinary steel horse-shoe magnet, no electric current being now used. The results were feeble, but still sufficient to show the perfect identity of action between electro-magnets and common magnets in this their power over light.

2158. Two magnetic poles were employed end-ways, *i. e.* the cores of the electro-magnets were hollow iron cylinders, and the ray of polarized light passed along their axes and through the diamagnetic placed between them: the effect was the same.


2159. One magnetic pole only was used, that being one end of a powerful cylinder electro-magnet. When the heavy glass was beyond the magnet, being close to it but between the magnet and the polarizing reflector, the rotation was in one direction, dependent on the nature of the pole; when the diamagnetic was on the near side, being close to it but between it and the eye, the rotation for the same pole was in the con-

trary direction to what it was before; and when the magnetic pole was changed, both these directions were changed with it. When the heavy glass was placed in a corresponding position to the pole, but above or below it, so that the *magnetic curves* were no longer passing through the glass parallel to the ray of polarized light, but rather perpendicular to it, then no effect was produced. These particularities may be understood by reference to fig. 1, where *a* and *b* represent the first positions of the diamagnetic, and *c* and *d* the latter positions, the course of the ray being marked by the dotted line. If also the glass were placed directly at the end of the magnet, then no effect was produced on a ray passing in the direction here described, though it is evident, from what has been already said (2155.), that a ray passing *parallel* to the magnetic lines through the glass so placed, would have been affected by it.



2160. Magnetic lines, then, in passing through silicated borate of lead, and a great number of other substances (2173.), cause these bodies to act upon a polarized ray of light when the lines are parallel to the ray, or in proportion as they are parallel to it: if they are perpendicular to the ray, they have no action upon it. They give the diamagnetic the power of rotating the ray; and the *law* of this action on light is, that if a magnetic line of force be *going from* a north pole, or *coming from* a south pole, along the path of a polarized ray coming to the observer, it will rotate that ray to the right-hand; or, that if such a line of force be coming from a north pole, or going from a south pole, it will rotate such a ray to the left-hand.

2161. If a cork or a cylinder of glass, representing the diamagnetic, be marked at its ends with the letters N and S, to represent the poles of a magnet, the line joining these letters may be considered as a magnetic line of force; and further, if a line be traced round the cylinder with arrow heads on it to represent direction, as in the figure, such a simple model, held up before the eye, will express the whole of the law, and give every position and consequence of direction resulting from it. If a watch be considered as the diamagnetic, the north pole of a magnet being imagined





against the face, and a south pole against the back, then the motion of the hands will indicate the direction of rotation which a ray of light undergoes by magnetization.

2162. I will now proceed to the different circumstances which affect, limit, and define the extent and nature of this new power of action on light.

2163. In the first place, the rotation appears to be in proportion to the extent of the diamagnetic through which the ray and the magnetic lines pass. I preserved the strength of the magnet and the interval between its poles constant, and then interposed different pieces of the same heavy glass (2151.) between the poles. The greater the extent of the diamagnetic in the line of the ray, whether in one, two, or three pieces, the greater was the rotation of the ray; and, as far as I could judge by these first experiments, the amount of rotation was exactly proportionate to the extent of diamagnetic through which the ray passed. No addition or diminution of the heavy glass on the *side* of the course of the ray made any difference in the effect of that part through which the ray passed.

2164. The power of rotating the ray of light *increased* with the intensity of the magnetic lines of force. This general effect is very easily ascertained by the use of electro-magnets; and within such range of power as I have employed, it appears to be directly proportionate to the intensity of the magnetic force.

2165. Other bodies, besides the heavy glass, possess the same power of becoming, under the influence of magnetic force, active on light (2173.). When these bodies possess a rotative power of their own, as is the case with oil of turpentine, sugar, tartaric acid, tartrates, &c., the effect of the magnetic force is to add to, or subtract from, their specific force, according as the natural rotation and that induced by the magnetism is right- or left-handed (2231.).

2166. I could not perceive that this power was affected by any degree of motion which I was able to communicate to the diamagnetic, whilst jointly subject to the action of the magnetism and the light.

2167. The interposition of copper, lead, tin, silver, and other ordinary non-magnetic bodies in the course of the magnetic curves, either between the pole and the diamagnetic, or in other positions, produced no effect either in kind or degree upon the phenomena.

2168. Iron frequently affected the results in a very considerable degree; but it always appeared to be, either by altering the direction of the magnetic lines, or disposing within

itself of their force. Thus, when the two contrary poles were on one side of the polarized ray (2150.), and the heavy glass in its best position between them and in the ray (2152.), the bringing of a large piece of iron near to the glass on the other side of the ray, caused the power of the diamagnetic to fall. This was because certain lines of magnetic force, which at first passed through the glass parallel to the ray, now crossed the glass and the ray; the iron giving two contrary poles opposite the poles of the magnet, and thus determining a new course for a certain portion of the magnetic power, and that across the polarized ray.

2169. Or, if the iron, instead of being applied on the opposite side of the glass, were applied on the same side with the magnet, either near it or in contact with it, then, again, the power of the diamagnetic fell, simply because the power of the magnet was diverted from it into a new direction. These effects depend much of course on the intensity and power of the magnet, and on the size and softness of the iron.

2170. The electro-helices (2190.) without the iron cores were very feeble in power, and indeed hardly sensible in their effect. With the iron cores they were powerful, though no more electricity was then passing through the coils than before (1071.). This shows, in a very simple manner, that the phænomena exhibited by light under these circumstances, is directly connected with the magnetic form of force supplied by the arrangement. Another effect which occurred illustrated the same point. When the contact at the voltaic battery is made, and the current sent round the electro-magnet, the image produced by the rotation of the polarized ray does not rise up to its full lustre immediately, but increases for a couple of seconds, gradually acquiring its greatest intensity; on breaking the contact, it sinks instantly and disappears apparently at once. The gradual rise in brightness is due to the *time* which the iron core of the magnet requires to evolve all that magnetic power which the electric current can develop in it; and as the magnetism rises in intensity, so does its effect on the light increase in power; hence the progressive condition of the rotation.

2171. I cannot as yet find that the heavy glass (2151.), when in this state, *i. e.* with magnetic lines of force passing through it, exhibits any increased degree, or has any specific magneto-inductive action of the recognized kind. I have placed it in large quantities, and in different positions, between magnets and magnetic needles, having at the time very delicate means of appreciating any difference between it and air, but could find none.

2172. Using water, alcohol, mercury, and other fluids contained in very large delicate thermometer-shaped vessels, I could not discover that any difference in volume occurred when the magnetic curves passed through them.

2173. It is time that I should pass to a consideration of this power of magnetism over light as exercised, not only in the silicated borate of lead (2151.), but in many other substances; and here we perceive, in the first place, that if all transparent bodies possess the power of exhibiting the action, they have it in very different degrees, and that up to this time there are some that have not shown it at all.

2174. Next, we may observe, that bodies that are exceedingly different to each other in chemical, physical, and mechanical properties, develop this effect; for solids and liquids, acids, alkalies, oils, water, alcohol, æther, all possess the power.

2175. And lastly, we may observe, that in all of them, though the degree of action may differ, still it is always the same in kind, being a rotative power over the ray of light; and further, the direction of the rotation is, in every case, independent of the nature or state of the substance, and dependent upon the direction of the magnetic line of force, according to the law before laid down (2160.).

2176. Amongst the substances in which this power of action is found, I have already distinguished the *silico-borate of lead* (2151.) as eminently fitted for the purpose of exhibiting the phænomena. I regret that it should be the best, since it is not likely to be in the possession of many, and few will be induced to take the trouble of preparing it. If made, it should be well-annealed, for otherwise the pieces will have considerable power of depolarizing light, and then the particular phænomena under consideration are much less strikingly observed. The *borate of lead*, however, is a substance much more fusible, softening at the heat of boiling oil, and therefore far more easily prepared in the form of glass plates and annealed; and it possesses as much magneto-rotative power over light as the silico-borate itself. *Flint-glass* exhibits the property, but in a less degree than the substances above. *Crown-glass* shows it, but in a still smaller degree.

2177. Whilst employing crystalline bodies as diamagnetics, I generally gave them that position in which they did not affect the polarized ray, and then induced the magnetic curves through them. As a class, they seemed to resist the assumption of the rotating state. *Rock-salt* and *fluor-spar* gave evidence of the power in a slight degree; and I think that a crystal of alum did the same, but its ray length in the transparent

part was so small that I could not ascertain the fact decisively. Two specimens of transparent fluor, lent me by Mr. Tennant, gave the effect.

2178. Rock-crystal, four inches across, gave no indications of action on the ray, neither did smaller crystals, nor cubes about three-fourths of an inch in the side, which were so cut as to have two of their faces perpendicular to the axis of the crystal (1692, 1693.), though they were examined in every direction.

2179. *Iceland spar* exhibited no signs of effect, either in the form of rhomboids, or of cubes like those just described (1695.).

2180. *Sulphate of baryta, sulphate of lime, and carbonate of soda*, were also without action on the light.

2181. A piece of fine clear *ice* gave me no effect. I cannot however say there is none, for the effect of water in the same mass would be very small, and the irregularity of the flattened surface from the fusion of the ice and flow of water, made the observation very difficult.

2182. With some degree of curiosity and hope, I put gold-leaf into the magnetic lines, but could perceive no effect. Considering the extremely small dimensions of the length of the path of the polarized ray in it, any positive result was hardly to be expected.

2183. In experiments with liquids, a very good method of observing the effect, is to inclose them in bottles from  $1\frac{1}{2}$  to 3 or 4 inches in diameter, placing these in succession between the magnetic poles (2150.), and bringing the analysing eye-piece so near to the bottle, that, by adjustment of the latter, its cylindrical form may cause a diffuse but useful image of the lamp-flame to be seen through it: the light of this image is easily distinguished from that which passes by irregular refraction through the striæ and deformations of the glass, and the phænomena being looked for in this light are easily seen.

2184. Water, alcohol, and æther, all show the effect; water most, alcohol less, and æther the least. All the fixed oils which I have tried, including almond, castor, olive, poppy, linseed, sperm, elaine from hog's lard, and distilled resin oil, produce it. The essential oils of turpentine, bitter almonds, spike lavender, lavender, jessamine, cloves, and laurel, produce it. Also naphtha of various kinds, melted spermaceti, fused sulphur, chloride of sulphur, chloride of arsenic, and every other liquid substance which I had at hand and could submit in sufficient bulk to experiment.

2185. Of aqueous solutions I tried 150 or more, including the soluble acids, alkalies and salts, with sugar, gum, &c., the

list of which would be too long to give here, since the great conclusion was, that the exceeding diversity of substance caused no exception to the general result, for all the bodies showed the property. It is indeed more than probable, that in all these cases the water and not the other substance present was the ruling matter. The same general result was obtained with alcoholic solutions.

2186. Proceeding from liquids to air and gaseous bodies, I have here to state that, as yet, I have not been able to detect the exercise of this power in any one of the substances in this class. I have tried the experiment with bottles 4 inches in diameter, and the following gases: oxygen, nitrogen, hydrogen, nitrous oxide, olefiant gas, sulphurous acid, muriatic acid, carbonic acid, carbonic oxide, ammonia, sulphuretted hydrogen, and bromine vapour, at ordinary temperatures; but they all gave negative results. With air, the trial has been carried, by another form of apparatus, to a much higher degree, but still ineffectually (2212.).

2187. Before dismissing the consideration of the substances which exhibited this power, and in reference to those in which it was superinduced upon bodies possessing, naturally, rotative force (2165. 2231.), I may record, that the following are the substances submitted to experiment: castor oil, resin oil, oil of spike lavender, of laúrel, Canada balsam, alcoholic solution of camphor, alcoholic solution of camphor and corrosive sublimate, aqueous solutions of sugar, tartaric acid, tartrate of soda, tartrate of potassa and antimony, tartaric and boracic acid, and sulphate of nickel, which rotated to the right-hand; copaiba balsam, which rotated the ray to the left-hand; and two specimens of camphine or oil of turpentine, in one of which the rotation was to the right-hand, and in the other to the left. In all these cases, as already said (2165.), the superinduced magnetic rotation was according to the general law (2160.), and without reference to the previous power of the body.

2188. Camphor being melted in a tube about an inch in diameter, exhibited high natural rotative force, but I could not discover that the magnetic curves induced additional force in it. It may be, however, that the shortness of the ray length and the quantity of coloured light left, even when the eye-piece was adjusted to the most favourable position for darkening the image produced by the naturally rotated ray, rendered the small magneto-power of the camphor insensible.

#### ¶ ii. *Action of electric currents on light.*

2189. From a consideration of the nature and position of the lines of magnetic and electric force, and the relation of a magnet to a current of electricity, it appeared almost certain

that an electric current would give the same result of action on light as a magnet; and, in the helix, would supply a form of apparatus in which great lengths of diamagnetics, and especially of such bodies as appeared to be but little affected between the poles of the magnet, might be submitted to examination and their effect exalted: this expectation was, by experiment, realized.

2190. Helices of copper wire were employed, three of which I will refer to. The first, or *long helix*, was 0·4 of an inch internal diameter; the wire was 0·03 of an inch in diameter, and having gone round the axis from one end of the helix to the other, then returned in the same manner, forming a coil sixty-five inches long, double in its whole extent, and containing 1240 feet of wire.

2191. The second, or *medium helix*, is nineteen inches long, 1·87 inch internal diameter, and three inches external diameter. The wire is 0·2 of an inch in diameter, and eighty feet in length, being disposed in the coil as two concentric spirals. The electric current, in passing through it, is not divided, but traverses the whole length of the wire.

2192. The third, or *Woolwich helix*, was made under my instruction for the use of Lieut.-Colonel Sabine's establishment at Woolwich. It is 26·5 inches long, 2·5 inches internal diameter, and 4·75 inches external diameter. The wire is 0·17 of an inch in diameter, and 501 feet in length. It is disposed in the coil in four concentric spirals connected end to end, so that the whole of the electric current employed passes through all the wire.

2193. The long helix (2190.) acted very feebly on a magnetic needle placed at a little distance from it; the medium helix (2191.) acted more powerfully, and the Woolwich helix (2192.) very strongly; the same battery of ten pairs of Grove's plate being employed in all cases.

2194. Solid bodies were easily subjected to the action of these electro-helices, being for that purpose merely cut into the form of bars or prisms with flat and polished ends, and then introduced as cores into the helices. For the purpose of submitting liquid bodies to the same action, tubes of glass were provided, furnished at the ends with caps; the cylindrical part of the cap was brass, and had a tubular aperture for the introduction of the liquids, but the end was a flat glass plate. When the tube was intended to contain aqueous fluids, the plates were attached to the caps, and the caps to the tube by Canada balsam; when the tube had to contain alcohol, æther or essential oils, a thick mixture of powdered gum with a little water was employed as the cement.

2195. The general effect produced by this form of appa-

ratus may be stated as follows:—The tube within the long helix (2190.) was filled with distilled water and placed in the line of the polarized ray, so that by examination through the eye-piece (2150.), the image of the lamp-flame produced by the ray could be seen through it. Then the eye-piece was turned until the image of the flame disappeared, and, afterwards, the current of ten pairs of plates sent through the helix; instantly the image of the flame reappeared, and continued as long as the electric current was passing through the helix; on stopping the current the image disappeared. The light did not rise up gradually, as in the case of electro-magnets (2170.), but instantly. These results could be produced at pleasure. In this experiment we may, I think, justly say that a ray of light is electrified and the electric forces illuminated.

2196. The phænomena may be made more striking, by the adjustment of a lens of long focus between the tube and the polarizing mirror, or one of short focus between the tube and the eye; and where the helix, or the battery, or the substance experimented with, is feeble in power, such means offer assistance in working out the effects: but, after a little experience, they are easily dispensed with, and are only useful as accessories in doubtful cases.

2197. In cases where the effect is feeble, it is more easily perceived if the Nichol eye-piece be adjusted, not to the perfect extinction of the ray, but a little short of or beyond that position; so that the image of the flame may be but just visible. Then, on the exertion of the power of the electric current, the light is either increased in intensity, or else diminished, or extinguished, or even re-illuminated on the other side of the dark condition; and this change is more easily perceived than if the eye began to observe from a state of utter darkness. Such a mode of observing also assists in demonstrating the rotatory character of the action on light; for, if the light be made visible beforehand by the motion of the eye-piece in one direction, and the power of the current be to *increase* that light, an instant only suffices, after stopping the current, to move the eye-piece in the other direction until the light is apparent as at first, and then the power of the current will be to *diminish* it; the tints of the lights being affected also at the same time.

2198. When the current was sent round the helix in one direction, the rotation induced upon the ray of light was one way; and when the current was changed to the contrary direction, the rotation was the other way. In order to express the direction, I will assume, as is usually done, that the cur-

rent passes from the zinc through the acid to the platinum in the same cell (663. 667. 1627.): if such a current pass under the ray towards the right, upwards on its right side, and over the ray towards the left, it will give left-handed rotation to it; or, if the current pass over the ray to the right, down on the right side, and under it towards the left, it will induce it to rotate to the right-hand.

2199. The LAW, therefore, by which an electric current acts on a ray of light is easily expressed. When an electric current passes round a ray of polarized light in a plane perpendicular to the ray, it causes the ray to revolve on its axis, as long as it is under the influence of the current, in the *same direction* as that in which the current is passing.

2200. The simplicity of this law, and its identity with that given before, as expressing the action of magnetism on light (2160.), is very beautiful. A model is not wanted to assist the memory; but if that already described (2161.) be looked at, the line round it will express at the same time the direction both of the current and the rotation. It will indeed do much more; for if the cylinder be considered as a piece of iron, and not a piece of glass or other diamagnetic, placed between the two poles N and S, then the line round it will represent the direction of the currents, which, according to Ampère's theory, are moving round its particles; or if it be considered as a core of iron (in place of a core of water), having an electric current running round it in the direction of the line, it will also represent such a magnet as would be formed if it were placed between the poles whose marks are affixed to its ends.

2201. I will now notice certain points respecting the degree of this action under different circumstances. By using a tube of water (2194.) as long as the helix, but placing it so that more or less of the tube projected at either end of the helix, I was able, in some degree, to ascertain the effect of length of the diamagnetic, the force of the helix and current remaining the same. The greater the column of water subjected to the action of the helix, the greater was the rotation of the polarized ray; and the amount of rotation seemed to be directly proportionate to the length of fluid round which the electric current passed.

2202. A short tube of water, or a piece of heavy glass, being placed in the axis of the Woolwich helix (2192.), seemed to produce equal effect on the ray of light, whether it were in the middle of the helix or at either end; provided it was always within the helix and in the line of the axis. From this it would appear that every part of the helix has the same effect;



and, that by using long helices, substances may be submitted to this kind of examination which could not be placed in sufficient length between the poles of magnets (2150.).

2203. A tube of water as long as the Woolwich helix (2192.), but only 0.4 of an inch in diameter, was placed in the helix parallel to the axis, but sometimes in the axis and sometimes near the side. No apparent difference was produced in these different situations; and I am inclined to believe (without being quite sure) that the action on the ray is the same, wherever the tube is placed, within the helix, in relation to the axis. The same result was obtained when a larger tube of water was looked through, whether the ray passed through the axis of the helix and tube, or near the side.

2204. If bodies be introduced into the helix possessing, naturally, rotating force, then the rotating power given by the electric current is superinduced upon them, exactly as in the cases already described of magnetic action (2165. 2187.).

2205. A helix, twenty inches long and 0.3 of an inch in diameter, was made of uncovered copper wire, 0.05 of an inch in diameter, in close spirals. This was placed in a large tube of water, so that the fluid, both in the inside and at the outside of the helix, could be examined by the polarized ray. When the current was sent *through* the helix, the water within it received rotating power; but no trace of such an action on the light was seen on the outside of the helix, even in the line most close to the uncovered wire.

2206. The water was inclosed in brass and copper tubes, but this alteration caused not the slightest change in the effect.

2207. The water in the brass tube was put into an *iron* tube, much longer than either the Woolwich helix or the brass tube, and quite one-eighth of an inch thick in the side; yet when placed in the Woolwich helix (2192.), the water rotated the ray of light apparently as well as before.

2208. An iron bar, one inch square and longer than the helix, was put into the helix, and the small water-tube (2203.) upon it. The water exerted as much action on the light as before.

2209. Three iron tubes, each twenty-seven inches long and one-eighth of an inch in thickness in the side, were selected of such diameters as to pass easily one into the other, and the whole into the Woolwich helix (2192.). The smaller one was supplied with glass ends and filled with water; and being placed in the axis of the Woolwich helix, had a certain amount of rotating power over the polarized ray. The

second tube was then placed over this, so that there was now a thickness of iron equal to two-eighths of an inch between the water and the helix; the water had *more* power of rotation than before. On placing the third tube of iron over the two former, the power of the water *fell*, but was still very considerable. These results are complicated, being dependent on the new condition which the character of iron gives to its action on the forces. Up to a certain amount, by increasing the development of magnetic forces, the helix and core, *as a whole*, produce increased action on the water; but on the addition of more iron and the disposal of the forces through it, their action is removed in part from the water and the rotation is lessened.

2210. Pieces of heavy glass (2151.), placed in iron tubes in the helices, produced similar effects.

2211. The bodies which were submitted to the action of an electric current in a helix, in the manner already described, were as follows:—Heavy glass (2151. 2176.), water, solution of sulphate of soda, solution of tartaric acid, alcohol, æther, and oil of turpentine; all of which were affected, and acted on light exactly in the manner described in relation to magnetic action (2173.).

2212. I submitted *air* to the influence of these helices carefully and anxiously, but could not discover any trace of action on the polarized ray of light. I put the long helix (2190.) into the other two (2191. 2192.), and combined them all into one consistent series, so as to accumulate power, but could not observe any effect of them on light passing through air.

2213. In the use of helices, it is necessary to be aware of one effect, which might otherwise cause confusion and trouble. At first, the wire of the long helix (2190.) was wound directly upon the thin glass tube which served to contain the fluid. When the electric current passed through the helix it raised the temperature of the metal, and that gradually raised the temperature of the glass and the film of water in contact with it, and so the cylinder of water, warmer at its surface than its axis, acted as a lens, gathering and sending rays of light to the eye, and continuing to act for a time after the current was stopped. By separating the tube of water from the helix, and by other precautions, this source of confusion is easily avoided.

2214. Another point of which the experimenter should be aware, is the difficulty, and almost impossibility, of obtaining a piece of glass which, especially after it is cut, does not depolarize light. When it does depolarize, difference of position makes an immense difference in the appearance. By always referring to the parts that do not depolarize, as the black

cross, for instance, and by bringing the eye as near as may be to the glass, this difficulty is more or less overcome.

2215. For the sake of supplying a general indication of the amount of this induced rotating force in two or three bodies, and without any pretence of offering correct numbers, I will give, generally, the result of a few attempts to measure the force, and compare it with the natural power of a specimen of oil of turpentine. A very powerful electro-magnet was employed, with a *constant* distance between its poles of  $2\frac{1}{2}$  inches. In this space was placed different substances; the amount of rotation of the eye-piece observed several times and the average taken, as expressing the rotation for the ray length of substance used. But as the substances were of different dimensions, the ray lengths were, by calculation, corrected to one standard length, upon the assumption that the power was proportionate to this length (2163.). The oil of turpentine was of course observed in its natural state, *i. e.* without magnetic action. Making water 1, the numbers were as follows:—

Oil of turpentine . . . . .	11·8
Heavy glass (2151.) . . . . .	6·0
Flint-glass . . . . .	2·8
Rock-salt . . . . .	2·2
Water . . . . .	1·0
Alcohol . . . . .	less than water.
Æther . . . . .	less than alcohol.

2216. In relation to the action of magnetic and electric forces on light, I consider, that to know the conditions under which there is no apparent action, is to add to our knowledge of their mutual relations; and will, therefore, very briefly state how I have lately combined these forces, obtaining no apparent result (955.).

2217. Heavy glass, flint-glass, rock-crystal, Iceland spar, oil of turpentine, and air, had a polarized ray passed through them; and, at the same time, lines of electro-static tension (2149.) were, by means of coatings, the Leyden jar, and the electric machine, directed across the bodies, parallel to the polarized ray, and perpendicular to it, both in and across the plane of polarization; but without any visible effect. The tension of a rapidly recurring, induced secondary current, was also directed upon the same bodies and upon water (as an electrolyte), but with the same negative result.

2218. A polarized ray, powerful magnetic lines of force, and the electric lines of force (2149.) just described, were combined in various directions in their action on heavy glass

(2151. 2176.), but with no other result than that due to the mutual action of the magnetic lines of light, already described in this paper.

2219. A polarized ray and electric currents were combined in every possible way in electrolytes (951-954). The substances used were distilled water, solution of sugar, dilute sulphuric acid, solution of sulphate of soda, using platinum electrodes; and solution of sulphate of copper, using copper electrodes; the current was sent along the ray, and perpendicular to it in two directions at right angles with each other; the ray was made to rotate, by altering the position of the polarizing mirror, that the plane of polarization might be varied; the current was used as a continuous current, as a rapidly intermitting current, and as a rapidly alternating double current of induction; but in no case was any trace of action perceived.

2220. Lastly, a ray of polarized light, electric currents, and magnetic lines of force, were directed in every possible way through dilute sulphuric acid and solution of sulphate of soda, but still with negative results, except in those positions where the phænomena already described were produced. In one arrangement, the current passed in the direction of radii from a central to a circumferential electrode, the contrary magnetic poles being placed above and below; and the arrangements were so good, that when the electric current was passing, the fluid rapidly rotated; but a polarized ray sent horizontally across this arrangement was not at all affected. Also, when the ray was sent vertically through it, and the eyepiece moved to correspond to the rotation impressed upon the ray in this position by the magnetic curves alone, the superinduction of the passage of the electric current made not the least difference in the effect upon the ray.

### ¶ iii. *General considerations.*

2221. Thus is established, I think for the first time \*, a

\* I say, for the first time, because I do not think that the experiments of Morrichini on the production of magnetism by the rays at the violet end of the spectrum prove any such relation. When in Rome with Sir H. Davy in the month of May 1814, I spent several hours at the house of Morrichini, working with his apparatus and under his directions, but could not succeed in magnetising a needle. I have no confidence in the effect as a *direct* result of the action of the sun's rays; but think, that when it has occurred it has been secondary, incidental, and perhaps even accidental; a result that might well happen with a needle that was preserved during the whole experiment in a north and south position.

January 2, 1846.—I should not have written "for the first time" as above, if I had remembered Mr. Christie's experiments and papers on the Influence of the Solar Rays on Magnets, communicated in the Philosophical Transactions for 1826, p. 219, and 1828, p. 379.—M. F.

true, direct relation and dependence between light and the magnetic and electric forces; and thus a great addition made to the facts and considerations which tend to prove that all natural forces are tied together, and have one common origin (2146.). It is, no doubt, difficult in the present state of our knowledge to express our expectation in exact terms; and, though I have said that another of the powers of nature is, in these experiments, directly related to the rest, I ought, perhaps, rather to say that another form of the great power is distinctly and directly related to the other forms; or that the great power manifested by particular phænomena in particular forms, is here further identified and recognised, by the direct relation of its form of light to its forms of electricity and magnetism.

2222. The relation existing between *polarized* light and magnetism and electricity, is even more interesting than if it had been shown to exist with common light only. It cannot but extend to common light; and, as it belongs to light made, in a certain respect, more precise in its character and properties by polarization, it collates and connects it with these powers, in that duality of character which they possess, and yields an opening, which before was wanting to us, for the appliance of these powers to the investigation of the nature of this and other radiant agencies.

2223. Referring to the conventional distinction before made (2149.), it may be again stated, that it is the magnetic lines of force *only* which are effectual on the rays of light, and they *only* (in appearance) when parallel to the ray of light, or as they tend to parallelism with it. As, in reference to matter not magnetic after the manner of iron, the phænomena of electric induction and electrolysation show a vast superiority in the energy with which electric forces can act as compared to magnetic forces, so here, in another direction and in the peculiar and correspondent effects which belong to magnetic forces, they are shown, in turn, to possess great superiority, and to have their full equivalent of action on the same kind of matter.

2224. The magnetic forces do not act on the ray of light directly and without the intervention of matter, but through the mediation of the substance in which they and the ray have a simultaneous existence; the substances and the forces giving to and receiving from each other the power of acting on the light. This is shown by the non-action of a vacuum, of air or gases; and it is also further shown by the special degree in which different matters possess the property. That magnetic force acts upon the ray of light always with the same

character of manner and in the same direction, independent of the different varieties of substance, or their states of solid or liquid, or their specific rotative force (2232.), shows that the magnetic force and the light have a direct relation: but that substances are necessary, and that these act in different degrees, shows that the magnetism and the light act on each other through the intervention of the matter.

2225. Recognizing or perceiving *matter* only by its powers, and knowing nothing of any imaginary nucleus, abstract from the idea of these powers, the phænomena described in this paper much strengthen my inclination to trust in the views I have on a former occasion advanced in reference to its nature\*.

2226. It cannot be doubted that the magnetic forces act upon and affect the internal constitution of the diamagnetic, just as freely in the dark as when a ray of light is passing through it; though the phænomena produced by light seem, as yet, to present the only means of observing this constitution and the change. Further, any such change as this must belong to opaque bodies, such as wood, stone, and metal; for as diamagnetics, there is no distinction between them and those which are transparent. The degree of transparency can at the utmost, in this respect, only make a distinction between the individuals of a class.

2227. If the magnetic forces had made these bodies magnets, we could, by light, have examined a transparent magnet; and that would have been a great help to our investigation of the forces of matter. But it does not make them magnets (2171.), and therefore the molecular condition of these bodies, when in the state described, must be specifically distinct from that of magnetized iron, or other such matter, and must be a *new magnetic condition*; and as the condition is a state of tension (manifested by its instantaneous return to the normal state when the magnetic induction is removed), so the *force* which the matter in this state possesses and its mode of action, must be to us a *new magnetic force* or *mode of action* of matter.

2228. For it is impossible, I think, to observe and see the action of magnetic forces, rising in intensity, upon a piece of heavy glass or a tube of water, without also perceiving that the latter acquire properties which are not only *new* to the substance, but are also in subjection to very definite and precise laws (2160. 2199.), and are equivalent in proportion to the magnetic forces producing them.

2229. Perhaps this state is a state of *electric tension tending*

\* A speculation, &c. Philosophical Magazine, 1844, vol. xxiv. p. 136.

to a current; as in magnets, according to Ampère's theory, the state is a state of *current*. When a core of iron is put into a helix, every thing leads us to believe that currents of electricity are produced within it, which rotate or move in a plane perpendicular to the axis of the helix. If a diamagnetic be placed in the same position, it acquires power to make light rotate in the same plane. The state it has received is a state of tension, but it has not passed on into currents, though the acting force and every other circumstance and condition are the same as those which do produce currents in iron, nickel, cobalt, and such other matters as are fitted to receive them. Hence the idea that there exists in diamagnetics, under such circumstances, a tendency to currents, is consistent with all the phænomena as yet described, and is further strengthened by the fact, that, leaving the loadstone or the electric current, which by inductive action is rendering a piece of iron, nickel, or cobalt magnetic, perfectly unchanged, a mere change of temperature will take from these bodies their extra power, and make them pass into the common class of diamagnetics.

---

2230. The present is, I believe, the first time that the molecular condition of a body, required to produce the circular polarization of light, has been artificially given; and it is therefore very interesting to consider this known state and condition of the body, comparing it with the relatively unknown state of those which possess the power naturally: especially as some of the latter rotate to the right-hand and others to the left; and, as in the cases of quartz and oil of turpentine, the same body chemically speaking, being in the latter instance a liquid with particles free to move, presents different specimens, some rotating one way and some the other.

2231. At first one would be inclined to conclude that the natural state and the state conferred by magnetic and electric forces must be the same, since the effect is the same; but on further consideration it seems very difficult to come to such a conclusion. Oil of turpentine will rotate a ray of light, the power depending upon its particles and not upon the arrangement of the mass. Whichever way a ray of polarized light passes through this fluid, it is rotated in the same manner; and rays passing in every possible direction through it *simultaneously* are all rotated with equal force and according to one common law of direction; *i. e.* either all right-handed or else all to the left. Not so with the rotation superinduced on the *same* oil of turpentine by the magnetic or electric forces: it exists only in one direction, *i. e.* in a plane perpendicular

to the magnetic line; and being limited to this plane, it can be changed in direction by a reversal of the direction of the inducing force. The direction of the rotation produced by the natural state is connected invariably with the direction of the ray of light; but the power to produce it appears to be possessed in every direction and at all times by the particles of the fluid: the direction of the rotation produced by the induced condition is connected invariably with the direction of the magnetic line or the electric current, and the condition is possessed by the particles of matter, but strictly limited by the line or the current, changing and disappearing with it.

2332. Let  $m$ , in fig. 3, represent a glass cell filled with oil of turpentine, possessing naturally the power of producing right-hand rotation, and  $a b$  a polarized ray of light. If the ray proceed from  $a$  to  $b$ , and the eye be placed at  $b$ , the rotation will be right-handed, or according to the direction expressed by the arrow-heads on the circle  $c$ ; if the ray proceed from  $b$  to  $a$ , and the eye be placed at  $a$ , the rotation will still be right-handed to the observer, *i. e.* according to the direction indicated on the circle  $d$ . Let now an electric current pass round the oil of turpentine in the direction indicated on the circle  $c$ , or magnetic poles be placed so as to produce the same effect (2155); the particles will acquire a further rotative force (which no motion amongst themselves will disturb), and a ray coming from  $a$  to  $b$  will be seen by an eye placed at  $b$




Fig. 3.

to rotate to the right-hand more than before, or in the direction on the circle  $c$ ; but pass a ray from  $b$  to  $a$ , and observe with the eye at  $a$ , and the phenomenon is no longer the same as before; for instead of the new rotation being according to the direction indicated on the circle  $d$ , it will be in the contrary direction, or to the observer's left-hand (2199). In fact the induced rotation will be added to the natural rotation as respects a ray passing from  $a$  to  $b$ , but it will be subtracted from the natural rotation as regards the ray passing from  $b$  to  $a$ . Hence the particles of this fluid which rotate by virtue of their natural force, and those which rotate by virtue of the induced force, cannot be in the same condition.

2233. As respects the power of the oil of turpentine to rotate a ray in whatever direction it is passing through the liquid, it may well be, that though all the particles possess



the power of rotating the light, only those whose planes of rotation are more or less perpendicular to the ray affect it; and that it is the resultant or sum of forces in any one direction which is active in producing rotation. But even then a striking difference remains, because the resultant in the same plane is not absolute in direction, but relative to the course of the ray, being in the one case as the circle *c*, and in the other as the circle *d*, fig. 3; whereas the resultant of the magnetic or electric induction is absolute, and not changing with the course of the ray, being always either as expressed by *c* or else as indicated by *d*.

2234. All these differences, however, will doubtless disappear or come into harmony as these investigations are extended; and their very existence opens so many paths, by which we may pursue our inquiries, more and more deeply, into the powers and constitution of matter.

2235. Bodies having rotating power of themselves, do not seem by that to have a greater or a less tendency to assume a further degree of the same force under the influence of magnetic or electric power.

2236. Were it not for these and other differences, we might see an analogy between these bodies, which possess at all times the rotating power, as a specimen of quartz which rotates only in one plane, and those to which the power is given by the induction of other forces, as a prism of heavy glass in a helix, on the one hand; and, on the other, a natural magnet and a helix through which the current is passing. The natural condition of the magnet and quartz, and the constrained condition of the helix and heavy glass, form the link of the analogy in one direction; whilst the supposition of currents existing in the magnet and helix, and only a tendency or tension to currents existing in the quartz and heavy glass, supplies the link in the transverse direction.

2237. As to those bodies which seem as yet to give no indication of the power over light, and therefore none of the assumption of the new magnetic conditions; these may be divided into two classes, the one including air, gases and vapours, and the other rock crystal, Iceland spar, and certain other crystalline bodies. As regards the latter class, I shall give, in the next series of these researches, proofs drawn from phenomena of an entirely different kind, that they do acquire the new magnetic condition; and these being so disposed of for the moment, I am inclined to believe that even air and gases have the power to assume the peculiar state, and even to affect light, but in a degree so small that as yet it has not been made sensible. Still the gaseous state is such a remarkable condi-

tion of matter, that we ought not too hastily to assume that the substances which, in the solid and liquid state, possess properties even general in character, always carry these into their gaseous condition.

2238. Rock-salt, fluor-spar, and, I think, alum, affect the ray of light; the other crystals experimented with did not; these are equiaxed and singly refracting, the others are unequiaxed and doubly refracting. Perhaps these instances, with that of the rotation of quartz, may even now indicate a relation between magnetism, electricity, and the crystallizing forces of matter.

2239. All bodies are affected by helices as by magnets, and according to laws which show that the causes of the action are identical as well as the effects. This result supplies another fine proof in favour of the identity of helices and magnets, according to the views of Ampère.

2240. The theory of static induction which I formerly ventured to set forth (1161, &c.), and which depends upon the action of the contiguous particles of the dielectric intervening between the inductric and the inducteous bodies, led me to expect that the same kind of dependence upon the intervening particles would be found to exist in magnetic action; and I published certain experiments and considerations on this point seven years ago (1709—1736). I could not then discover any peculiar condition of the intervening substance or diamagnetic; but now that I have been able to make out such a state, which is not only a state of tension (2227), but dependent entirely upon the magnetic lines which pass through the substance, I am more than ever encouraged to believe that the view then advanced is correct.

2241. Although the magnetic and electric forces *appear* to exert no power on the ordinary or on the depolarized ray of light, we can hardly doubt but that they have some special influence, which probably will soon be made apparent by experiment. Neither can it be supposed otherwise than that the same kind of action should take place on the other forms of radiant agents as heat and chemical force.

2242. This mode of magnetic and electric action, and the phænomena presented by it, will, I hope, greatly assist hereafter in the investigation of the nature of transparent bodies, of light, of magnets, and their action one on another, or on magnetic substances. I am at this time engaged in investigating the new magnetic condition, and shall shortly send a further account of it to the Royal Society. What the possible effect of the force may be in the earth as a whole, or in magnets, or in relation to the sun, and what may be the best

means of causing light to evolve electricity and magnetism, are thoughts continually pressing upon the mind; but it will be better to occupy both time and thought, aided by experiment, in the investigation and development of real truth, than to use them in the invention of suppositions which may or may not be founded on, or consistent with fact.

Royal Institution, Oct. 29, 1845.

---

L. *On the Cause of remarkably Mild Winters which occasionally occur in England.* By Lieut.-Colonel SABINE, R.A., For. Sec. R.S.

*To the Editors of the Philosophical Magazine and Journal.*

GENTLEMEN,

THE unusual character of the winter which we have just experienced, together with its effects which we are now witnessing upon our gardens and fields, and its influence on the public health as evidenced by the bills of mortality, should make it an object not only of scientific, but of general interest, to endeavour to trace out the cause of so remarkable a phenomenon. By a memorandum with which the Astronomer Royal has been so obliging as to furnish me, it appears that the mean temperature in December, January and February, exceeded the mean temperature of the same months in the preceding year by the amounts respectively of  $8^{\circ}7$ ,  $5^{\circ}3$ ,  $11^{\circ}2$ ; on an average above  $8^{\circ}$  for three months. An excess of temperature of such amount and such continuance, must surely, one would suppose, have some sufficiently notable cause. I am not aware that any probable cause has yet been suggested; but should you oblige me by inserting this communication, it may at least be of use in commencing the discussion, and possibly in eliciting the opinions of others, whose views on the subject the public may naturally desire to know.

The winter which within my recollection most nearly resembled the present, was that of 1821-1822, and undoubtedly the resemblance is in many respects very striking. For the peculiarity in that year there was a cause assigned, adequate I believe to account for all the phenomena, and of which the existence was proved: I allude to the extension of the Gulf-stream in that year to the coast of Europe, instead of its terminating as it usually does about the meridian of the Azores. In the winter of 1821-1822, the warm water of the Gulf-stream spread itself beyond its usual bounds over a space of ocean which may be roughly estimated as exceeding 600 miles in latitude and 1000 in longitude, carrying with it water several de-

gress higher than the temperature of the sea in ordinary years in the same parallels. The facts, both in respect to the Gulf-stream, and to the peculiarities of the winter in that year, were stated in the volume of *Pendulum* and other observations which I published in 1825; perhaps the statement of them now will be most satisfactorily given in the words which were then used: and I have the less hesitation in introducing an extract from that work, because it was published many years ago, and is, I believe, but little known, at least in this country. The statement was as follows:—

“In the passage of the *Iphigenia* from England to the coast of Africa, a remarkable and very interesting evidence was obtained, by observations on the temperature of the sea, of the accidental presence in that year of the water of the Gulf-stream, in longitudes much to the eastward of its ordinary extension.

“The *Iphigenia* sailed from Plymouth on the 4th of January [1822], after an almost continuous succession of very heavy westerly and south-westerly gales, by which she had been repeatedly driven back and detained in the ports of the Channel; the following memorandum exhibits her position at noon on each day of her subsequent voyage from Plymouth to Madeira, and from thence to the Cape Verd Islands, the temperature of the air in the shade and to windward, and that of the surface of the sea; it also exhibits in comparison, the ordinary temperature of the ocean at that season, in the respective parallels, which Major Rennell has been so kind as to permit me to insert on his authority, as an approximation founded on his extensive inquiries; the last column shows the excess or defect in the temperature observed in the *Iphigenia*'s passage.

Date.	Latitude N.	Longitude W.	Air.	Surface-water.		Excess or defect.	
				Observed.	Usual.		
Plymouth to Madeira.	1822. Jan. 5	47 30	7 30	47·0	49·0	50·0	—1
	6	44 20	9 30	52·5	55·7	52·5	+3·2
	7	41 22	11 37	54·0	58·2	54·0	+4·2
	8	38 54	13 20	54·2	61·7	55·7	+6
	9	No observation.		56·0	63·0	58·0	+5
Madeira to Cape Verd Islands.	10	33 40	15 30	60·7	64·0	60·0	+4
	19	26 00	17 50	66·0	65·5	67·0	—1·5
	20	24 30	18 50	68·0	67·0	68·4	—1·4
	21	23 06	20 00	69·0	69·0	69·5	—0·5
	22	21 02	21 27	69·5	69·5	71·2	—1·7
	23	19 20	23 00	70·6	70·2	71·6	—1·4

“It is seen by the preceding memorandum, that in the passage from Plymouth to Madeira, the Iphigenia found the temperature of the sea, between the parallels of  $44\frac{1}{3}^{\circ}$  and  $33\frac{2}{3}^{\circ}$ , several degrees warmer than its usual temperature in the same season; namely  $3^{\circ}2$  in  $44\frac{1}{3}^{\circ}$ , increasing to  $6^{\circ}$  in  $39^{\circ}$ , and again diminishing to  $4^{\circ}$  in  $33\frac{2}{3}^{\circ}$ ; whilst at the same period, the general temperature of the ocean in the adjoining parallels, both to the northward and to the southward, even as far as the Cape Verd Islands in  $19\frac{2}{3}^{\circ}$ , was colder by a degree and upwards than the usual average. The evidence of many careful observers at different seasons and in different years, whose observations have been collected and compared by Major Rennell, has satisfactorily shown, that the water of the Gulf-stream, distinguished by the high temperature which it brings from its origin in the Gulf of Mexico, is not usually found to extend to the eastward of the Azores. Vessels navigating the ocean between the Azores and the continent of Europe, find at all seasons a temperature progressively increasing as they approach the sun; the absolute amount varies according to the season, the maximum in summer being about  $14^{\circ}$  warmer than the maximum in winter; but the progression in respect to latitude is regular, and is nearly the same in winter as in summer, being an increase of  $3^{\circ}$  of Fahr. for every  $5^{\circ}$  of latitude. It is further observed, that the ordinary condition of the temperature, in the part of the ocean under notice, is little subject to disturbance, and that in any particular parallel and season, the limits of variation in different years are very small; after westerly winds of much strength or continuance, the sea in all the parallels is rather colder than the average temperature, on account of the increased velocity communicated to the general set of the waters of the north-eastern Atlantic towards the south. To the heavy westerly gales which had prevailed almost without intermission in the last fortnight in November, and during the whole of December, may therefore be attributed the colder temperatures observed in the latitude of  $47\frac{1}{2}^{\circ}$ , and in those between  $26^{\circ}$  and  $19\frac{1}{3}^{\circ}$ .

“If doubt could exist in regard to the higher temperatures between  $44\frac{1}{3}^{\circ}$  and  $33\frac{2}{3}^{\circ}$  being a consequence of the extension in that year of the Gulf-stream in the direction of its general course, it might be removed by a circumstance well-deserving of notice, namely, that the greatest excess above the natural temperature of the ocean was found in or about the latitude of  $39^{\circ}$ , being the parallel where the middle of the stream, indicated by the warmest water, would arrive, by continuing to flow to the eastward of the Azores, in the prolongation of the great circle in which it is known to reach the mid-Atlantic.

“One previous and similar instance is on record, in which the water of the Gulf-stream was traced by its temperature quite across the Atlantic to the coast of Europe; this was by Dr. Franklin, in a passage from the United States to France, in November 1776\*. The latter part of his voyage, *i. e.* from the meridian of  $35^{\circ}$  to the Bay of Biscay, was performed, with little deviation, in the latitude of  $45^{\circ}$ ; in this run exceeding 1200 miles, in a parallel of which the usual temperature, towards the close of November, is about  $55\frac{1}{2}^{\circ}$ , he found  $63^{\circ}$  in the longitude of  $35^{\circ}$  W., diminishing to  $60^{\circ}$  in the Bay of Biscay; and  $61^{\circ}$  in  $10^{\circ}$  west longitude, near the same spot where the *Iphigenia* found  $55^{\circ}\cdot7$  on the 6th of January, being about five weeks later in the season. At this spot then, where the *Iphigenia* crossed Dr. Franklin’s track, the temperature in November 1776 was  $5\frac{1}{2}^{\circ}$ , and in January 1822,  $3^{\circ}\cdot2$  above the ordinary temperature of the season.

“There can be little hesitation in attributing the unusual extension of the stream in particular years to its greater initial velocity, occasioned by a more than ordinary difference in the levels of the Gulf of Mexico and of the Atlantic; it has been computed by Major Rennell, from the known velocity of the stream at various points of its course, that in the summer months, when its rapidity is greatest, the water requires about eleven weeks to run from the outlet of the Gulf of Mexico to the Azores, being about 3000 geographical miles; and he has further supposed, in the case of the water of which the temperature was examined by Dr. Franklin, that perhaps not less than three months were occupied, in addition, by its passage to the coasts of Europe, being altogether a course exceeding 4000 geographical miles. On this supposition, the water of the latter end of November 1776 may have quitted the Gulf of Mexico, with a temperature of  $83^{\circ}$ , in June; and that of January 1822, towards the end of July, with nearly the same temperature. The summer months, particularly July and August, are those of the greatest initial velocity of the stream, because it is the period when the level of the Caribbean Sea and Gulf of Mexico is most deranged.

“It is not difficult to imagine that the space between the Azores and the coasts of the old continent, being traversed by the stream, slowly as it must be, at a much colder season in the instance observed by the *Iphigenia* than in that by Dr. Franklin, its temperature may have been cooled thereby to a nearer approximation to the natural temperature of the ocean in the former than in the latter case; and that the difference

\* Franklin’s Works, 8vo, London, 1806, vol. ii. pp. 200, 201.

between the excess of  $5^{\circ}5$  in November, and of  $3^{\circ}2$  in January, may be thus accounted for.

“If the explanation of the apparently very unusual facts observed by Dr. Franklin in 1776, and by the *Iphigenia* in 1822, be correct, how highly curious is the connexion thus traced between a more than ordinary strength of the winds within the tropics in the summer, occasioning the derangement of the level of the Mexican and Caribbean seas, and the high temperature of the sea between the British Channel and Madeira, in the following winter!

“Nor is the probable meteorological influence undeserving of attention, of so considerable an increase in the temperature of the surface water over an extent of ocean exceeding 600 miles in latitude and 1000 in longitude, situated so importantly in relation to the western parts of Europe. It is at least a remarkable coincidence, that in November and December 1821, and in January 1822, the state of the weather was so unusual in the southern parts of Great Britain and in France, as to have excited general observation; in the meteorological Journals of the period it is characterized as ‘most extraordinarily hot, damp, stormy, and oppressive;’ it is stated, ‘that an unusual quantity of rain fell both in November and December, but particularly in the latter;’ that ‘the gales from the west and south-west were almost without intermission,’ and that in December, the mercury in the barometer was lower than it had been known for thirty-five years before\*.”

\* “The following description of this very remarkable winter is extracted from Mr. Daniell’s *Essay on the Climate of London* (Meteorological Essays. London, 1823, pages 297 and 298), and becomes highly curious when viewed in connexion with the unusual temperature of the ocean in the direction in which the principal winds proceeded.

“November 1821 differed from the mean, and from both the preceding years, in a very extraordinary way. The average temperature was  $5^{\circ}$  above the usual amount; and although its dryness was in excess” (the relative dryness in consequence of the increased temperature) “the quantity of rain exceeded the mean quantity by one-half. The barometer on the whole was not below the mean. All the low lands were flooded, and the sowing of wheat very much interrupted by the wet.

“In December the quantity of rain was very nearly double its usual amount. The barometer averaged considerably below the mean, and descended lower than had been known for thirty-five years. Its range was from 30.27 inches to 28.12 inches. The temperature was still high for the season, and the weather continued, as in the last month, in an uninterrupted course of wind and rain; the former often approaching to a hurricane, and the latter inundating all the low grounds. The water-sodden state of the soil, in many parts, prevented wheat-sowing, or following the land at the regular season. The mild temperature pushed forward all the early-sown wheats to a height and luxuriance scarcely ever before witnessed. The grass and every green production increased in an equal proportion.

“January, 1822. This most extraordinary season still continued above

It is impossible to read this description of the winter of 1821-1822 without being struck with the many features which it has in common with that of the present year. The excess of heat in both amounted to several degrees, and continued through several months. There were similar floods in many parts of England in the early part and middle of this winter; and these were not confined to England only, but extended, as in 1821-1822, to many of the rivers of western Europe. The tension of the vapour conveyed to the shores of the British Channel in December, January and February last, was nearly  $\frac{1}{3}$ rd part greater, as appears by the Greenwich Observations, than in the same months of the preceding year; although in consequence of the much higher temperature, the humidity of the air, or the ratio of the humidity to saturation, has been less. This was also the case in 1821-1822. We have had an unusual prevalence of westerly and south-westerly winds at the season when they are ordinarily replaced in a much greater proportion by the dry and cold winds which come to us from the interior of the great continents of Europe and Asia. If in the southern parts of Britain, and on the shores of the British Channel, we have been less severely affected by storms and extreme barometrical depressions than was the case in 1821-1822, we may possibly owe the comparative exemption to the fact that the excess of heat above the mean has been greater in the winter of 1845-1846 than it was in 1821-1822; whence we may infer perhaps that the conflict of the opposing currents of the atmosphere has been removed in the present year further to the north and north-east than on the former occasion; it is at the limits which are reached by the warm and humid current proceeding from the south-west, and in the localities where it encounters the dry and cold stream pressing from the east and north-east, that the greater atmospherical derangements are produced, and these have been experienced in the northern parts of Britain.

The similarity of the two winters having thus been shown, and specially their agreement in those features in which they differ from ordinary winters, it will naturally be asked, what evidence we have to prove or disprove an extension of the Gulf-

the mean temperature, but the rain, as if exhausted in the preceding month, fell much below the usual quantity in this. There was not one day on which the frost lasted during the twenty-four hours.

“ ‘ Serious apprehensions were entertained, lest the wheats, drawn up as they had been by the warm and moist weather, without the slightest check from frost, should be exhausted by excessive vegetation, and ultimately be more productive in straw than corn.

“ ‘ The month of February, still 5° above the mean temperature, ended a winter which never has been paralleled.’ ”



stream in the present year, similar to that which took place in 1821. To this it must be replied, that strange as it may appear, this remarkable phænomenon may take place in any year without our having other knowledge of it than by its effects, although it occurs at so short a distance from our ports, from whence so many hundred vessels are continually crossing and recrossing the part of the ocean where a few simple observations with a thermometer would serve to make it known. We have no organized means of learning an occurrence which, whether it be or be not the cause of the present extremely mild winter, cannot fail whenever it does occur to affect materially and for a considerable length of time the climate of an extensive district of the globe including our own islands. History has recorded two instances in which the extension of the Gulf-stream is known to have taken place; and in both we owe our knowledge of it to the casual observations of an accidental voyager. Some one there may be in the present winter whose curiosity may have induced him, in the well-frequented passage between England and Madeira, to dip a thermometer in the sea once or twice a day, and who may therefore, perhaps unconsciously, be in possession of the very facts which it is desirable to know; in such case this communication, should it meet his eye, may be the means of inducing their publication. It is desirable however that we should not be thus altogether dependent on accident for information which may have even greater practical than scientific value; happily it is well known that suggestions of this nature, when really deserving attention, do not pass unheeded by our excellent Hydrographer, to whose department such subjects seem naturally to belong\*.

But not only might we by such means be occasionally informed in November or December that we had probably to expect a continuance of very mild weather through January and February; it is not unreasonable to suppose that such winters might well be anticipated at a still earlier period of the year; ships sail faster than the Gulf-stream flows, and a more than usual difference existing in the levels of the Gulf of Mexico and the Atlantic, or a more than usual initial velocity of the stream itself, with the consequent probability of a winter of unusual mildness in Europe, might be known in England in the summer or in the early autumn; or even going

\* It is much to be wished that a society existed in England which should charge itself with the many interesting and important considerations belonging to *physical* geography. Did the object and scope of the Royal Geographical Society embrace physical as well as descriptive geography, it cannot be doubted that science and the public would be greatly benefited.

back to a yet earlier stage of the phænomenon, we might be apprised that the causes which operate in producing the derangement of the level of the Caribbean and Mexican seas were prevailing in any particular year in an unusually high degree.

I wish in conclusion to guard against the possibility of being understood to suppose that amidst the variety of incidents by which our climate is affected, there may not be others which may be influential in the production of winters of unusual mildness in an equal, or even in a greater degree than the extension of the Gulf-stream; or, that whenever the stream reaches the coasts of Europe, its influence on our climate must necessarily occasion winters like that of 1821-1822, or 1845-1846. It is reasonable to believe that there may be degrees of initial velocity between that which is usual and that which is extreme. There may also be counteracting or qualifying causes with which we are as yet wholly unacquainted. The object of this communication is rather to recall to recollection, on the occasion of the present remarkable winter, the coincidence that was discovered between the similar winter in 1821-1822, and the extension of the stream; and to promote the adoption of such simple means as may supply additional evidence, whereby we may discern between coincidence observed on a single occasion, and connexion which may be established by the observation of repeated coincidence.

I am, Gentlemen,

Your obedient Servant,

Woolwich, March 17th, 1846.

EDWARD SABINE.

LI. *Observations on the Recent Researches of Prof. Faraday.*

By M. POUILLET\*.

FOR some months past there has been much talk about a new series of researches by M. Faraday, the result of which was one of the most important discoveries, the action of magnetism on light. Two authentic documents have at length reached us on this subject; one is published in the January number of the Philosophical Magazine, being an abstract of the sitting of the Royal Society of the 27th of November, the other was communicated at the last meeting of the French Academy by M. Dumas in the name of Mr. Faraday himself. Various results are announced in these two publications, but only one fact is presented with some development,

\* Translated from the *Comptes Rendus*, for January 26, 1846.—The researches of Prof. Faraday, referred to in this paper, are given entire in our present Number.

it is that which relates to the action of an electro-magnet on a ray of polarized light in turning its plane of polarization either to the right or left, according to the relative directions of the luminous ray and of the resultant of the magnetic actions.

This fact is justly considered by Faraday as a fundamental one, for hitherto there is nothing analogous in science; and it constitutes of itself a discovery of the very highest importance. Undoubtedly many persons have hastened to repeat and investigate it in order to ascertain its perfect accuracy, and to find out the most marked characters and the most essential conditions. Immediately after having read the *Philosophical Magazine* I set to work, as I stated at the last meeting of the Academy, but my first trials having been without result, and other persons not having been more fortunate than myself in the attempt, it appeared to me necessary to resume them with greater attention, varying the mode of experiment and making up in the best manner I was able for the want of precision in the directions which had come to my knowledge.

I hasten to lay before the Academy today the result of these researches, and with a twofold motive; in the first place, to render homage to the author of the discovery, and in the next place, to furnish those physicists who may desire to follow this new path of science with some indications that may be of service to them, if, as I believe, they add anything to what has hitherto been published on the subject.

The apparatus employed by me is composed,—1st, of a Bunsen's battery; 2nd, of one or more electro-magnets; 3rd, of M. Soleil's instrument for exhibiting the least angular displacements of the planes of polarization; 4th, of the various substances to be submitted to examination. The elements of the Bunsen's battery are of the ordinary size; in the majority of cases ten suffice to render the phenomenon perceptible; but to measure it and to compare the intensities with a certain approximation, forty, fifty, and even 100 elements must be employed. The electro-magnets are capable of supporting 1600 pounds when excited by a battery of twenty pairs. They are soft iron cylinders, seven to eight centimetres in diameter and about fifty centimetres in length, which are curved in a horse-shoe form, the distance of the axes of the two arms or poles being not more than fifteen to twelve centimetres. From 500 to 600 metres of copper wire, coated twice with silk, are wound round each arm. The instrument of M. Soleil is composed of two parts, one objective, the other ocular. The objective part, or that which is turned towards the light, is nothing more than a Nicol's prism, behind which

is a system of two juxtaposed plates of quartz cemented by one margin, and worked together in order to fulfill a double condition of giving them exactly the same thickness and rendering each perfectly perpendicular to the axis. The surface of junction of these plates being parallel to the pencil of light and occupying the centre of its breadth, it is evident that the first half of the pencil traverses one of the plates only, and the second half the other plate; and as they were selected of opposite rotatory power, the first half of the polarized pencil is found to have its planes of polarization deviated, for instance towards the right, by a certain angle, and the second half, on the contrary, has its planes of polarization deviated towards the left by a perfectly equal angular magnitude. The magnitude of these deviations depends on the common thickness of the two plates, which is usually from five to six millimetres.

The ocular portion, or that directed towards the eye, consists, in the first place, of a thick plate of rock-crystal, likewise perpendicular to the axis, having for instance a rotatory power to the right, and a thickness of five millimetres very accurately determined by the spherometer. Behind this plate is the compensator, composed of two equal prismatic plates, provided with a similar rotatory power towards the left, *i. e.* in a contrary direction to the first. These two prisms, opposed like two wedges by their acute angle, are moved simultaneously by the same spring; they slide one upon the other, to be arranged sometimes by their less sometimes by their greater thickness, and thus always form a system equivalent to a parallel plate, but one which would vary from the thickness 0 to nearly double that of the base of each prism. To avoid the deviations which the light might experience from the variable distance of these prisms and the obliquity of the surfaces, each one is compensated by a glass prism.

Lastly, behind the compensator is a doubly refracting achromatic prism and a small Galilean telescope, to which the eye is applied to observe the pencil of light which has passed both the objective portion, the intermediate bodies submitted to examination, and the ocular portion of the instrument.

The graduation of the compensator is easily made; and when this has once been done with sufficient care, the instrument indicates that the cause, whatever it be, which produces the deviation in the plane of polarization has an intensity equivalent to that of a plate of quartz of a known thickness; always, be it understood, on the condition that this cause exercises on the various simple lights actions comparable to that which the quartz exercises.

The instrument of M. Soleil, the construction of which I have just described, must be separated into two parts for the experiments under consideration. The objective and the ocular parts were mounted separately on my frame of diffraction\*, which is most readily adapted for all the researches in which it is required to centre the apparatus on the same axis. A common lamp is placed before the objective part, and a strong magnifier gives a pencil of light closely parallel, which being propagated in the direction of the common axis, traverses successively the object-glass, the pieces subjected to the test, and the ocular; the distance between the objective and the ocular may vary between tolerably distant limits, for it may extend to nearly two metres, or only to a few centimetres, according to the nature of the observations.

It is important to remark that the pencil of light is always horizontal, and the apparatus was accidentally arranged so that the light was propagated from south to north, which may assist us to define more easily the relative positions of the polarized ray, of the electro-magnets, and of the bodies on which they act.

The electro-magnet is horizontal, that is to say, the plane of the axes of its two branches is horizontal, and precisely at the height of the pencil of light which traverses the apparatus; moreover, the vertical plane, formed by the extremities of the two branches or by the poles of the electro-magnet, is parallel to this pencil, and may approach it more or less. This being settled, if it be desired to submit to experiment, for example, a parallelepipedon of flint-glass of ten or twelve centimetres in length and terminated perpendicularly to its length by two parallel planes, this parallelepipedon is first arranged so that the ray polarized by the objective traverses it in the direction of its axis, and if the flint-glass is pure and well-annealed, as it must be for the success of the experiment, its interposition produces neither deviation nor coloration in the ray of light. The electro-magnet is then approached, arranging it in the same manner as if the piece of flint-glass were a piece of iron to close it, and there is even no inconvenience in arranging it so that the two poles of the electro-magnet are in contact with the flint-glass; the middle of the length of the latter corresponds consequently to the interval which exists between the two arms of the electro-magnet.

When these arrangements have been made, a current is passed, and suddenly it is seen that the two tints of the red image, which correspond to the two opposed plates of the quartz of the object-glass, cease to be identical. Let us sup-

\* See my *Eléments de Physique*, 4th edition, vol. ii. pl. 26.

pose, for example, that the one on the right has turned blue; if the current is passed in a contrary direction, it is that on the left which this time turns blue in the same manner. Thus, by reversing the poles of the electro-magnet, the action which it exercises on the flint-glass, or on the light which traverses it, is also suddenly reversed. Here then we see the action in question rendered evident in the most striking and incontestable manner.

In the circumstances of which I have just spoken, ten elements are more than sufficient to exhibit it to a practised eye; but with a hundred elements, it assumes such an intensity that persons the most unaccustomed to these kind of observations could not fail to perceive it as a perfectly characterized phenomenon.

Before seeking to ascertain whether this effect, at once so novel and so extraordinary, results from a direct action of the magnetic fluid upon light, or from an indirect action, in which the ponderable matter of the flint-glass intervenes, or at least the collective forces to which this matter is subjected in order to exist in molecular equilibrium, it is necessary first to determine precisely what is the nature of the effect produced, and to seek above all to measure its intensity, in order to ascertain what are the conditions under which the phenomenon is shown with the greatest energy.

For this purpose, instead of observing directly the coloured tints which the quartz gives by the lamp perpendicular to the axis, it is necessary to recompose what M. Biot has called *the tint of passage*. This is done by placing before the objective several systems of blue and greenish glasses; but I found in the cabinet of the Conservatoire some glasses very slightly coloured blue, which give to this tint a sensitiveness still greater than that obtainable by other means. When these glasses are interposed in the pencil, the tints of the quartz become of a light lilac, on which the least changes of shade are appreciable; the uncertainties which are presented by the zero of the compensator disappear, and it becomes possible not only to perceive, but to measure the effects which correspond to thicknesses of quartz of a hundredth of a millimetre.

The instrument thus modified, the compensator being at zero, and the polarizing prisms of the object-glass and of the ocular being suitably regulated in their relative positions, the experiment may be proceeded with; only there is one thing which requires mention, not to pay attention to the yellow image, but to look exclusively at the lilac image, the two halves of which are then exactly of the same shade.

As soon as the current passes, one of the halves of this image, for instance that on the right, is seen to turn blue; we observe that this tint is persistent as the current itself, and we may be convinced that, from the first instant, it acquired its whole value, that is to say, that the prolonged duration of the action adds nothing perceptible to it. The compensator is then moved in the proper direction; the difference of the tints gradually disappears in proportion as it advances, and with a little practice the point at which the equality is re-established is soon found. The number of divisions is noted down, and we thus obtain a measure, or at least an approximate measure of the effect produced,—say twenty divisions.

When subsequently the current is passed in a contrary direction, it is the other half of the tint, that to the left, which turns blue, and it is in the other direction that the compensator has to be moved to re-establish the equality. No interval of time is appreciable between the change of the current and the change of effect upon the light, and it is again instantaneously that the tint takes all its value. When the optical apparatus is well-adjusted, and the electrical communications are equally good in both directions, the ground gone over by the compensator is the same in the two cases, that is to say, that if it progressed in the first twenty divisions to the right, it should in the second proceed twenty degrees to the left.

These opposite effects and the corresponding measures may be repeated indefinitely, either with the same or a different number of pairs of the battery; and a few hours are sufficient, during which the action of the battery is nearly constant, to pass in review a great number of diaphanous substances, and to obtain a first approximation on the relative sensitiveness with which they obey the magnetic influence.

When the substances submitted to the test are more or less coloured, it is necessary to vary the systems of glasses intended to produce the tint of passage, and we do not always succeed in composing a tint equally delicate and easy of observation. It might happen consequently that some substances, even slightly coloured, when submitted to these modes of observation, would appear much less energetic than they are in reality.

Let us pause then at the diaphanous substances, and observe that in the experiment with the flint-glass cited above, it was necessary to advance the compensator twenty divisions to the right and twenty divisions to the left, according as the current passed in one direction or the other. Let us bear in mind, that if, instead of interposing on the passage of the pencil a prism of flint-glass submitted to the electro-magnet,

there had been interposed, without magnetic action, a lamina of quartz perpendicular to the axis, of a proper thickness, turning to the right in the first case and to the left in the second, it is certain that the equality of the tints would have been re-established by the same movements of the compensator. Now, it is known that the effect produced by such a lamina of quartz would have been to turn the plane of polarization from right to left, whence it seems very natural and legitimate to conclude, that the flint-glass subjected to the magnetic action has produced the same effect as this lamina of quartz, that is to say, that it has also turned the plane of polarization to the right for one direction of current, and to the left for the contrary direction. This is, in fact, the conclusion to which Mr. Faraday has come, and he has characterized this new action of magnetism upon light, by stating that the magnetism turns the plane of polarization of the luminous ray submitted to its influence under certain conditions, and that the direction of this rotation is connected with that of the current.

Quartz, and the other substances which, of themselves, by their nature or structure have, without the intervention of magnetism, the permanent property of turning the planes of polarization, exert this action with variable intensities on the different elements constituting white light; and there are dispersive powers for this rotation, as there are different dispersive powers for refraction. It would be very important to make in this respect some researches upon the substances which acquire this property by the magnetic action, analogous to those very remarkable ones which M. Biot made upon the former. The apparatus which I have used must be very much modified to be adapted to this class of experiments; it serves to show the phænomena very distinctly, rather than to measure them in their more delicate details. Such an investigation, however, cannot be undertaken with phænomena so little developed as those which I have obtained; for in such limits they might perhaps be as well explained by partial depolarizations towards the right and left as by the rotation itself of the plane of polarization, which, moreover, would not detract anything, and would perhaps add to their importance.

As I just stated, the plane of polarization, in that specimen of the flint-glass which gave the most energetic effects, was diverted by the magnetic action as much as it would have been by the action of a plate of quartz two-tenths of a millimetre in thickness; now, since by changing the direction of the current, the rotation takes place in an opposite direction, it is seen



that the total effect obtained by passing from the magnetic action which is exerted in one direction to that which is exerted in the other, is equal to that which would be produced by a plate of quartz four-tenths of a millimetre in thickness. Such, up to the present time, is the maximum effect which I have been able to obtain. As we have now a means of comparing the intensities of this force, it will be very easy to see how it will be modified by the different relative positions of the electro-magnet and of the piece of flint-glass.

The following are the observations I have made with respect to this point:—

1. If, instead of placing the electro-magnet in contact with the piece of flint-glass, it is removed parallel to itself in the same horizontal plane, and so that the vertical plane separating the two arms corresponds always to the middle of the flint-glass, the action diminishes, but feebly in proportion as the distance increases, so that at the distance of ten centimetres, it is still a considerable proportion of what it was when it was actually in contact.

2. If the electro-magnet is again placed in contact, and the piece of flint-glass slid in the direction of the ray of light to subject it to the action of one only of the poles of the magnet, a moment arrives when the action is wholly null; then, if it is still slid in the same direction, removing it more and more from its primitive position, until it is placed beyond the pole to which it is submitted, the action begins anew; but then it is contrary to what it was at first.

These observations appear to lead to three important consequences:—

It first results, that if we consider the unknown action of the magnet on the flint-glass as being produced by attractions and repulsions, the effect is null when the resultant of these attractive and repulsive forces is perpendicular to the direction of the polarized ray; and it is at its maximum, on the contrary, when this resultant is parallel to the ray. We may thus, from these considerations, form a just idea of the direction in which it acts, for in considering, *always hypothetically*, the piece of flint-glass as a piece of soft iron, acquiring two poles from the influence of the magnet, the movement of the plane of polarization occurs to the right when the light enters by the south pole, and proceeds from the south to the north pole, and it occurs to the left when the light enters by the north pole; consequently, whatever be the position of the piece of flint-glass, if two observations are made on it without touching it, and without disarranging the electrical apparatus, but merely turning the optical apparatus to cause the light

to enter successively in the two directions, we shall see, in the first case, the effect to the right, and in the second the effect to the left, which establishes, as Mr. Faraday has pointed out, a difference, at least apparent, between the substances which have the permanent property of turning the planes of polarization and those which acquire it by magnetic action.

In the second place it results, that on experimenting in this way care must be taken to give to the pieces subjected to the electro-magnet a length greater than the distance of the axes of the two arms; for the portions which would exceed those axes would receive similar modifications among one another, and opposed to that which the central portion would receive; it may even be presumed that the compensation might obtain exactly, so that with a connecting piece exceeding the breadth of the magnet, the action might be perfectly null. This result seems to me opposed to that which is pointed out by Mr. Faraday, namely, that the effect is proportional to the length of the piece subjected to the experiments.

It results, lastly, that in order to obtain a greater effect, two electro-magnets, opposed to one another, may be presented to the piece of flint-glass, so that the poles of the same name face one another. This I have verified, and it is even by the assistance of two electro-magnets thus opposed that I have obtained the maximum effect of which I have spoken above. By placing thus several similar systems in succession on the same pencil, the effect would be, without doubt, tripled, &c.

It has appeared to me very important to examine whether the position of the plane of polarization, with relation to the horizontal plane of the electro-magnet, had any influence on the energy of the action; but whether the plane of polarization be itself horizontal, vertical or intermediate, the results appeared to me to remain perceptibly the same.

I have hitherto spoken only of flint-glass, but I have subjected to experiment all the other solid transparent bodies which I have been able to procure; for instance, various kinds of flint-glass, and doubtless of different composition, crown-glass, and glass of all kinds, coloured with copper, gold, chromium, &c., and also rock-salt. All these bodies present, although with less intensity, the same phænomena as the flint-glass; unfortunately the samples of crown-glass are sometimes so annealed as to modify the colours, and which does not allow of their being compared with other bodies; nevertheless, after the attempts which I have been able to make on some less imperfect specimens, I am led to think that the action of the crown-glass has an intensity comprised between the half and the two-thirds of that of the flint-glass.

The chloride of sodium has an action very analogous to that of the flint-glass.

I have also subjected to experiment some transparent or coloured liquids; these experiments were made in a trough formed of parallel glasses, having a length of thirteen centimetres, equal to the distance of the axes of the electro-magnets, a breadth of three centimetres, and a depth of five centimetres. The trough being empty, and the electro-magnets being in action, no sensible effect was produced by the parallel glasses which formed the extremities.

The intensity of all these liquids is very nearly equal to that of the crown-glass; the most energetic however appeared to me to be olive oil, distilled water, concentrated ammonia, and pure nitric acid; and the less energetic, acetic acid, sulphuric acid, ferrocyanide of potassium, and ferrocyanate of magnesia. It appeared to me certain that several bodies dissolved in distilled water weakened its effects.

Mr. Faraday states that manganese, chromium, and cerium are magnetic after the manner of iron, and that all the compounds of these bodies preserve this property more or less. I had long ago proved the first fact for manganese, and in the course of last summer I proved it for very pure chromium obtained by the battery, both from chromic acid and from sulphate of chromium. With regard to the magnetic compounds, I have studied them recently by a very simple and very easy process, which consists in arranging a powerful electro-magnet, with its poles at top, forming a horizontal plane; a thin paper is stretched over each pole, in contact with the iron itself, and it is then only requisite to throw upon this paper some very fine particles of the substance to be examined, and to give the paper some slight vibrations, which put them in motion. The particles arrange and fix themselves on the circle which corresponds to the terminal bar of the iron of the electro-magnet, and describe the circle with great precision. By this means I have ascertained that almost all the compounds of magnetic metals are, in fact, more or less magnetic; prussian blue and the sesquichloride of chromium (of M. Peligot) are so in a remarkable manner. There are some compounds however which are exceptions to this rule; such, for example, are the double cyanide of iron and of potassium, the chromate of silver, and the bichromate of potass.

Other metals, as platina sponge and arsenic, exhibit a perceptible action; but it would require to be verified on perfectly pure specimens.

Bismuth presents other phenomena; instead of forming a

circle, like the magnetic metals, it forms two concentric circles, leaving thus a narrow white band, in the very place where the other metals form a circle, as if it were repelled by the more lively action of the iron armature of the magnet. The effect is so marked, that on mixing, for example, some sesquichloride of chromium very finely pulverized with some bismuth, likewise in very fine powder, the violet circle of the chloride is seen, and the two circles of the bismuth which are separate from them, although very near. Amber seems to give the same appearances as the bismuth, though in a much weaker degree.

No attractive or repulsive effect is observed by this means, either on very pure antimony or on the other metals, binary or other compounds (among the rare metals, I have only experimented on tellurium and the uranium of M. Peligot), nor on the alkalis, sulphur, iodine, charcoal, and diamond. I regret that I had not at my disposal either cerium or any of its compounds.

These negative results cannot invalidate in the least the general proposition of M. Faraday, who has doubtless operated with more delicate means or with more energetic magnets. I merely mention them here to point out the easy process which I have employed, and the limit of its sensibility.

There is another process for investigating the magnetic properties,—that which was employed by Coulomb when he discovered that all bodies are subject to the influence of magnets, and which has been since employed in the same view by many experimentalists, and very recently by M. Ed. Becquerel (*Comptes Rendus*, vol. xx. p. 1708). Mr. Faraday appears to have employed it; but doubtless, from the weakness of my electro-magnets, although excited by a battery of 100 pairs, I have not obtained the same results as he; in my experiments, bismuth and amber are the only two substances which took a direction perpendicular to the line of the poles, and without doubt the relation existing between this direction of the bismuth and the effect of repulsion which the fine powder of that body experiences from the part of the armature of the magnet will appear highly remarkable. These two mechanical actions of magnetism upon bodies—the attraction and repulsion of fine powders, placed almost in contact with one of the poles, and the direction given to more considerable masses, oscillating in the presence of the two poles—appear therefore to be dependent one upon the other; but in what degree are they connected with the third action, the optical action which Mr. Faraday has just discovered?

Admitting with this philosopher that all the substances

which are not magnetic after the manner of iron, are *diamagnetic* or magnetic after the manner of bismuth, we should be led to conclude immediately that the optical action being concomitant with a certain mechanical action, it is at least presumable that this action is exerted upon the bodies, and not directly and immediately on the light which passes through them.

But if it happens, as in my experiments, either from the relative weakness of my magnets or from the imperfection of the methods which I have employed, or from other causes—if it happens that the various kinds of glasses, distilled water, the fatty bodies, &c., which are so sensitive to the optical action, are nevertheless insensible to the mechanical action of the magnetism, it would not be a reason to conclude that magnetism acts directly upon the light itself; a conclusion which, moreover, would only have a precise meaning in the system of emission, for in the undulatory theory, which seems at present so completely demonstrated, it is the æther of the body submitted to the experiment which would be modified by the magnetism, and it would doubtless be very difficult to recognise whether it is modified without any participation of the ponderable matter of the body with which it is so intimately connected.

LII. *On the Aberration of Light.* By G. G. STOKES, M.A.,  
Fellow of Pembroke College, Cambridge\*.

I WISH to say a few words more on the subject of aberration, to prevent misapprehension. It is evident from Prof. Challis's last communication, that we differ merely as to the phænomenon which we understand by the term "aberration of light." When the position of a star has been corrected for refraction, precession, and nutation, and proper motion if it has any, let  $s$  be its mean annual place referred to the celestial sphere,  $s_1$  the point to which the star is referred by astronomical measurement, and  $s_2$  the point in which the sphere is cut by the line along which the light comes from the star, produced backwards,  $s_2$  being corrected in the same manner as  $s_1$ . It is shown by observation that  $s_1$  is displaced from  $s$  towards the point towards which the earth is moving, through an angle equal to the ratio of the velocity of the earth to that of light multiplied by the sine of the earth's way. This is the phænomenon which I understand by the *aberration of light*, and which it was the object of one of my former communi-

\* Communicated by the Author.

cations to account for on the theory of undulations. But it is evident that what Prof. Challis means by aberration, is the circumstance that  $s_1$  is displaced from  $s_2$  through the angle which I have mentioned. Prof. Challis's reasoning, by his own confession, does not explain aberration in the sense in which I used the word; for he says that it follows from *observation* (not theory alone), that  $s_2$  coincides with  $s$ .

---

### LIII. *Intelligence and Miscellaneous Articles.*

#### ANALYSIS OF DIASPORE FROM SIBERIA.

BY M. A. DAMOUR.

THE remarkable characters of diaspore have frequently attracted the attention of mineralogists, and have been extremely well described and analysed by MM. Children, Dufrenoy, and Hess. The author observes, that he should therefore have abstained from referring to them, if he had not had occasion lately to observe a singular property of this mineral which had not been previously noticed. The diaspore is a well-known hydrate of alumina. It is shown by the experiments of M. Dufrenoy, that this mineral, even when long boiled in sulphuric acid, not only resists its action, but retains all its water. M. Damour, on repeating this experiment, obtained the same result; but he afterwards found that the diaspore, when deprived of its water by calcination, was almost totally soluble in sulphuric acid when assisted by heat.

This property is the inverse of that which chemists always observe with respect to hydrates, and in general with respect to substances which have not been calcined. In fact, the greater number of these substances lose their solubility in acids after they have been heated to redness. In this case the contrary occurs: the peculiar molecular condition of the crystallized hydrate of alumina, constituting the diaspore, appears then to be the only obstacle to the natural affinity of this hydrate for the sulphuric acid; for calcination, by destroying this arrangement of the molecules, restores the usual properties of alumina.

M. Damour took advantage of this circumstance in order to simplify the method of analysing diaspore.

The mineral was first purified by digesting it, reduced to very fine powder in dilute hydrochloric acid at a moderate heat. There was dissolved a notable quantity of oxide of iron accidentally mixed with it. The powder, after washing, was perfectly white. The proportion of water was found to be nearly similar in three different operations: to determine this the dried powder of the mineral was suffered to remain under a receiver over a stratum of pumice moistened with sulphuric acid, this powder was weighed and placed in a small covered platina crucible; in order to prevent the projection of the powdered mineral, the crucible was placed in another of the same metal; the whole being weighed, the crucibles were submitted to

the highest temperature which could be produced by the flame of an alcohol eolipyle. The crucibles were cooled in a receiver with a glass stopper, containing fragments of chloride of calcium. When perfectly cool they were again weighed, and the difference between the first weighing and that after calcination was attributed to the quantity of water disengaged, and was 14.97, 14.96, and 14.90 in three experiments.

In order to act upon the diaspoire deprived of water, hydrated sulphuric acid was poured upon the mineral remaining in the crucible in which it had been calcined. The whole was heated in a sand-bath so as to volatilize the greater part of the sulphuric acid; when the matter had become of a pasty consistence, water was added, which dissolved a great quantity of sulphate of alumina; the solution was poured off, and more sulphuric acid was added, and this operation was repeated five times. The aluminous solution was filtered in order to separate a small portion of a white earthy deposit; this, which had resisted the prolonged action of sulphuric acid, still contained much alumina; when moistened with nitrate of cobalt and heated to redness, it acquired a very decided blue tint, and readily dissolved in the salt of phosphorus.

The solution of sulphate of alumina was supersaturated with carbonate of ammonia; the alumina was collected, washed and heated for a long time to strong redness. It was very white, and nitrate of cobalt gave a fine blue tint to it.

One hundred parts of diaspoire yielded—

Alumina .....	79.91
Water .....	14.90
Mineral unacted upon ....	5.80—100.61

M. Damour admits that this analysis is superfluous after those of MM. Dufrenoy and Hess, and gives it merely to exhibit a property worthy of attention, and which had not been previously noticed with respect to any mineral whatever.—*Ann. de Ch. et de Phys.*, Mars 1846.

#### ON BORACIC ÆTHER.

M. Ebelmen having ascertained that boracic acid is volatilized by the vapour of water and of alcohol, succeeded in preparing, after some trials, boracic æther by the following process:—fused and finely-powdered boracic acid was put into a tubulated retort, and an equal weight of absolute alcohol was added to it. In a few minutes the temperature of the mixture became 122° Fahr., that of the atmosphere being only 64°. The retort was heated, and a thermometer placed in it showed that the liquid did not begin to boil until heated to about 203°, and its temperature continued rising from this point. At about 230° the distillation was stopped to cohobate the distilled liquid, and it was again distilled at 230°. The boracic acid swelled much during the operation, and the liquid which covered it while the distillation was going on, had completely imbibed it the following day. The distilled liquid had the slightly alliaceous smell of absolute alcohol, became very turbid on admixture with water, deposited

*Phil. Mag.* S. 3. Vol. 28. No. 187. April 1846. 2 A

boracic acid, burnt with a perfectly green flame, and yielded abundant white fumes of boracic acid.

The semi-solid mass remaining in the retort was bruised and digested during twenty-four hours in anhydrous æther, which completely disintegrated it; the æthereal solution, when clear, was poured into a retort placed in an oil-bath, and fitted with a condensing apparatus. It was requisite to employ a temperature of about 392° Fahr. to obtain the last traces of æther and of alcohol. There remained in the retort a large quantity of a viscid liquid, of a slight amber colour, yielding at 392° Fahr. thick vapours in contact with the air, and which became solid on cooling.

The author considers this product as boracic æther, and it approximates in physical properties to boracic acid and the borates, which are well-known to assume the vitreous state by fusion. It is in fact a true transparent glass, but one which is rather soft at common temperatures; at about 104° Fahr. it may be drawn into fine threads. It has a weak æthereal odour and burning taste; when applied to the skin, it occasions a strong sensation of heat, and is converted into a white powder, which is hydrated boracic acid; the same effect is produced by the contact of air with the boracic æther, but when the fragments are of considerable size, it takes place slowly, eventually however they become quite opaque. When boracic æther is triturated with water, it is very rapidly decomposed with the extrication of much heat; alcohol is reproduced and may be obtained by distilling the aqueous liquid.

Boracic æther is volatile, but not distillable; at about 392° Fahr. it emits thick vapours into the air; but when distillation is attempted, it is decomposed, leaving a considerable residue of fused boracic acid. When it is dissolved in absolute alcohol and the mixture distilled, the alcohol volatilizes such a quantity of boracic æther, that on the addition of water it becomes almost a solid mass.

It is combustible, and burns with a white smoke and a fine green flame, leaving a residue of fused boracic acid. It is soluble in æther and alcohol in all proportions, and retains these fluids with great affinity, for it is requisite to heat them to 392° Fahr. to remove the last traces of them; these solutions become solid masses by the addition of water.

When boracic æther is heated, it first fuses, then decomposes, swells, and becomes less and less liquid. There are simultaneously disengaged alcohol, which retains a large quantity of boracic æther, and a colourless gas which burns with a green flame before it is washed in water. After having been passed through water, the gas burns with a bright flame, and possesses all the properties of olefiant gas. The residue of the decomposition is but unmixed anhydrous boracic acid, much swelled with carbonaceous matter; it is requisite to heat it to redness for a long time to expel the inflammable gas.

Great difficulty attends the analysis of boracic æther; it was effected by converting the boracic acid, first into borate of ammonia, and afterwards into anhydrous boracic acid. The mean of several experiments gave—



Boric acid.....	66·7
Carbon .....	19·8
Hydrogen .....	4·4
Oxygen (estimated by loss) .	9·1—100·0

The author observes that the carbon and hydrogen are in the same proportions as in æther, while the oxygen is obviously in excess. The formula  $\text{BO}^6\text{C}^4\text{H}^5\text{O}$  is the nearest approach to the above mean; it gives—

Boric acid ..	872·0	65·4
Carbon .....	300·0	22·4
Hydrogen ....	62·5	4·7
Oxygen .....	100·0	7·6
	<u>1334·5</u>	<u>100·0</u>

M. Ebelmen observes that the difference between the results of experiment and calculation are too considerable to be attributed to errors of analysis. It must be admitted, he says, that the boracic æther contains a certain excess of boracic acid disseminated uniformly throughout the vitreous mass; this supposition, he further observes, is not at all improbable, when the mode of preparing boracic æther is considered.—*Ann. de Ch. et de Phys.*, Fevrier 1846.

#### ACTION OF BORACIC ACID ON PYROXYLIC SPIRIT.

M. Ebelmen states that the action of boracic acid upon pyroxylic spirit is similar to that which it exerts upon alcohol; when equal weights of them are mixed, great increase of temperature is produced. On heating the retort from  $212^\circ$  to  $230^\circ$  Fahr., but little distilled product is obtained; on allowing the retort to cool, and treating the matter which it contains with anhydrous æther, and operating in other respects as for boracic æther of alcohol, boracic methylic æther is obtained, the properties of which are perfectly comparable with those of boracic æther. It is soft and may be drawn into threads at common temperatures; when treated with water it is immediately decomposed, with the disengagement of much heat, into boracic acid and pyroxylic spirit; it burns like boracic æther, with a fine green flame.

Pyroxylic spirit is preferable to alcohol as a reagent for determining the presence of boracic acid by the colour of the flame; when the alcoholic solution does not contain much boracic acid, the edges only of the flame are green, and it is often difficult to discover it. But with pyroxylic spirit it requires only a small quantity of the acid to give the whole flame a green colour; this result is doubtless dependent upon the fact, that the flame of the pyroxylic spirit by itself has less colour than that of alcohol.

When pyroxylic spirit is distilled with a great excess of boracic acid, a colourless gas is obtained which is soluble in water, and whose properties resemble those of boracic methylic æther,  $\text{C}^2\text{H}^3\text{O}$ ; the mode in which boracic methylic æther is decomposed is therefore entirely different from that of the corresponding compound of alcohol.

M. Ebelmen found that boracic methylic æther yielded 69·5 and 70·6 per cent. of fused boracic acid by ammonia; the acid was black and contained a small quantity of charcoal disseminated through it;

the proportion of acid correspondent to the formula  $\text{BO}^6 \text{C}^2 \text{H}^3 \text{O}$  would be 75.2 per cent. The product obtained was evidently a little impure, and contained, besides some boracic methylic æther, some of the compounds of boracic acid with the pyrogenous compounds, which it is so difficult to separate from pyroxylic spirit.—*Ibid.*

ON A SIMPLE METHOD OF PROTECTING FROM LIGHTNING,  
BUILDINGS WITH METALLIC ROOFS. BY PROF. HENRY.

On the principle of electrical induction, houses thus covered are evidently more liable to be struck than those furnished either with shingle or tile. Fortunately, however, they admit of very simple means of perfect protection. It is evident, from well-established principles of electrical action, that if the outside of a house were encased entirely in a coating of metal, the most violent discharge which might fall upon it from the clouds would pass silently to the earth without damaging the house, or endangering the inmates. It is also evident, that if the house be merely covered with a roof of metal, without projecting chimneys, and this roof were put in metallic connexion with the ground, the building would be perfectly protected. To make a protection, therefore, of this kind, the Professor advises that the metallic roof be placed in connexion with the ground, by means of the tin or copper gutters which serve to lead the water from the roof to the earth. For this purpose, it is sufficient to solder to the lower end of the gutter a riband of sheet copper, two or three inches wide, surrounding it with charcoal, and continuing it out from the house until it terminates in moist ground. The upper ends of these gutters are generally soldered to the roof; but if they are not in metallic contact, the two should be joined by a slip of sheet copper. The only part of the house unprotected by this arrangement will be the chimneys; and to secure these, it will only be necessary to erect a short rod against the chimney, soldered at its lower end to the metal of the roof, and extending fifteen or twenty inches above the top of the flue.

Considerable discussion in late years has taken place in reference to the transmission of electricity along a conductor; whether it passes through the whole capacity of the rod, or is principally confined to the surface. From a series of experiments presented to the American Philosophical Society, by Professor Henry, on this subject, it appears that the electrical discharge passes, or tends to pass, principally at the surface; and as an ordinary-sized house is commonly furnished with from two to four perpendicular gutters (two in front and two in the rear), the surface of these will be sufficient to conduct, silently, the most violent discharge which may fall from the clouds.

Professor Henry also stated, that he had lately examined a house struck by lightning, which exhibited some effects of an interesting kind. The lightning struck the top of the chimney, passed down the interior of the flue to a point opposite a mass of iron placed on the floor of the garret, where it pierced the chimney; thence it passed explosively, breaking the plaster, into a bedroom below, where it came in contact with a copper bell-wire, and passed along this horizontally and silently for about six feet; thence it leaped explosively

through the air a distance of about ten feet, through a dormer window, breaking the sash, and scattering the fragments across the street. It was evidently attracted to this point by the upper end of a perpendicular gutter, which was near the window. It passed silently down the gutter, exhibiting scarcely any mark of its passage until it arrived at the termination, about a foot from the ground. Here again an explosion appeared to have taken place, since the windows of the cellar were broken. A bed, in which a man was sleeping at the time, was situated against the wall, immediately under the bell-wire; and although his body was parallel to the wire, and not distant from it more than four feet, he was not only uninjured, but not sensibly affected. The size of the hole in the chimney, and the fact that the lightning passed along the copper wire without melting it, show that the discharge was a small one, and yet the mechanical effects, in breaking the plaster, and projecting the window-frame across the street, were astonishingly great.

These effects the Professor attributes to a sudden repulsive energy, or expansive force developed in the air along the path of the discharge. Indeed, he conceives that most of the mechanical effects which are often witnessed in cases of buildings struck by lightning, may be referred to the same cause. In the case of a house struck within a few miles of Princeton, the discharge entered the chimney, burst open the flue, and passed along the *cockloft* to the other end of the house; and such was the explosive force in this confined space, that nearly the whole roof was blown off. This effect was, in all probability, due to the same cause which suddenly expands the air in the experiment with Kinnorsly's electrical air thermometer.—From the *Proc. of the American Philosophical Society*, June 20, 1845.

#### OBSERVATIONS ON CAPILLARITY. BY PROF. HENRY.

In 1839, the author presented the results of some experiments on the permeability of lead to mercury; and subsequent observation had led him to believe that the same property was possessed by other metals in reference to each other. His first attempt to verify this conjecture was made with the assistance of Dr. Patterson, at the United States Mint. For this purpose, a small globule of gold was placed on a plate of sheet iron, and submitted to the heat of an assaying furnace; but the experiment was unsuccessful; for, although the gold was heated much above its melting-point, it exhibited no signs of sinking into the pores of the iron. The idea afterward suggested itself, that a different result would have been obtained had the two metals been made to adhere previous to heating, so that no oxide could have been formed between the surfaces. In accordance with this view, Prof. Henry inquired of Mr. Cornelius, of Philadelphia, if, in the course of his experience in working silver-plated copper, in his extensive manufactory of lamps, he had ever observed the silver to disappear from the copper when the metal was heated. The answer was, that the silver always disappears when the plate is heated above a certain temperature, leaving a surface of copper exposed; and that it was generally believed by the workmen, that the silver evaporates at this temperature.

Professor Henry suggested that the silver, instead of evaporating, merely sunk into the pores of the copper, and that by carefully removing the surface of the latter by the action of an acid, the silver would reappear. To verify this by experiment, Mr. Cornelius heated one end of a piece of thick plated copper to nearly the melting-point of the metal; the silver at this end disappeared, and when the metal was cleaned by a solution of dilute sulphuric acid, the end which had been heated presented a uniform surface of copper, whilst the other end exhibited its proper coating of silver. The unsilvered end of the plate was next placed, for a few minutes, in a solution of muriate of zinc, by which the exterior surface of copper was removed, and the surface of silver was again exposed. This method of recovering the silver before the process of plating silver by galvanism came into use, would have been of much value to manufacturers of plated ware, since it often happened that articles were spoiled, in the process of soldering, by heating them to the degree at which silver disappears.

It is well-known to the jeweller, that articles of copper, plated with gold, lose their brilliancy after a time, and that this can be restored by boiling them in ammonia; this effect is probably produced by the ammonia acting on the copper, and dissolving off its surface, so as to expose the gold, which, by diffusion, has entered into the copper.

A slow diffusion of one metal through another probably takes place in cases of alloys. Silver coins, after having lain long in the earth, have been found covered with a salt of copper. This may be explained by supposing that the alloy of copper, at the surface of the coin, enters into combination with the carbonic acid of the soil, and being thus removed, its place is supplied by a diffusion from within; and in this way it is not improbable that a considerable portion of the alloy may be exhausted in the process of time, and the purity of the coin be considerably increased.

Perhaps, also, the phænomenon of what is called *segregation*, or the formation of nodules of flint in masses of carbonated lime, and of indurated marl in beds of clay, may be explained on the same principle. In breaking up these masses, it is almost always observed, that a piece of shell or some extraneous matter occupies the middle, and probably formed the nucleus, around which the matter was accumulated by attraction. The difficulty consists in explaining how the attraction of cohesion, which becomes insensible at sensible distances, should produce this effect. To explain this, let us suppose two substances uniformly diffused through each other by a slight mutual attraction, as in the case of a lump of sugar dissolved in a large quantity of water, every particle of the water will attract to itself its proportion of the sugar, and the whole will be in a state of equilibrium. If the diffusion at its commencement had been assisted by heat, and this cause of the separation of the homogeneous particles no longer existed, the diffusion might be one of unstable equilibrium; and the slightest extraneous force, such as the attraction of a minute piece of shell, might serve to disturb the quiescence, and draw to itself the diffused particles which were immediately contiguous to it. This would leave a vacuum of the atoms around the attracting mass: for example, as in the case of the sugar, there would be a portion of the

water around the nucleus deprived of the sugar; this portion of the water would attract its portion of sugar from the layer without, and into this layer the sugar from the layer next without would be diffused, and so on until, through all the water, the remaining sugar would be uniformly diffused. The process would continue to be repeated, by the nucleus again attracting a portion of the sugar from the water immediately around it, and so on until a considerable accumulation would be formed around the foreign substance.

We can in this way conceive of the manner by which the molecular action, which is insensible at perceptible distances, may produce results which would appear to be the effect of attraction acting at a distance.—From the *Proc. of the American Philosophical Society*.

OBITUARY.—The University of Königsberg has sustained a severe loss by the death of the celebrated astronomer Bessel, who died, after long suffering, on the 17th of March, in the 62nd year of his age.

METEOROLOGICAL OBSERVATIONS FOR FEB. 1846.

*Chiswick*.—February 1. Very fine: rain. 2. Fine. 3, 4. Overcast: rain. 5, 6. Very fine. 7. Overcast: windy, with showers. 8. Clear: cloudy: very clear at night. 9. Frosty: fine, but cold. 10. Frosty: cloudy and cold. 11. Frosty: fine: partially overcast. 12. Foggy: cloudy and fine. 13. Densely clouded. 14, 15. Cloudy and fine. 16. Densely overcast. 17, 18. Overcast and fine. 19. Hazy. 20. Overcast. 21. Exceedingly fine. 22. Cloudy: boisterous, with rain at night. 23, 24. Rain. 25. Heavy clouds and mild. 26. Cloudy in the morning: afterward cloudless and exceedingly fine. 27. Slight haze: showery. 28. Very fine.

Mean temperature of the month .....	43°·32
Mean temperature of February 1845 .....	33·07
Average mean temperature for the last twenty years .....	39·36
Average amount of rain.....	1·61 inch.

*Boston*.—Feb. 1. Fine. 2. Fine: rain early A.M. 3. Cloudy. 4. Fine. 5. Cloudy: rain early A.M. 6. Fine. 7. Stormy: rain early A.M. 8. Fine: rain early A.M. 9. Fine: snow early A.M.: snow A.M. and P.M. 10. Fine: snow on the ground. 11. Cloudy: snow on the ground. 12. Fine: snow on the ground. 13. Cloudy: snow all gone: melted snow. 14—22. Cloudy. 23. Cloudy: rain early A.M. 24. Cloudy. 25. Fine: rain early A.M. 26. Cloudy. 27. Fine: rain A.M. 28. Fine. This month has been usually fine.

*Sandwich Manse, Orkney*.—Feb. 1. Sleet-showers. 2. Cloudy. 3. Cloudy: sleet-showers. 4, 5. Hail-showers. 6. Showers: rain. 7. Showers: snow-showers. 8. Snow-showers. 9. Snow-showers: frost. 10. Snow: showers. 11. Clear: cloudy. 12. Cloudy: damp. 13. Showers. 14, 15. Cloudy: showers. 16. Rain: cloudy. 17. Showers: cloudy: drizzle. 18. Showers: drizzle: cloudy: drizzle. 19. Bright: cloudy. 20. Clear: cloudy. 21. Rain: cloudy. 22. Rain. 23. Clear. 24. Damp: showers. 25—27. Clear: cloudy. 28. Cloudy: showers: clear.

*Applegarth Manse, Dumfries-shire*.—Feb. 1. Occasional showers. 2. Fair and fine. 3. Heavy rain. 4. Sleet and rain P.M. 5. Showers. 6, 7. Heavy showers. 8. Slight fall of snow. 9. Frost: fine: clear. 10. Frost: fine. 11. Thaw: fair: mild. 12. Slight frost. 13. Very slight frost. 14, 15. Fine. 16. Very fine. 17. Fine. 18. Frost A.M. 19. Fine, but cloudy: shower. 20. Slight showers: mild. 21. Wet and stormy. 22, 23. Damp and drizzling. 24, 25. Heavy rain. 26. Wet. 27. Remarkably fine. 28. Damp and drizzling.

Mean temperature of the month .....	43°·4
Mean temperature of February 1845 .....	34·5
Mean temperature of Feb. for twenty-three years .	37·0
Mean rain in February for eighteen years.....	2·0 inches.

*Meteorological Observations made by Mr. Thompson at the Garden of the Horticultural Society at CHISWICK, near London; by Mr. Veall, at BOSTON; by the Rev. W. Dunbar, at Applegarth Manse, DUMFRIES-SHIRE; and by the Rev. C. Clouston, at Sandwick Manse, ORKNEY.*

Days of Month.	Barometer.						Thermometer.						Wind.						Rain.								
	Chiswick.		Dumfries-shire.		Orkney, Sandwick.		Chiswick.		Boston.		Dumfries-shire.		Orkney, Sandwick.		Boston.		Dumfries-shire.		Orkney, Sandwick.		Boston.		Chiswick.		Orkney, Sandwick.		
	Max.	Min.	8 $\frac{1}{2}$ a.m.	9 a.m.	9 p.m.	8 $\frac{1}{2}$ p.m.	Max.	Min.	8 $\frac{1}{2}$ a.m.	9 a.m.	9 p.m.	8 $\frac{1}{2}$ a.m.	9 a.m.	9 p.m.	Max.	Min.	8 $\frac{1}{2}$ a.m.	9 a.m.	9 p.m.	8 $\frac{1}{2}$ a.m.	9 a.m.	9 p.m.	Max.	Min.	8 $\frac{1}{2}$ a.m.	9 a.m.	9 p.m.
1846.																											
Feb.																											
1.	29.834	29.830	29.35	29.53	29.55	29.41	51	39	44	50 $\frac{1}{2}$	41	43	40 $\frac{1}{2}$	43	40 $\frac{1}{2}$	nw.	w.	wnw.	wnw.	wnw.	wnw.	wnw.	wnw.	.52	.20	.20	.06
2.	29.932	29.614	29.28	29.58	29.68	29.61	48	35	38	48	39 $\frac{1}{2}$	45	47	41	37	nw.	w.	ws.	w.	w.	w.	w.	w.	.06	.06	.06	.06
3.	29.894	29.821	29.49	29.50	29.38	29.24	52	39	43	49	42	45	43	45	37	sw.	w.	w.	w.	w.	w.	w.	w.	.01	.15	.15	.06
4.	30.053	29.956	29.52	29.65	29.67	29.41	49	36	40.5	43 $\frac{1}{2}$	35 $\frac{1}{2}$	38	42	38	42	w.	w.	w.	w.	w.	w.	w.	w.	.31	.57	.57	.06
5.	29.833	29.744	29.42	29.50	29.60	29.40	49	30	41	44	38	40	38	40	38	nw.	w.	w.	w.	w.	w.	w.	w.	.04	.16	.16	.06
6.	29.988	29.937	29.53	29.68	29.46	29.44	50	33	38	46 $\frac{1}{2}$	35	37 $\frac{1}{2}$	44 $\frac{1}{2}$	37	44 $\frac{1}{2}$	w.	w.	ws.	ws.	ws.	ws.	ws.	ws.	.08	.17	.17	.06
7.	29.994	29.829	29.26	29.29	29.69	29.58	52	31	48	48 $\frac{1}{2}$	43 $\frac{1}{2}$	41 $\frac{1}{2}$	34	41	34	w.	w.	wnw.	wnw.	wnw.	wnw.	wnw.	wnw.	.08	.38	.38	.06
8.	30.129	29.979	29.54	29.79	30.00	29.83	46	29	40	41	33 $\frac{1}{2}$	34	34	34	34	w.	w.	wnw.	nne.	nne.	nne.	nne.	nne.	.04	.14	.14	.06
9.	30.341	30.113	29.77	30.18	30.23	30.44	45	26	33	49	32	34 $\frac{1}{2}$	33 $\frac{1}{2}$	33	33	ne.	calm	n.	n.	n.	n.	n.	n.	.04	.24	.24	.06
10.	30.341	30.070	30.05	30.31	30.22	30.31	41	22	32	40	29 $\frac{1}{2}$	39	44	41	43 $\frac{1}{2}$	ne.	calm	n.	n.	n.	n.	n.	n.	.04	.03	.03	.06
11.	30.158	30.074	29.78	30.00	30.00	30.05	44	25	34	46 $\frac{1}{2}$	35	41	43 $\frac{1}{2}$	46	46	nw.	calm	nw.	nw.	nw.	nw.	nw.	nw.	.04	.02	.02	.06
12.	30.142	30.118	29.83	30.03	29.99	29.99	45	36	34	50	33 $\frac{1}{2}$	46	46	46	46	w.	calm	nw.	nw.	nw.	nw.	nw.	nw.	.25	.06	.06	.06
13.	30.137	30.105	29.73	29.96	29.99	29.91	45	27	40.5	49	34	45	43 $\frac{1}{2}$	38	45	w.	calm	nw.	nw.	nw.	nw.	nw.	nw.	.00	.06	.06	.06
14.	30.185	30.118	29.70	30.01	30.10	30.06	48	24	39	49	37 $\frac{1}{2}$	42 $\frac{1}{2}$	38	42	38	w.	calm	w.	w.	w.	w.	w.	w.	.00	.13	.13	.06
15.	30.260	30.227	29.90	30.12	30.07	30.02	50	31	38	48	33	45	46	46	46	w.	calm	w.	w.	w.	w.	w.	w.	.00	.13	.13	.06
16.	30.194	30.182	29.80	30.09	30.09	30.12	48	41	43	51	41 $\frac{1}{2}$	46	45	45	44 $\frac{1}{2}$	nw.	calm	w.	w.	w.	w.	w.	w.	.00	.05	.05	.06
17.	30.177	30.010	29.77	30.05	29.93	30.04	47	39	41	49 $\frac{1}{2}$	39 $\frac{1}{2}$	44 $\frac{1}{2}$	45	44	45	w.	calm	nw.	nw.	nw.	nw.	nw.	nw.	.00	.05	.05	.06
18.	30.005	29.978	29.62	29.82	29.85	29.85	48	39	42	47 $\frac{1}{2}$	31	45	44 $\frac{1}{2}$	45	44 $\frac{1}{2}$	w.	calm	n.	w.	w.	w.	w.	w.	.00	.03	.03	.06
19.	30.061	29.993	29.62	29.82	29.85	29.85	46	39	42	45	39	45	45	45	45	sw.	calm	w.	w.	w.	w.	w.	w.	.00	.02	.02	.06
20.	30.095	30.074	29.72	29.88	29.82	29.91	50	40	42	47	40	43 $\frac{1}{2}$	42	46	46	sw.	calm	s.	w.	w.	w.	w.	w.	.01	.10	.10	.06
21.	30.109	30.091	29.69	29.78	29.80	29.72	58	36	45.5	48 $\frac{1}{2}$	45	48	46	46	46	sw.	calm	s.	w.	w.	w.	w.	w.	.01	.15	.15	.06
22.	29.988	29.861	29.55	29.50	29.50	29.42	57	47	47	53	42 $\frac{1}{2}$	47	47	47	47	s.	calm	s.	w.	w.	w.	w.	w.	.07	.24	.24	.06
23.	29.800	29.690	29.38	29.49	29.45	29.41	58	50	52.5	51	47 $\frac{1}{2}$	50	45	45	45	sw.	calm	s.	w.	w.	w.	w.	w.	.16	.22	.22	.06
24.	29.603	29.405	29.15	29.18	29.08	29.27	59	49	52.5	55 $\frac{1}{2}$	48	46	46	46	46	sw.	calm	sw.	sw.	sw.	sw.	sw.	sw.	.03	.22	.22	.06
25.	29.801	29.489	29.00	28.93	29.43	28.99	58	34	52	54	47	47	46	46	46	sw.	w.	sw.	sw.	sw.	sw.	sw.	sw.	.01	.22	.22	.06
26.	29.800	29.663	29.37	29.46	29.36	29.48	60	39	45	52	42	46	46	46	46	s.	w.	w.	w.	w.	w.	w.	w.	.01	.03	.03	.06
27.	29.648	29.623	29.30	29.40	29.44	29.45	62	38	49	56 $\frac{1}{2}$	46	47 $\frac{1}{2}$	47	47	47	se.	w.	w.	w.	w.	w.	w.	w.	.05	.03	.03	.06
28.	29.952	29.664	29.23	29.47	29.62	29.51	64	42	52	52	41 $\frac{1}{2}$	47	46	46	46	w.	w.	w.	w.	w.	w.	w.	w.	.02	.02	.02	.06
Mean.	30.016	29.902	29.54	29.701	29.775	29.681	51.07	35.07	42.3	48.8	39.0	43.30	42.80	43.30	42.80									1.61	0.57	1.52	3.34

THE  
LONDON, EDINBURGH AND DUBLIN  
PHILOSOPHICAL MAGAZINE  
AND  
JOURNAL OF SCIENCE.

[THIRD SERIES.]

MAY 1846.

LIV. *Thoughts on Ray-vibrations.* By MICHAEL FARADAY,  
Esq., D.C.L., F.R.S., Fullerian Prof., &c. &c.

*To Richard Phillips, Esq.*

DEAR SIR,

AT your request I will endeavour to convey to you a notion of that which I ventured to say at the close of the last Friday-evening Meeting, incidental to the account I gave of Wheatstone's electro-magnetic chronoscope; but from first to last understand that I merely threw out as matter for speculation, the vague impressions of my mind, for I gave nothing as the result of sufficient consideration, or as the settled conviction, or even probable conclusion at which I had arrived.

The point intended to be set forth for the consideration of the hearers was, whether it was not possible that the vibrations which in a certain theory are assumed to account for radiation and radiant phenomena may not occur in the lines of force which connect particles, and consequently masses of matter together; a notion which, as far as it is admitted, will dispense with the æther which, in another view, is supposed to be the medium in which these vibrations take place.

You are aware of the speculation\* which I some time since uttered respecting that view of the nature of matter which considers its ultimate atoms as centres of force, and not as so many little bodies surrounded by forces, the bodies being considered in the abstract as independent of the forces and capable of existing without them. In the latter view, these little particles have a definite form and a certain limited size; in the former view such is not the case, for that which represents size may be considered as extending to any distance to which the lines of force of the particle extend: the particle indeed is

\* Phil. Mag. 1844, vol. xxiv. p. 136.

supposed to exist only by these forces, and where they are it is. The consideration of matter under this view gradually led me to look at the lines of force as being perhaps the seat of the vibrations of radiant phænomena.

Another consideration bearing conjointly on the hypothetical view both of matter and radiation, arises from the comparison of the velocities with which the radiant action and certain powers of matter are transmitted. The velocity of light through space is about 190,000 miles in a second; the velocity of electricity is, by the experiments of Wheatstone, shown to be as great as this, if not greater: the light is supposed to be transmitted by vibrations through an æther which is, so to speak, destitute of gravitation, but infinite in elasticity; the electricity is transmitted through a small metallic wire, and is often viewed as transmitted by vibrations also. That the electric transference depends on the forces or powers of the matter of the wire can hardly be doubted, when we consider the different conductivity of the various metallic and other bodies; the means of affecting it by heat or cold; the way in which conducting bodies by combination enter into the constitution of non-conducting substances, and the contrary; and the actual existence of one elementary body, carbon, both in the conducting and non-conducting state. The power of electric conduction (being a transmission of force equal in velocity to that of light) appears to be tied up in and dependent upon the properties of the matter, and is, as it were, existent in them.

I suppose we may compare together the matter of the æther and ordinary matter (as, for instance, the copper of the wire through which the electricity is conducted), and consider them as alike in their essential constitution; *i. e.* either as both composed of little nuclei, considered in the abstract as matter, and of force or power associated with these nuclei, or else both consisting of mere centres of force, according to Boscovich's theory and the view put forth in my speculation; for there is no reason to assume that the nuclei are more requisite in the one case than in the other. It is true that the copper gravitates and the æther does not, and that therefore the copper is ponderable and the æther is not; but that cannot indicate the presence of nuclei in the copper more than in the æther, for of all the powers of matter gravitation is the one in which the force extends to the greatest possible distance from the supposed nucleus, being infinite in relation to the size of the latter, and reducing that nucleus to a mere centre of force. The smallest atom of matter on the earth acts directly on the smallest atom of matter in the sun, though they are 95,000,000



of miles apart; further, atoms which, to our knowledge, are at least nineteen times that distance, and indeed, in cometary masses, far more, are in a similar way tied together by the lines of force extending from and belonging to each. What is there in the condition of the particles of the supposed æther, if there be even only *one* such particle between us and the sun, that can in subtilty and extent compare to this?

Let us not be confused by the *ponderability* and *gravitation* of heavy matter, as if they proved the presence of the abstract nuclei; these are due not to the nuclei, but to the force superadded to them, if the nuclei exist at all; and, if the *æther* particles be without this force, which according to the assumption is the case, then they are more material, in the abstract sense, than the matter of this our globe; for matter, according to the assumption, being made up of nuclei and force, the æther particles have in this respect proportionately more of the nucleus and less of the force.

On the other hand, the infinite elasticity assumed as belonging to the particles of the æther, is as striking and positive a force of it as gravity is of ponderable particles, and produces in its way effects as great; in witness whereof we have all the varieties of radiant agency as exhibited in luminous, calorific, and actinic phænomena.

Perhaps I am in error in thinking the idea generally formed of the æther is that its nuclei are almost infinitely small, and that such force as it has, namely its elasticity, is almost infinitely intense. But if such be the received notion, what then is left in the æther but force or centres of force? As gravitation and solidity do not belong to it, perhaps many may admit this conclusion; but what is gravitation and solidity? certainly not the weight and contact of the abstract nuclei. The one is the consequence of an *attractive* force, which can act at distances as great as the mind of man can estimate or conceive; and the other is the consequence of a *repulsive* force, which forbids for ever the contact or touch of any two nuclei; so that these powers or properties should not in any degree lead those persons who conceive of the æther as a thing consisting of force only, to think any otherways of ponderable matter, except that it has more and other *forces* associated with it than the æther has.

In experimental philosophy we can, by the phænomena presented, recognise various kinds of lines of force; thus there are the lines of gravitating force, those of electro-static induction, those of magnetic action, and others partaking of a dynamic character might be perhaps included. The lines of electric and magnetic action are by many considered as exerted

through space like the lines of gravitating force. For my own part, I incline to believe that when there are intervening particles of matter (being themselves only centres of force), they take part in carrying on the force through the line, but that when there are none, the line proceeds through space\*. Whatever the view adopted respecting them may be, we can, at all events, affect these lines of force in a manner which may be conceived as partaking of the nature of a shake or lateral vibration. For suppose two bodies, A B, distant from each other and under mutual action, and therefore connected by lines of force, and let us fix our attention upon one resultant of force having an invariable direction as regards space; if one of the bodies move in the least degree right or left, or if its power be shifted for a moment within the mass (neither of these cases being difficult to realize if A and B be either electric or magnetic bodies), then an effect equivalent to a lateral disturbance will take place in the resultant upon which we are fixing our attention; for, either it will increase in force whilst the neighbouring resultants are diminishing, or it will fall in force as they are increasing.

It may be asked, what lines of force are there in nature which are fitted to convey such an action and supply for the vibrating theory the place of the æther? I do not pretend to answer this question with any confidence; all I can say is, that I do not perceive in any part of space, whether (to use the common phrase) vacant or filled with matter, anything but forces and the lines in which they are exerted. The lines of weight or gravitating force are, certainly, extensive enough to answer in this respect any demand made upon them by radiant phænomena; and so, probably, are the lines of magnetic force: and then who can forget that Mossotti has shown that gravitation, aggregation, electric force, and electro-chemical action may all have one common connexion or origin; and so, in their actions at a distance, may have in common that infinite scope which some of these actions are known to possess?

The view which I am so bold as to put forth considers, therefore, radiation as a high species of vibration in the lines of force which are known to connect particles and also masses of matter together. It endeavours to dismiss the æther, but not the vibrations. The kind of vibration which, I believe, can alone account for the wonderful, varied, and beautiful phænomena of polarization, is not the same as that which occurs on the surface of disturbed water, or the waves of sound in gases or liquids, for the vibrations in these cases are direct,

\* Experimental Researches in Electricity, pars. 1161, 1613, 1663, 1710, 1729, 1735, 2443.

or to and from the centre of action, whereas the former are lateral. It seems to me, that the resultant of two or more lines of force is in an apt condition for that action which may be considered as equivalent to a *lateral* vibration; whereas an uniform medium, like the æther, does not appear apt, or more apt than air or water.

The occurrence of a change at one end of a line of force easily suggests a consequent change at the other. The propagation of light, and therefore probably of all radiant action, occupies *time*; and, that a vibration of the line of force should account for the phænomena of radiation, it is necessary that such vibration should occupy time also. I am not aware whether there are any data by which it has been, or could be ascertained whether such a power as gravitation acts without occupying time, or whether lines of force being already in existence, such a lateral disturbance of them at one end as I have suggested above, would require time, or must of necessity be felt instantly at the other end.

As to that condition of the lines of force which represents the assumed high elasticity of the æther, it cannot in this respect be deficient: the question here seems rather to be, whether the lines are sluggish enough in their action to render them equivalent to the æther in respect of the time known experimentally to be occupied in the transmission of radiant force.

The æther is assumed as pervading all bodies as well as space: in the view now set forth, it is the forces of the atomic centres which pervade (and make) all bodies, and also penetrate all space. As regards space, the difference is, that the æther presents successive parts or centres of action, and the present supposition only lines of action; as regards matter, the difference is, that the æther lies between the particles and so carries on the vibrations, whilst as respects the supposition, it is by the lines of force between the centres of the particles that the vibration is continued. As to the difference in intensity of action within matter under the two views, I suppose it will be very difficult to draw any conclusion, for when we take the simplest state of common matter and that which most nearly causes it to approximate to the condition of the æther, namely the state of rare gas, how soon do we find in its elasticity and the mutual repulsion of its particles, a departure from the law, that the action is inversely as the square of the distance!

And now, my dear Phillips, I must conclude. I do not think I should have allowed these notions to have escaped from me, had I not been led unawares, and without previous

consideration, by the circumstances of the Evening on which I had to appear suddenly and occupy the place of another. Now that I have put them on paper, I feel that I ought to have kept them much longer for study, consideration, and, perhaps, final rejection; and it is only because they are sure to go abroad in one way or another, in consequence of their utterance on that evening, that I give them a shape, if shape it may be called, in this reply to your inquiry. One thing is certain, that any hypothetical view of radiation which is likely to be received or retained as satisfactory, must not much longer comprehend alone certain phænomena of light, but must include those of heat and of actinic influence also, and even the conjoined phænomena of sensible heat and chemical power produced by them. In this respect, a view, which is in some degree founded upon the ordinary forces of matter, may perhaps find a little consideration amongst the other views that will probably arise. I think it likely that I have made many mistakes in the preceding pages, for even to myself, my ideas on this point appear only as the shadow of a speculation, or as one of those impressions on the mind which are allowable for a time as guides to thought and research. He who labours in experimental inquiries knows how numerous these are, and how often their apparent fitness and beauty vanish before the progress and development of real natural truth.

I am, my dear Phillips,

Ever truly yours,

M. FARADAY.

Royal Institution, April 15, 1846.

LV. *On the Wax of the Chamærops.*

By J. E. TESCHEMACHER, Esq.\*

**A**BOUT three millions of palm leaves are annually imported into the United States of America, for the purpose of being manufactured into hats. They come tied in bundles, called in Spanish *Esteras*, each *estera* weighing from 50 to 60 pounds; these are the palmate part of the leaf with a small portion of the petiole; this last weighs one-eighth of the leaf. The palm from which the leaves are cut in Cuba and other parts of the West Indies for this purpose is a Chamærops, a low-growing species, not differing I believe from the *C. humilis* of the southern sections of the United States, except in being much more robust in habit. The *C. humilis* of the United States is too soft and yielding for this manufacture.

\* Communicated by the Chemical Society; having been read December 1, 1845.

I have cultivated the plant from Cuba for five or six years, and was unable to discover any difference in foliage; but I have never seen the fruit of either. The leaf of the *Chamærops* spreads out nearly horizontal with folds, precisely like those of a lady's fan. On opening these folds, when they arrive in the United States in their dried state, there is a quantity of white flaky powder, under this is the bright varnish which covers the whole surface of the leaf; both these are true vegetable wax. From one of these palm leaves I obtained, by passing the thumb down the folds, 90 grains of the white wax in powdery flakes, and by boiling the leaf, after cutting in pieces, in alcohol, 300 grains more of a gray coloured wax.

At the manufactory the leaves are often bleached by the fumes of sulphurous acid gas, and then split by machinery into very thin strips; this division cracks off of course a large portion of the brittle varnish, which, together with the white powder, falls to the ground, is swept together and burnt or thrown away. The weight of this substance destroyed annually probably exceeds one hundred thousand pounds.

On treating this substance with a small quantity of boiling alcohol, it may, like other wax, be separated into cerine and myricine.

The powdery flakes first obtained contain about 80 per cent. myricine and 20 per cent. cerine, but the wax obtained from boiling the leaf in alcohol contains scarcely any myricine. This is easily accounted for; the flakes, being the brittle and more resinous part, break off readily; while the alcohol, which acts on the leaves, dissolves only the cerine, leaving the myricine undissolved; this might no doubt be obtained by increasing the quantity of alcohol and continuing the process, if it were desirable. In bees' wax the proportions of these two substances vary also, the cerine from 70 to 90 per cent., myricine from 10 to 30 per cent.; and it is probable that the more or less brittle quality of all wax depends on the relative quantity of these two ingredients.

The wax of *Ceroxylon andicola*, a very lofty palm, found by Humboldt at Quindin on the Andes, has been analysed, and found very nearly to resemble bees' wax in its ultimate principles.

	Bees' wax.	Palm wax.
Carbon . . . . .	80·14	80·28
Hydrogen . . . . .	14·08	13·20
Oxygen . . . . .	5·78	6·52

To obtain this wax, the outer portion of the trunk is rasped or scraped, the raspings are heated in water, the wax swims at the top, the other parts fall to the bottom, the wax is collected, made into small balls, and dried in the sun; it has a

deep yellow colour, and when the resinous part (myricine?) is melted it has the appearance of amber; after the separation of the wax and resin from the produce of Ceroxylon, there remains in the alcohol a bitter yellow substance, supposed to be a vegetable alkaloid. This yellow substance separates also from the wax of the leaf of *Chamærops*, but I think it is not an ingredient in the wax, but of other parts of the juices dissolved by the alcohol.

The production from the juices of plants by a purely vegetable function of wax scarcely differing from that deposited in their hives by bees is calculated to throw light on the question of the formation of this substance by these insects, and also merits the careful examination of those who are entering into the study of the various transformations of the vegetable juices at different periods of their progress towards maturity.

LVI. *Analysis of a Cobalt Ore found in Western India.*

By J. MIDDLETON, Esq., F.G.S.\*

**B**EING engaged in analyses of the metallic ores of North-western India, with a view to the ascertainment of the constitutions of those most remarkable among them, and also with the hope of detecting others whose existence in the country has not heretofore been even suspected, I am desirous of submitting to the Chemical Society the results of my inquiries whenever they appear to me of sufficient interest to justify my troubling them with them. I may mention, that should the Society desire any information from me on this or any other subject that I may be qualified and in a position to furnish, I shall most gladly meet their wishes.

The hilly districts of Rajpootanah are remarkably prolific in metallic ores, many of these, too, exceedingly rich and abundant. Within a narrow compass in the independent state of Syepoore, are to be found the following minerals:—

Sulphuret of copper, sulphate of copper, sulphuret of cobalt, alum.

The native method of mining for the first of these ores, and which is the same as that adopted for the others, may be interesting to some of your members.

“The mine of copper is very deep, and difficult of access. The miners enter with burning lamps on their heads and with chisels, iron hammers, and baskets in their hands. They dig out the ore with their chisels by the light of their lamps, and bring it up with great labour and difficulty to the surface. They then pound and grind it small in a mill, after which

\* Communicated by the Chemical Society; having been read December 15, 1845.

they mix it with moist cow-dung, and this mixture being made into balls is placed in the sun to dry. When this has been accomplished the lumps are burnt, after which they are not broken up, but being mixed with an equal quantity of charcoal and as much iron filings, are put into a crucible, and a strong heat kept up by blowing with a leathern bellows till the dross separates and the copper settles at the bottom in the form of a solid disc. This product is again heated with charcoal until perfectly pure copper is produced\*."

The mineral possessing greatest interest amongst those above enumerated is the sulphuret of cobalt. It is found in the copper mines in considerable abundance, and exists in a primitive schist in the form of bands and disseminated grains, the colour of which is a steel gray inclining to yellow. The grains appear to be crystallized, and are probably the cube and its derivatives. What is particularly remarkable in this ore is its purity, so far surpassing in this respect any that, so far as I am aware, is to be met with anywhere else. The only substance in combination with it, after separation of the matrix, is an iron pyrites, which is however but mechanically mixed, and so highly magnetic as to be readily removable by the magnet. The relative proportions in which these two exist are—

Cobalt pyrites . . . .	90·78 per cent.
Iron . . . . .	9·22 ...

The iron pyrites consists of black amorphous granules without metallic lustre, and, as above stated, it is highly magnetic, having at the same time the low specific gravity of 2·58. It gives on analysis—

Iron . . . . .	62·27 per cent.
Sulphur . . . . .	37·73 ...

The analysis was carefully made, and repeated for verification, so that, notwithstanding the specific gravity is so much lower than that assigned as characteristic of iron pyrites, there can be no doubt such is the constitution of this constituent of the ore in question.

The cobalt pyrites exhibits the usual characteristic reactions, generally subject to some modifications, which do not deserve notice, as I found them to be mostly owing to the high temperature at which my experiments were made: one however is rather remarkable, and not assignable to this cause, but probably to the particular natural constitution of the mineral, which, as I have found, in other cases modifies the behaviour of substances occasionally.

Ferrocyanide of potassium produces in acid solutions a

\* This description is the translation of a native one given to me with the minerals.

bluish-green precipitate, which completely dissolves up in forty-eight hours, provided the solution be not highly concentrated, to a brilliant emerald-green fluid, which is not affected by acids or by standing, but the colour of which changes to greenish yellow, without precipitation, by ammonia.

By very careful and repeated analysis, the reduction process having been adopted for the metal, I found the proportion of the constituents to be, taking the average,—

Cobalt . . . . .	64·64 per cent.
Sulphur . . . . .	35·36 ...

from which it is obvious the substance is a sub-sulphuret, that its constitution is  $\text{Co}_2\text{S}$ , a rather remarkable result, considering that the iron compound, doubtless of simultaneous formation, is different.

The cobalt pyrites has the specific gravity of 5·45. It is used by Indian jewellers for staining gold of a delicate rose-red colour; the *modus operandi* which they follow I have been unable to learn; it is a secret with them, which they are unwilling to disclose.

#### LVII. On the Structural Relations of Organized Beings.

By H. E. STRICKLAND, M.A., F.G.S.\*

I PROPOSE to make a few observations on the Relations which subsist between different organized beings in respect of the *similarities* of their physical structures. This limitation will exclude—first, the relations between individuals, such as that of parent to offspring, for in individuals of the same species the essential points of structure are not *similar*, but *identical*; and secondly, the relations between an organized being and the external circumstances of soil, climate, or food, to which it is adapted, in other words, between structure and function; for these adaptations of the one to the other, however interesting and admirable in themselves, are not relations of *similarity*.

On comparing together the innumerable species of organized beings, we find their structures to present every possible degree of variation, from an almost perfect identity to the utmost amount of difference which the mind can conceive any two organized bodies to possess. These agreements and differences are not however devoid of laws and principles; they admit of being classed under certain general heads, and we thus discover the traces of Divine workmanship not merely

\* Read before the Ashmolean Society of Oxford, March 10, 1845, and communicated by the Author.



in the structure of an individual organism, but in the mutual relations of those organisms, the due combinations of which constitute the Natural Systems of Botany and Zoology.

When the human mind first began to observe and to compare the structures of organic life, to generalize the points of agreement, and thus to lay the foundation of the Science of Natural History, no inherent principles of classification were even suspected to exist, characters were compared and generalized at random, and the arrangements which resulted were of the rudest and most unphilosophical kind. The most superficial and arbitrary characters were selected as the basis of classification, and no man was able to give a reason why one mode of arrangement should not be as correct and as true to Nature as another. Thus we find the older naturalists classing Lizards, Tortoises and Frogs with terrestrial Mammalia, under the name of "Four-footed Beasts," while Serpents were made into a distinct Class; and Whales, whose physiological organization is as highly developed as in any other Mammal, were dismissed among the cold-blooded Class of Fish, into which the humble Lobster and the Oyster entered from the other side to keep them company. By some authors we find the *Echinus* and the Hedge-hog approximated, because both are covered with spines; the Ammonite and the Rock-crystal were described in the same chapter "de lapidibus"; Shrew-mice and Spiders were classed together, because both were supposed to be venomous; Bats were referred to Birds, Corals to Plants, and so on.

In the course of the seventeenth century, the few who cultivated natural science began to be conscious that these crude arrangements were not satisfactory, or consistent with the realities of Nature; and in the works of Ray and of Lister, we perceive many instances of an instinctive preference for essential instead of arbitrary characters. But it was Linnæus who first pointed out in express terms the great principle of the *Subordination of Characters*. This principle teaches us to give to each point of structure its due weight, and to attach more value to those peculiarities whose immediate influence on the mysteries of Life often renders them the most difficult for our senses to appreciate, than to those external characters which, though most conspicuous to the eye, are but remotely connected with the real Essence of the creature. This principle has been further developed by later naturalists, especially by Cuvier, and accordingly we now find that in the modern systems of Zoology the *primary* divisions of the Animal Kingdom are based on characters derived chiefly from the *nervous* system, as being the most important feature in organization,

the *secondary* subdivisions are grounded on the *organs of respiration*, groups of a lower rank on the *digestive* system, and so on, the most superficial peculiarities, such as external form and colour, being reserved to characterize the ultimate groups of genera and species. These improved principles of classification are gradually bringing the systems of Zoology and Botany into a state of permanence, consistent with Nature, and satisfactory to that Truth-seeking Instinct which is inherent in the human mind.

A further advance of philosophical Classification has shown that the characters of organized beings require not only to be subordinated according to their importance, but subdivided according to their kinds. There are many instances of correspondence of structural characters in organic beings which can never by any process of subordination become elements in a natural classification, and it is important to distinguish those which *can* from those which *cannot* be so employed. Zoologists had long been aware that certain sets of characters produced an arbitrary or artificial method if employed for classification, while others seemed to lead to a natural system, but the question was involved in obscurity till the time of MacLeay, who was the first to give us clear definitions on the distinction between AFFINITY and ANALOGY. He applied his views indeed in support of a theory, the *Quinary System*, which few naturalists are now disposed to support, and with which we are not now concerned; but his elucidation of Affinities and Analogies is not the less valuable on that account. Although I am not disposed to take the same view of these principles as that of Mr. MacLeay, yet as the principles themselves are at the foundation of all sound classification, whether in Zoology or Botany, I may be allowed to make a few further remarks upon this subject.

It appears to me that the instances of resemblance or agreement of structure between any two species of organized beings should be reduced, not into *two*, but into *three* distinct classes, *Affinity*, *Analogy*, and a third, for which I propose to adopt the name of *Iconism*\*.

I. The highest class of these structural agreements is that of *Affinities*, which appear to be the direct result of those Laws of Organic Life which the Creator has enacted for his own guidance in the act of Creation. Affinity consists in an *essential* and physiological agreement in the corresponding parts of organic beings, resulting from a uniformity of plan

\* This term, suggested by the Rev. Dr. Ingram, President of Trinity College, appears preferable to *Mimesis*, which I had originally proposed to use.

which pervades the System of Nature\*. These essential agreements of parts consist rather in a similarity of organic composition and of relative situation, than of form. A microscopic examination of the primary tissue, or a chemical analysis of its substance, will often demonstrate the true affinities of a structure when its external form would only mislead us. And when we have proved an affinity to subsist between the structures of two organic beings, we then apply the term to the beings themselves, and say that an affinity subsists between them, greater or less, according to the number and importance of the organs in which such affinity is shown. Take for example the long, straight weapon of offence in the Narwhal, its general appearance is that of a *horn*, and such the vulgar accordingly call it; but if we examine its organization and its chemical composition, we find that both are utterly unlike those of real horns, but correspond to the structure of teeth. Further, if we examine the mode of its connexion with the skull, we find that it is inserted into a socket like other teeth, instead of being attached in the manner of horns, and we accordingly pronounce it to be not a horn but a tooth, developed for purposes of offence to an extraordinary extent. And having thus shown that the weapon of the Narwhal has no affinity to real horns, we no longer appeal to this structure in proof of any affinity between the Narwhal and the truly horned animals. Again, the Narwhal in its external form much resembles a Fish; but when we look to its nervous, circulatory, and reproductive organizations, which rank much higher in the scale of characters than external form, we find that it is no Fish, but a true Mammal, agreeing in every essential point with the warm-blooded quadrupeds of the land, to which its affinities are real and direct. Similar instances of the discordance between outward form and real affinity might be multiplied to a great extent; and it forms a constant employment for the scientific zoologist to distinguish real affinities from apparent ones, and thus to refer every organized being to its true position in the Natural System.

It will thus be seen that every instance of asserted affinity

\* We may suppose, for instance, that it was a law of organic creation that all Birds should have the anterior extremities modified into the form of wings; and in obedience with this law we find that there is no Bird which is absolutely without wings, though there are several kinds in which the wings are perfectly incapable of flight. Again, it is a law that Mammalia have neither more nor less than seven cervical vertebræ; and we find this law to hold good, without an exception, through the whole Class of Mammals, from the slender-necked Giraffe to the Whale, which can hardly be said to have any neck at all. The above, out of countless other examples, will show what is meant by *laws of organization*.

between two organic beings is merely a corollary deduced from an observed affinity between the corresponding organs in each; and though it is not usual to apply the term *affinity* to the similarities between parts, yet as the similarity between the wholes results from the similarities of their parts, the word *affinity* may be as correctly applied to the one as to the other. In works of comparative anatomy it is customary to speak of those members which are essentially equivalent in two organic beings as *analogous* organs, but we shall soon see that the word *analogy* has a very different sense; and as the relation between equivalent organs is one of real *affinity*, and forms the sole ground on which we assert the affinity of the whole beings, we may introduce the adjective *affine* or *homologous* in place of *analogous*, when referring to structures which essentially correspond in different organic beings.

When we say that Affinity consists in an essential agreement of structure resulting from a fixity of purpose in the mind of Creative Wisdom, it must not be supposed that all affinities are equally strong, direct, and palpable. Any agreement, however slight, or however concealed by more palpable differences, which forms part of the plan of organic existence, is a true affinity; and the principle of subordination of characters before referred to is merely the arranging of these affinities in the true order of their proximities. The proximity of affinities is in the inverse ratio of their essential importance, the most important agreements of characters being those which have the widest extent, and which therefore form affinities between the remotest points in the System of Organized Beings. We will illustrate this by an example showing the successive series of affinities which the same species bears to others, commencing with the most remote, and proceeding to the closest affinity which can subsist between two distinct species. We will take as an example the species Raven (*Corvus corax*).

A Raven has an Affinity to an	it is the same Affinity which exists between	and is derived from the Affinity between their respective	supplying the diagnostic characters of the
1. Oak-tree; 2. Locust; 3. Salmon; 4. Swan; 5. Humming Bird; 6. Sparrow; 7. Jay; 8. Magpie; 9. Carrion Crow;	all Animals and all Plants, Vertebrata and Insects, Birds and Fish, Insectores and Natatores, Conirostres and Tenuirostres, Corvidæ and Fringillidæ, Corvinæ and Garrulinæ, Corvus and Pica, one species of Corvus and another,	Organic Life, &c. nervous systems, &c. vertebral columns, &c. circulatory systems, &c. structure of feet, &c. conical beaks, &c. structure of nostrils, &c. short elevated beaks, &c. even tails, black plumage, &c.	Organic EMPIRE. Animal KINGDOM. PROVINCE, <i>Vertebrata</i> . CLASS, <i>Birds</i> . ORDER, <i>Insectores</i> . TRIBE, <i>Conirostres</i> . FAMILY, <i>Corvidæ</i> . SUBFAMILY, <i>Corvinæ</i> . GENUS, <i>Corvus</i> .

The affinities in this series are seen to accumulate successively as we proceed from the remotest organism to the approximate species. The Raven and Carrion Crow not only possess that superficial resemblance of form which constitutes their generic character, but they have in addition all the other points of affinity which extend from them to a greater or less distance into the realms of organic existence. Thus we find that

The Raven has		
organization	in common with all	Organized Beings.
a nervous system	...	Animals.
a vertebral column	...	Vertebrata.
a peculiar circulatory system	...	Birds.
perching feet	...	Insectores.
a conical beak	...	Conirostres.
the nostrils covered by feathers	...	Corvidæ.
ridge of the beak arched	...	Corvinæ.
an even tail	...	Corvus.
and		
a wholly black plumage	...	Carrion Crow.

It will be seen from the above example, that the whole process of classification consists in observing the affinities of structure in different beings, in estimating their importance, and in arranging them according to that estimate. It follows that a clear comprehension of *affinities*, as distinguished from the other kinds of resemblance, is essential to the objects of the scientific zoologist.

Although affinity consists in an essential and intimate agreement in the structure of certain organs, yet it by no means implies an identity of function in those organs. The modifications of external form are so various that we frequently find the same organ applied by different animals to purposes the most remote from its normal function; and on the other hand we see very different organs applied to discharge the same function. Thus, as a general proposition, it is certain that the proper function of wings is flying, of legs walking, of fins swimming; and yet we find examples where each of these organs is applied to any other function but its own, as in the case of the Bat, Seal, Ostrich, Penguin, Gurnard, and Flying Fish. Hence, although it is generally true that certain organs are destined to perform certain definite functions, yet the exceptions are so frequent as to make us attach a minor degree of importance to *function*, while we give the fullest weight to those essential properties which form the only test of real affinity.

II. We have next to consider that class of structural agreements known by the name of *Analogies*. These consist in a similarity of external form and of function connected with it, but without that agreement of essence which constitutes *Affinity*. These analogous agreements are equally the result of natural laws, but of laws of a different class from the former. Agreements of affinity are produced in conformity with the laws of the organic Creation, while analogies have a reference to the laws and properties of external and often inorganic matter. In obedience to these laws, it follows that whenever an instrument is required to produce a given effect upon external objects, or to resist their influences in a given manner, there is in general one method, and one only, of effecting the object in the best and most effectual way. Accordingly, whatever be the organ or instrument employed, that organ must have a certain and definite mechanical structure bestowed upon it to obtain the desired end. As a general rule, the same end is attained in different organic beings by means of the same set of organs; but when those organs are required for any other purpose, or are so modified as to be unfit for that special end, then some other set of organs are endowed with the requisite external structure and are called upon to act as substitutes for the legitimate instruments. Examples of this adaptation of organs to purposes remote from their normal destination are numerous and well-known; and I cannot do better than refer to the late Mr. John Duncan's work on the *Analogies of Organized Beings*, where there are numerous examples of such analogies arranged in a tabular and highly perspicuous form. We need only take the Elephant as an instance. We may suppose that this animal required horns for the purpose of defence, but it belongs to an order, the Pachydermata, in which horns are uniformly absent, and the laws of Affinity forbade their introduction. To supply this defect, the incisor teeth are removed from their usual duties of mastication, and are so developed as to assume the form and discharge the function of horns. Further, the great size and weight of these lengthened tusks required a great strength and shortness of neck, and the animal was consequently unable to reach the ground with his mouth. A hand was therefore required to convey the food to the mouth, but the vast weight of the animal required a massive structure in the feet, which forbade them to be adapted to the purpose of hands. To supply this want then the *nose* is lengthened out, furnished with muscles, divided at the end into a finger and thumb, and in this proboscis behold a hand! almost equal in delicacy of manipulation to the hand of Man. And thus we see the Ele-

phant, endowed in one respect with an analogy to the Ox, and in another respect to Man, yet having no immediate affinity with either.

As then Analogy consists in an agreement of function, and only of form so far as it tends to discharge that function, it follows that real and genuine *Analogies* may take place between the works of Nature and the works of Man, while no such relation of *Affinity* can possibly exist. When, for instance, the inventive powers of Man are called upon to imitate any of the operations of Nature, the external matter to be acted on being in both cases the same, a similar arrangement of form is adopted by both. If the problem be to make a floating body adapted for rapid motion through water, Man either by practical experiment or mathematical calculation produces the form of a boat, and thus unconsciously imitates the structure of the Whale and Seal among Mammals, the Penguin among Birds, the Ichthyosaurus and Turtle among Reptiles, the Fish among Vertebrata, the Dytiscus among Coleoptera, the Notonecta among Hemiptera, Sepia among Mollusca, Physalia among Acalephæ, &c. &c. Nor is the analogy between a ship and a Fish confined to the external form only; the keel of the one represents the spine of the other, the "ribs" of both agree in name as in nature, the rudder coincides with the tail, the oars with the fins, the masts with the spinous processes, the running rigging with the tendons, the seamen with the muscles, the *look-out man* on the fore-castle with the eye, and the captain in the cabin with the mental faculties in the Fishes' brain. Again, what can be more striking than the analogy between a locomotive steam-engine and a living Animal? We see in both an analogous respiratory and digestive system, the same necessity for food and drink and oxygen to sustain that internal combustion which is the source of the vital action, the same obedience of the organs of motion to the impulse of the governing mind, and the same wear and tear of the system, terminating in old age and sudden or gradual death. Yet in all these cases there is no set purpose on the part of Man to imitate the works of Nature, he merely applies the faculties which God has given him to elicit the properties which the same God has given to matter; and by this process alone he often arrives at the same or similar results to those at which Creative Wisdom had arrived before him. It appears to me therefore, that relations of Analogy, that is to say, agreements in structure in consequence solely of an agreement in the function to be performed, may be as truly and as correctly asserted to exist between ar-

tificial and natural productions, as between one object of the latter class and another. It is clear from this how much lower *Analogies* ought to stand in our estimation than *Affinities*. The latter form an essential part of that magnificent plan of Creation, which notwithstanding the amount of attention which Man has given to it, is of so transcendental a nature, that it may almost be said to be yet "to us invisible or dimly seen." Analogies, on the contrary, appear not to form any element whatever in the great System of Nature, but are merely examples of the recurrence of certain mechanical forms whenever the production of a certain mechanical action called for them; and so far from their being at or beyond the verge of human comprehension, we have seen that Man enjoys the high privilege of copying by these Analogies, at a humble distance, the far transcendent works of his Maker.

It would be an improvement in the language of Comparative Anatomy, if the term *analogous organs* were limited to the sense above defined. The serrations in the beak of a duck, for instance, are *analogous* in form and in function to *teeth*, but in their essential nature they are only a corneous modification of the *lips*. Most anatomists, however, would habitually say that the beak of a bird is *analogous* to the lips of a Mammal, though it must be evident how much more precise their language would become if they spoke of this *essential* relation as an *affinity*, and applied the word *analogous* to *formal* or *functional* relations only. A similar inaccuracy is committed by geologists in speaking of the *recent analogue* of a fossil species, meaning thereby that living species which has the nearest affinity to the extinct one. It would be more correct if they would term it the *recent affine*, or the *recent homologue*.

III. There is yet a third species of relation of structural similarity between organized beings which has usually been confounded with Analogy, but which appears to me to be distinct from it in kind, as well as far inferior to it in importance,—I refer to those cases where a resemblance in form or configuration exists, but without any perceptible identity either of essence or of function. Such, for example, are the resemblances between the flower of the *Bee Orchis* and a *Bee*, between the shell of *Murex haustellum* and a *Woodcock's head*, between a *Fungia* and a *Fungus*, *Ovulum* and an *egg*, *Haliotis* and an *ear*, &c. To this class also belong the numerous instances of *similarity of colour* between Birds whose affinities are remote, such as the resemblance of *Oriolus* to *Xanthornus*, of *Dicrurus* to *Corvus*, of *Cissopis* to *Pica*, of *Agelaius phœni-*



*ceus* to *Campephaga phœnicea*. Many errors of classification have been caused by mistaking these similarities for true affinities.

Not only are such cases of external resemblance unconnected with any agreement in the essential structures of the bodies compared, but there is no conceivable similarity in the functions which they are created to discharge. I think therefore that it is not going too far, nor departing from that veneration which the true naturalist will always feel for Nature's God, to call such superficial coincidences of form *accidental*. They seem to arise from the exuberant variety of the works of Nature which causes an occasional recurrence of similar forms, without any express design for such coincidences. Nothing can be inferred from such resemblances, either as to essential affinity or functional design; and they would almost have been beneath our notice, were it not that some authors have regarded them as examples of real analogies. The advocates of the Quinary theory of classification, who regard *Analogies* to be as important an element in the Natural System as *Affinities*, often speak of these mere resemblances in the light of true Analogies, and appeal to them in confirmation of their views. Regarding however, as I do, those views to be erroneous, I think it important that the distinction between *functional Analogy* and *mere resemblance* should be clearly pointed out; and to render the distinction more marked, I would distinguish the latter by the new term *Iconism*.

We must beware indeed of too hastily pronouncing an instance of resemblance to be an *Iconism*, merely because we cannot immediately detect any functional analogy. There may be real reasons for these resemblances, real agreements in the functions to be discharged, which we have not yet detected, and perhaps may never discover. A person might say, for instance, that the species of Mantis called the "walking leaf" presents a mere *Iconism* or accidental resemblance to true leaves; whereas it is highly probable that this very resemblance is given to the animal to enable it to remain concealed from its foes amid the verdant foliage. Such at least is undoubtedly the intention of numerous instances in which animals present an analogous colour to the surrounding surface, and even undergo corresponding changes with it, such as that of the Ptarmigan, which during summer is of a speckled gray plumage, like the lichen-covered rocks which it frequents, while in winter it becomes a pure white when those rocks are covered with snow.

I have now endeavoured to show that the relations of resemblance in organized beings are of three kinds, diminishing

successively in importance; that *Affinities* are expressions of the real and elementary and esoteric Plan of Creation which the Author of Nature has been pleased to follow; that *Analogies* are coincidences of structure consequent solely upon an identity of external physical conditions; and that *Iconisms* are merely accidental recurrences of similar forms resulting from the exuberance of Nature's riches. It is evident that these distinctions must be clearly understood before we can make any progress in Natural History as a Science, and the remarks above offered may perhaps aid in drawing attention to the subject or removing the difficulties which surround it.

LVIII. *Abstract of Meteorological Observations made during the year 1845 at Gongo Soco, in the interior of Brazil. By WILLIAM JORY HENWOOD, F.R.S., F.G.S., Member of the Geological Society of France, Chief Commissioner of the Gold Mines of Gongo Soco and Catta Preta, &c. &c.\**

**T**HE rich gold mines of Gongo Soco are situated in the province of Minas Geraës, about forty-eight miles north-west of the city of Ouro Preto† (Villa Rica), in long.  $43^{\circ} 30'$  west and lat.  $19^{\circ} 58' 30''$  south, in a vale bounded on the north by the wooded mountain-range of Tejuco, and on the south by undulating grassy lowlands, which at the distance of about eight miles are terminated by the mountain-chain of the Caraças, which rises from 4000 to 5000 feet above the plain.

Barometrical measurements‡ give Gongo Soco an elevation of about 3360 feet above the sea at Rio de Janeiro.

The thermometrical observations were made at such times as my occupations permitted, but the hours are probably not the best possible§. The midnight observations were made by Captains Blamey, Luke, and Guy, and the thermometer they used needed a constant correction of  $2^{\circ} \cdot 8 +$ ; all the others are my own, and the thermometer I employed was a standard one (No. 89) of the British Association. The thermometer is suspended in a wooden box pierced with numerous holes, and hangs at about six feet above the ground, in a shed open at all sides, and is well protected,—as well from reflected heat as from the direct rays of the sun.

\* Communicated by the Author.

† Mr. Caldcleugh estimates the elevation of Ouro Preto at 3969 feet above the sea.—*Daniell's Meteorological Essays*, p. 345.

‡ Made by the Austrian Mining Engineer, M. Virgil von Helmreichen.

§ The observations at 4 and 8 P.M. give higher results than would have been afforded at 3 and 9 respectively.

Table I.  
Hourly mean and extreme temperatures for every month.

	6 A.M.			9 A.M.			Noon.			4 P.M.			6 P.M.			8 P.M.			9 P.M.			Midnight.		
	Max.	Min.	Mean.	Max.	Min.	Mean.	Max.	Min.	Mean.	Max.	Min.	Mean.	Max.	Min.	Mean.	Max.	Min.	Mean.	Max.	Min.	Mean.	Max.	Min.	Mean.
January.	69.8	64.	66.7	76.2	65.5	70.	83.	65.	75.5	84.5	69.	75.9	80.5	68.8	73.2	76.8	67.8	71.	76.8	68.5	71.8	64.8	67.8	
February	71.	60.8	67.2	74.6	63.	70.	81.	66.	75.4	84.	64.6	76.6	77.5	63.5	73.4	75.8	58.	71.	75.8	68.3	72.8	61.8	68.1	
March...	71.	62.	66.3	73.2	64.	68.6	78.5	67.	73.3	81.2	69.5	74.8	78.	67.	73.	74.2	65.	70.1	73.2	64.8	69.	70.8	60.8	66.5
April ...	68.	61.	65.3	72.5	63.	67.7	75.	65.	71.6	77.	64.5	72.5	75.	63.	70.	71.5	62.8	68.1	70.2	63.2	67.1	69.8	59.8	66.2
May.....	68.	54.2	61.3	70.	61.8	65.4	75.	65.5	70.8	76.5	66.	71.5	73.	62.	67.5	69.	60.	64.3	66.	59.8	62.9	67.8	57.8	62.3
June ...	64.	47.5	52.9	65.	55.	60.	72.	60.	67.7	72.	66.5	68.5	66.5	58.	62.4	64.	54.	58.4	61.	54.5	56.3	62.8	50.8	55.4
July.....	60.5	42.	54.6	67.	52.8	58.9	68.	60.	65.2	71.	62.	66.2	66.5	57.5	61.8	64.	52.2	58.1	60.2	51.5	56.2	65.8	48.8	55.7
August..	64.2	49.	58.	67.	57.	61.9	73.	61.	69.1	75.5	61.	70.7	71.6	60.	66.8	69.	58.5	62.9	65.8	57.	62.4	63.8	52.8	58.7
Sept. ...	65.2	49.9	59.4	70.	55.	63.	79.3	58.	69.9	81.	58.2	70.8	77.	57.8	67.8	71.	56.9	63.5	70.3	58.	62.8	68.8	53.8	60.2
October	70.2	55.5	64.7	75.5	59.6	68.8	84.	64.	75.7	85.2	66.	76.3	80.	65.	73.2	74.4	62.	69.5	73.	61.8	67.5	70.8	56.8	65.8
Nov. ...	70.	61.	66.8	77.	62.6	71.1	85.	65.	77.3	87.	67.	77.8	84.	65.	74.9	76.5	64.5	71.7	74.	64.	70.4	72.8	58.8	67.5
Dec. ...	70.8	63.	66.2	77.5	67.	71.3	84.	67.	77.8	85.4	68.5	77.5	82.	67.8	76.	77.8	66.	71.4	74.5	64.4	70.	72.8	62.8	67.4

Table II.

Mean temperature of each month.

January .....	71.07	September ...	64.67
February .....	71.25	October .....	70.2
March.....	70.2	November ...	72.19
April .....	68.65	December.....	72.2
May .....	65.75		
June .....	60.2		
July .....	59.52		
August .....	63.81		

Table III.—Mean temperature of each of eight hours.

6 A.M. . . .	62°45	6 P.M. . . .	70°
9 ... . . .	66°39	8 ... . . .	66°66
Noon . . . .	72°44	9 ... . . .	65°11
4 P.M. . . .	73°26	Midnight . .	63°42

The foregoing results give a mean temperature of 67°46: as however the observations at 4 and 8 P.M. give higher results than would have been afforded if the observations had been made at 3 and 9 P.M. instead, and as we are without observations at 3 A.M., I consider the foregoing temperature about 2°3 above the true mean, which in this case will be about 65°14.

Circumstances have prevented my making many comparisons between the indications of the thermometer in the shade and when exposed to the direct action of the sun's rays. In the following instances, however, the instrument was suspended at about three feet above a surface of newly-turned garden-mould, of a deep red colour.

Table IV.

Date.	Therm. shade.	Therm. sunshine.	Remarks.
June 28, 3 P.M.	67°5	87°	Calm.
... 29, Noon	64°5	79°8	Calm.
Nov. 2, 2 P.M.	83°5	99°3	Light breeze E.
Dec. 29, 4 ...	82°2	97°2	Brisk breeze W.

My garden is a level spot of about one-third of an acre in area, and contains several orange and coffee trees, besides other shrubs; none of them, however, are very large; and the rain-gauge is placed on the ground at a distance from them all, in the centre of the garden.

Table V.—Quantity of rain.

	inches.
January . . . . .	23°32
February . . . . .	23°03
March . . . . .	12°84
April . . . . .	8°06
May . . . . .	1°60
June . . . . .	1°09
July . . . . .	1°28
August . . . . .	1°05
September . . . . .	3°88
October . . . . .	9°14
November . . . . .	27°
December . . . . .	12°46

Total in 1845. 124°75

Although the rain is occasionally very heavy, I have seen none to compare with the results recorded by Prof. Forbes\*. The heaviest showers I have seen, were

January 13,	6	P.M.,	when	1.12	inch	of	rain	fell	in	1	hour.
...	17,	2	...	...	0.72	...	...	20	minutes.		
November 13,	4	...	...	1.04	...	...	17	...			
...	24,	5	...	...	1.2	...	...	25	...		
...	25,	2	...	...	2.24	...	...	1½	hour.		

The heaviest falls of rain during twenty-four hours were, February 22, when 3.92 inches were collected; November 26, when 3.76 inches were collected.

At the commencement of the wet season, heavy thunderstorms precede the rains for several days; they usually begin early in the afternoon, but generally pass off as evening approaches. As the season advances, they become daily later; and towards its conclusion, the time of their appearance is very irregular.

From April to August there is usually but little rain, and the continued drought enables the farmer to burn the dry stubble of his maize and beans, and to clear his grounds for tillage; for several weeks during August and September the atmosphere is filled with the smoke from these burnings; and at this time violent thunderstorms, with heavy showers, are frequent†.

For two or three weeks, about the end of January and the beginning of February, there is usually a cessation of rain and a continuance of unclouded sunshine (the *veronica*); but no such interval occurred in 1845‡.

I am unprovided with a barometer, hygrometer, and many other instruments necessary for a regular course of meteorological observation, as on leaving England my stay in Brazil was not expected to have exceeded a few months; and I have not obtained them since, as the requisite attention to them would interfere with the indispensable duties of my office. I have, however, a considerable series of observations on the

\* Reports of the British Association (1832), p. 252, and (1840) p. 113-116.

† I believe it has been long known that thunderstorms and rains follow the fires on the great prairies of North America, but I am unable to refer to my authority for the remark.

‡ A season seldom passes without heavy hail-storms in this province: during the early part of the wet season of 1844, I saw two such here; but during the present year we have had none, although there have been some of great severity in the neighbourhood.

intensity of terrestrial magnetism, but their reduction must await more leisure.

I am aware of the poverty of my remarks, and nothing but the scantiness of recorded observations in the interior of Brazil would have emboldened me to submit them to your readers.

Gongo Soco Gold Mines,  
January 16, 1846.

W. J. HENWOOD.

LIX. On *Pegmine and Pyropine*, animal substances allied to *Albumen*. By ROBERT D. THOMSON, M.D., Lecturer on *Practical Chemistry in the University of Glasgow*.\*

**T**HIS paper was written for a government report, detailing the results of an extensive series of experiments made on the influence of different kinds of food in feeding cattle during the course of 1845.

The report was drawn up last year, but has not yet been published. In reference to the reducing powers of the animal system, it is remarked that "there is only one instance with which physiologists are at present acquainted that could be adduced as evidence in favour of any substance being rendered more complex in the animal system, viz. the production of fibrine or flesh from curd or caseine. So far as chemical experiments carry us, we are not in a condition to affirm that no fibrine exists in milk; but it must be admitted that none has as yet been detected. If these be correct, then it would appear to follow that the infant fed on milk must derive its flesh from the curd of that fluid; and that as curd contains no phosphorus, while fibrine does, the curd of the milk in order to form muscular fibrine is united to phosphorus in the animal system, and is thus built up instead of being, as is the rule with other substances, reduced to a smaller number of elements. The objection to this view of the subject is, that the experiments which have been made on fibrine do not prove that it contains phosphorus. They only show that phosphoric acid can be detected in it even when it is purified in the most careful manner suggested by chemical knowledge, and it would therefore be somewhat premature to adopt any such analogy as that which we have been considering."

In support of the view first suggested by Beccaria and advocated in recent times by Prout, that the animal system merely modifies the substances which it employs as food, and does not produce them from its elements, a series of experiments made by the writer four years ago, may be quoted, hitherto unpublished, which demonstrate, that in the ocean, as

\* Communicated by the Author.

on land, the higher subsist on the lower animals, *because* the latter consist of the same materials of which the higher systems are composed. Without the lower animals, therefore, it is obvious the larger could not exist, and hence we may infer that the inferior organizations first peopled the earth, an argument opposed to the idea of some geologists, that animals have not been developed in succession. As it is well known that oysters serve as food for larger fishes, and these again for more powerful species, experiments were made to determine the composition of oysters, herrings, and haddocks, since it is highly probable that these prey on each other. Portions of these fishes were well-washed in water, to remove the oil and soluble matters; the white residue was then treated with alcohol and repeated digestions in æther. The resulting matter, which was considered to be pure fibrine, was found to have the following composition in the three species of fish when burned with chromate of potash:—

	Oyster.	Herring.	Haddock.
Carbon . . .	53·98	53·77	53·67
Hydrogen . . .	} 46·02	7·44	7·00
Nitrogen . . .		16·23	16·89
Oxygen . . .			
Sulphur . . .		22·56	22·44
	100·00	100·00	100·00

It is obvious, therefore, that fibrine can be obtained with the greatest facility and of the purest form from fish.

*Can a substance be obtained from Albumen, &c. free from Sulphur?*

In all of these kinds of fibrine, sulphur could be readily detected; nor was it found possible by any of the methods which have been hitherto described, to obtain either from fibrinous matter or from albuminous substances, a simpler body destitute of sulphur. The analysis of the milk detailed in a previous part of the report, afforded excellent opportunities of testing the accuracy of the idea supported by some of the continental chemists, that a substance can be obtained by the action of potash upon albuminous substances which contains no sulphur. On repeating the experiments that have been detailed in books upon a considerable scale with caseine or curd of milk, which were carefully conducted by William Parry, Esq., late of H.M. 4th Regiment, it was uniformly found that the resulting product contained sulphur. By this statement, certainly it is not meant to infer that such a substance may not exist, but only that the writer has not been

able to procure such a substance as *proteine* by following most scrupulously the directions supplied by its original describer, and others who have copied his descriptions. His scepticism on this subject originated some years ago when engaged in researches on the brain, an abstract only of which has been published in Liebig's edition of Geiger's *Pharmacie*. The process of analysis for this intricate combination consisted in dissolving the albuminous part of the nervous system in dilute caustic potash; a reagent which produces no soluble power on the peculiar matter of the brain, but combines with it, forming an insoluble salt. The potash solution, on being withdrawn from the insoluble matters, yielded by neutralization with acetic acid, a substance which ought to have been *proteine*, because it was obtained by precisely the same process as that which has been described as the best for procuring that substance. But on dissolving after washing in potash, adding acetate of lead and boiling, it gave an abundant black precipitate, indicating the presence of sulphur. This experiment was shown to Prof. Liebig by the writer at the time (1842), and it is believed that that distinguished chemist considers the existence of *proteine* problematical.

#### PEGMINE.

About the same period (four years ago), the writer examined a product of the disease usually known under the name of the buffy coat of the blood, a coating of a buff colour, which usually exhibits itself on the surface of inflamed blood, and which has attracted much of the attention of writers upon pathological subjects. He found it to be a distinct body, and he has been in the habit of describing it in his lectures under the name of *pegmine* (from *πήγμα*, *coagulum*). It partially dissolves by long-continued boiling in water, but may be washed in cold water, like fibrine, without undergoing any decomposition. It therefore possesses an equal right with fibrine to the character of a body *sui generis*. When dissolved in potash and precipitated with acetate of lead, and the liquid is boiled, a black precipitate of the sulphuret of lead falls. The following are the results of the analysis of this substance made in 1842, and which the writer has been in the habit of quoting in his lectures.

##### I. *Pegmine* containing *Fat*.

The first specimen was prepared by simply washing the buffy coagulum with repeated additions of cold water. It was taken from a patient affected with a violent attack of



pleuritis. It is obvious from the analysis that it contained a considerable amount of fatty matter:

Carbon . . .	58·80	
Hydrogen . .	8·44	
Nitrogen . .	} 32·76	
Oxygen . . .		
Sulphur . . .		
	100·00	

### II. *Pure Pegmine.*

Another specimen procured from a different patient, also affected with an attack of inflammation of the membrane of the lungs, was treated with cold water, alcohol, and æther to remove all the fatty and oily matters mixed with it; when burned with chromate of lead, the following result was obtained:—

	I.	II.	Approximate true composition.
Carbon . . .	52·07		52·07
Hydrogen . .	7·80	7·14	7·14
Nitrogen . .	} 14·00	14·40	14·20
Oxygen . . .			
Sulphur . . .	}		
	100·00		100·00

It is possible that the nitrogen is somewhat undervalued.

In the first analysis, the pegmine was dried at 212°, in the second at 300°. The same substance is met with in the inferior animals, especially in the horse, although not, it is believed, in the healthy state of that animal, as has been asserted, but in a similar condition of the animal to that in which it appears in the human subject—*inflammation.*

### PYROPINE.




The only body which bears any resemblance in composition to the so-called proteine, is a beautiful substance which is found occasionally in the tusk of the elephant, occupying the hollow portion of the interior of that part of the animal. It possesses a fine ruby tint, and is sometimes tough, but when of the finest colour is brittle. Sections of it exhibit occasional traces of the remains of organization. It is insoluble in water, and thus differs from glue or gelatine, to which it has some affinities in its physical aspect. The writer has not been able to satisfy himself that it contains no sulphur, in consequence of its difficult solubility in caustic potash. The composition of pyropine by two analyses is as follows:—

	I.	II.
Carbon . . .	53·33	53·50
Hydrogen . . .	7·52	7·66
Nitrogen . . .	14·50	
Oxygen . . .	24·65	38·84
Sulphur . . .	24·65	
	100·00	100·00

These analyses were communicated to Prof. Liebig some years ago, and published by him in his edition of Geiger's *Pharmacie*, with the omission only of the nitrogen, which had not then been determined.

Liebig has suggested, with great plausibility to the writer, that this beautiful substance may be an altered form of blood, an idea which receives some support from the fact, that when pyropine is incinerated, it leaves 0·52 per cent. of a reddish ash,—a fact not sufficiently, perhaps, conclusive.

When pyropine is boiled in water, the liquid is not precipitated by infusion of nut-galls, a proof that it contains no gelatine or glue. Neither is it precipitated by acetate of lead. The colour of pyropine is not altered by this treatment, with the exception that a few scanty flocks of a membranous-looking matter floated about. When broken into coarse powder, it has a rich ruby colour; and hence its name (*pyrōpus*, a ruby). In fine powder it is brown; a minute portion of it dissolved in hot alcohol, and was deposited on cooling in the form of ferruginous flocks. The following formulæ would probably represent the relations of the preceding bodies to each other. They must, however, be considered as mere possible representations of their composition, calculated to exhibit the difference in reference to their increasing amount of oxygen.



In these formulæ, pyropine is represented as differing from fibrine in containing no sulphur, and pegmine from the preceding bodies by the presence of an additional quantity of oxygen.

The increased amount of oxygen in pegmine may be explained by the circumstance, that in inflammatory action respiration is more rapidly carried on, and in consequence a greater quantity of oxygen is introduced into the system than in the healthy condition of the body. In all cases of coagulation of blood in contact with oxygen, there is observable a light coloured portion situated on the surface of the coagulum, affording a proximate illustration of the production of the buffy coat.

LX. *On certain Definite Multiple Integrals.*

By the Rev. BRICE BRONWIN\*.

IN the integrals treated of in this paper, let the limits of integration be given by the equation

$$\frac{x^2}{\alpha^2} + \frac{y^2}{\beta^2} + \dots + \frac{z^2}{\lambda^2} = 1, \quad \dots \dots \dots (a.)$$

including both negative and positive values of the ( $n$ ) variables. Let  $P(r) = 1 \cdot 2 \cdot 3 \dots r$ ; and for convenience let  $D, D_1, \&c.$  stand for  $\frac{d}{dx}, \frac{d}{dy}, \&c.$  respectively. The general term of the series expressing the value of

$$\psi = \iint \dots \phi(g - x, h - y, \dots) dx dy \dots$$

is  $\frac{D^{2p} D_1^{2q} \dots}{P(2p) P(2q) \dots} \phi(g, h, \dots) \iint \dots x^{2p} y^{2q} \dots dx dy \dots,$

the odd powers of  $x, y, \&c.$  obviously vanishing. But by a well-known theorem, integrating for positive values of  $x, y, \&c.,$  and doubling the result; making  $s = p + q + \dots,$  this term becomes

$$\frac{P\left(p - \frac{1}{2}\right) P\left(q - \frac{1}{2}\right) \dots}{P(2p) P(2q) \dots} \frac{\alpha \beta \dots}{P\left(s + \frac{n}{2}\right)} (\alpha D)^{2p} (\beta D_1)^{2q} \dots \phi(g, h, \dots).$$

By another known theorem

$$P\left(p - \frac{1}{2}\right) = \frac{\pi^{\frac{1}{2}}}{2^{2p}} \frac{P(2p)}{P(p)}, \quad \&c.;$$

and the above term is changed into

$$2^n \pi^{\frac{n-1}{2}} \alpha \beta \dots \frac{P\left(s + \frac{n-1}{2}\right)}{P(2s+n)} \frac{(\alpha D)^{2p} (\beta D_1)^{2q} \dots}{P(p) P(q) \dots} \phi(g, h, \dots).$$

And the sum of all the terms of the order  $s$  is

$$2^n \pi^{\frac{n-1}{2}} \alpha \beta \dots \frac{P\left(s + \frac{n-1}{2}\right)}{P(2s+n)} \{(\alpha D)^2 + (\beta D_1)^2 + \dots\}^s \phi(g, h \dots) \\ = \alpha \beta \dots f(s) \text{ to abridge.}$$

Hence  $\psi = \alpha \beta \dots \sum f(s), \quad \dots \dots \dots (b.)$

$s$  having all integer values from zero to infinity.

Suppose  $\phi(g, h \dots)$  such that

$$(D^2 + D_1^2 + \dots) \phi(g, h \dots) = 0.$$

\* Communicated by the Author.

Then, separating the symbols of operation from those of quantity,

$$D^2 + D_1^2 + D_2^2 + \dots = 0 \quad D_n^2 = -(D^2 + D_1^2 \dots + D_{n-1}^2),$$

$$D_n^4 = -D_n^2(D^2 + D_1^2 + \dots) = -(D^2 + D_1^2 + \dots) D_n^2$$

$$= (D^2 + D_1^2 + \dots)^2, \quad D_n^6 = -(D^2 + D_1^2 + \dots)^3, \text{ \&c.}$$

Therefore

$$\{\alpha^2 D^2 + \epsilon^2 D_1^2 \dots + \lambda^2 D_n^2\}^s = (\alpha^2 D^2 + \epsilon^2 D_1^2 \dots + \theta^2 D_{n-1}^2)^s$$

$$+ s(\alpha^2 D^2 + \epsilon^2 D_1^2 \dots + \theta^2 D_{n-1}^2)^{s-1} \lambda^2 D_n^2$$

$$+ \frac{s(s-1)}{2} (\alpha^2 D^2 + \epsilon^2 D_1^2 \dots + \theta^2 D_{n-1}^2)^{s-2} \lambda^4 D_n^4 + \text{\&c.}$$

$$= (\alpha^2 D^2 + \epsilon^2 D_1^2 \dots + \theta^2 D_{n-1}^2)^s$$

$$- s(\alpha^2 D^2 \dots + \theta^2 D_{n-1}^2)^{s-1} (\lambda^2 D^2 + \lambda^2 D_1^2 \dots + \lambda^2 D_{n-1}^2) + \text{\&c.}$$

$$= \{\alpha^2 D^2 + \epsilon^2 D_1^2 \dots - \lambda^2 D^2 - \lambda^2 D_1^2 \dots\}^s$$

$$= \{(\alpha^2 - \lambda^2) D^2 + (\epsilon^2 - \lambda^2) D_1^2 \dots + (\theta^2 - \lambda^2) D_{n-1}^2\}^s.$$

In this case therefore

$$\psi = 2^n \pi^{\frac{n-1}{2}} \alpha \epsilon \dots \lambda \sum \frac{P\left(s + \frac{n-1}{2}\right)}{P(2s+n)} \left\{ (\alpha^2 - \lambda^2) D^2 + (\epsilon^2 - \lambda^2) D_1^2 \dots \right\}^s \phi(g, h \dots) \dots \dots (c)$$

Change in this last  $\alpha, \epsilon \dots \lambda$  into  $a, b \dots l$ , but so that

$$\alpha^2 - \lambda^2 = a^2 - l^2, \quad \epsilon^2 - \lambda^2 = b^2 - l^2, \text{ \&c.};$$

and let this change  $\psi$  into  $\phi$ , we have obviously

$$\psi = \frac{\alpha \epsilon \dots}{a b \dots} \phi. \dots \dots (d)$$

This is an extension of Laplace's theorem relative to the attraction of ellipsoids on a point exterior. And if  $\alpha, \epsilon, \text{ \&c.}$ ,  $a, b, \text{ \&c.}$  be independent of  $g, h, \text{ \&c.}$ , we have also

$$\left(\frac{d}{dg}\right)^m \left(\frac{d}{dh}\right)^n \dots (\psi) = \frac{\alpha \epsilon \dots}{a b \dots} \left(\frac{d}{dg}\right)^m \left(\frac{d}{dh}\right)^n \dots (\phi). \dots (e)$$

To give an example or two, let

$$R = \{(g-x)^2 + (h-y)^2 \dots\}^{\frac{1}{2}},$$

and let  $R_1$  stand for the same quantity when  $\alpha, \epsilon, \text{ \&c.}$  are changed into  $a, b, \text{ \&c.}$ ; then, since

$$\left(\frac{d^2}{dg^2} + \frac{d^2}{dh^2} \dots\right) \frac{1}{R^{n-2}} = 0,$$

$n$  being the number of variables, we have

$$\iint \dots \frac{dx dy \dots}{R^{n-2}} = \frac{\alpha \beta \dots}{a b \dots} \iint \dots \frac{dx dy \dots}{R_1^{n-2}} \dots \dots (1.)$$

In the next two examples, by way of distinction, let  $x, y, \&c.$  be changed into  $\alpha x, a x, \beta y, b y, \&c.$ ; and let there be three variables,

$$\left. \begin{aligned} & \iiint \frac{\alpha \beta \gamma dx dy dz}{\{(g-\alpha x)^2 + (h-\beta y)^2 + (k-\gamma z)^2\}^{\frac{3}{2}}} \\ &= \frac{\alpha \beta \gamma}{a b c} \iiint \frac{a b c dx dy dz}{\{(g-ax)^2 + (h-by)^2 + (k-cz)^2\}^{\frac{3}{2}}} \\ &= \iiint \frac{\alpha \beta \gamma dx dy dz}{\{(g-ax)^2 + (h-by)^2 + k^2\}^{\frac{3}{2}}} + Ac + Bc^2 + \&c. \\ &= \iint \frac{\alpha \beta \gamma dx dy \sqrt{1-x^2-y^2}}{\{(g-ax)^2 + (h-by)^2 + k^2\}^{\frac{3}{2}}} + Ac + Bc^2 + \&c. \\ &= \iint \frac{\alpha \beta \gamma dx dy \sqrt{1-x^2-y^2}}{\{(g-ax)^2 + (h-by)^2 + k^2\}^{\frac{3}{2}}} \end{aligned} \right\} (2.)$$

the equation of limits for the first member being  $x^2 + y^2 + z^2 = 1$ , that for the second  $x^2 + y^2 = 1$ . This result is obtained by developing into series relative to  $z$ , then integrating for this quantity, and lastly diminishing  $c$  without limit, the quantities  $A, B, \&c.$  being finite. In the second member it must be observed, that since  $c=0, a = \sqrt{\alpha^2 - \gamma^2}, b = \sqrt{\beta^2 - \gamma^2}$ .

If we differentiate (2.) for  $(k)$ , we have

$$\left. \begin{aligned} & \iiint \frac{(k-\gamma z) dx dy dz}{\{(g-\alpha x)^2 + (h-\beta y)^2 + (k-\gamma z)^2\}^{\frac{3}{2}}} \\ &= \iint \frac{k dx dy \sqrt{1-x^2-y^2}}{\{(g-ax)^2 + (h-by)^2 + k^2\}^{\frac{3}{2}}} \end{aligned} \right\} \dots \dots (3.)$$

By integrating the first members of (2.) and (3.) relative to  $z$ , we should obtain very singular results.

Let

$$R = \{(g-x)^2 + (h-y)^2 + (k-z)^2\}^{\frac{1}{2}}, \quad V = \iiint \frac{dx dy dz}{R},$$

U the same integral when  $\alpha, \beta, \gamma$  are changed into  $a, b, c$ .  
Make

$$\frac{\sqrt{(g-x)^2 + (h-y)^2}}{R} = \sin u, \quad \frac{g-x}{\sqrt{(g-x)^2 + (h-y)^2}} = \sin v.$$

Then

$$x = g - R \sin u \sin v, \quad y = h - R \sin u \cos v, \quad z = k - R \cos u, \\ dx dy dz = -d R \sin u du dv;$$

or rather

$$dx dy dz = d R \sin u du dv,$$

because (R) decreases while (x) increases. Therefore

$$U = \iiint R d R \sin u du dv = \frac{1}{2} \iint R^2 \sin u du dv.$$

Putting the values of x, y, and z in

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1,$$

and making

$$A = \frac{1}{a^2} \sin^2 u \sin^2 v + \frac{1}{b^2} \sin^2 u \cos^2 v + \frac{1}{c^2} \cos^2 u,$$

$$B = \frac{g}{a^2} \sin u \sin v + \frac{h}{b^2} \sin u \cos v + \frac{k}{c^2} \cos u,$$

$$\frac{g^2}{a^2} + \frac{h^2}{b^2} + \frac{k^2}{c^2} = 1, \dots \dots \dots (d.)$$

we find

$$R = \frac{2 B}{A}, \quad \text{and } U = 2 \iint \frac{B^2}{A^2} \sin u du dv.$$

Between  $u=0$  and  $u=\pi$ , R will be positive, and then negative up to  $u=2\pi$ . We must therefore integrate from  $u=0, v=0$  to  $u=\pi, v=\pi$ . If we leave out terms containing the first power of  $\cos u, \cos v$ , as these would give nothing in the value of the integral, we may make

$$B^2 = \frac{g^2}{a^4} \sin^2 u \sin^2 v + \frac{h^2}{b^4} \sin^2 u \cos^2 v + \frac{k^2}{c^4} \cos^2 u.$$

Put

$$\left(\frac{g^2}{a^4} + \frac{h^2}{b^4}\right) \sin^2 u + \frac{2 k^2}{c^4} \cos^2 u = m, \quad \left(\frac{h^2}{b^4} - \frac{g^2}{a^4}\right) \sin^2 u = n,$$

$$\left(\frac{1}{a^2} + \frac{1}{b^2}\right) \sin^2 u + \frac{2}{c^2} \cos^2 u = p, \quad \left(\frac{1}{b^2} - \frac{1}{a^2}\right) \sin^2 u = q;$$

and we have

$$B^2 = \frac{1}{2} (m + n \cos 2v), \quad A = \frac{1}{2} (p + q \cos 2v).$$

We may obviously integrate from

$$u = 0, v = 0 \text{ to } u = \frac{\pi}{2}; \quad v = \frac{\pi}{2},$$

multiplying the integral by 4. Therefore

$$U = 16 \iint \frac{m + n \cos 2v}{(p + q \cos 2v)^2} \sin u du dv = 8\pi \int \frac{m p - n q}{(p^2 - q^2)^{\frac{3}{2}}} \sin u du$$

$$\begin{aligned}
 &= 4\pi \int \left\{ \frac{m+n}{p+q} + \frac{m-n}{p-q} \right\} \frac{\sin u \, du}{\sqrt{(p+q)(p-q)}} \\
 &= 2\pi \int \left\{ \frac{b}{a} \frac{c^4 g^2 \sin^2 u + a^4 k^2 \cos^2 u}{c^2 \sin^2 u + a^2 \cos^2 u} \right. \\
 &\quad \left. + \frac{a}{b} \frac{c^4 h^2 \sin^2 u + b^4 k^2 \cos^2 u}{c^2 \sin^2 u + b^2 \cos^2 u} \right\} \frac{\sin u \, du}{\Delta},
 \end{aligned}$$

where  $\Delta = (c^2 \sin^2 u + a^2 \cos^2 u)^{\frac{1}{2}} (c^2 \sin^2 u + b^2 \cos^2 u)^{\frac{1}{2}}$ .

If we transform this by making  $\sin^2 u = \frac{x}{c^2 + x}$ , and to abridge  $\Delta = \sqrt{(a^2 + x)(b^2 + x)(c^2 + x)}$ , we find

$$\begin{aligned}
 U &= \pi a b c \int \left\{ \frac{g^2}{a^2} + \frac{h^2}{b^2} + \frac{1}{c^2} \frac{a^2 k^2 - c^2 g^2}{a^2 + x} \right. \\
 &\quad \left. + \frac{1}{c^2} \frac{b^2 k^2 - c^2 h^2}{b^2 + x} \right\} \frac{dx}{\Delta},
 \end{aligned}$$

the integral to be taken from  $x = 0$  to  $x = \infty$ . In this value of  $U$ ,  $v$  has been taken in the plane of  $x$  and  $y$ . But if we make  $a$  and  $c$ ,  $g$  and  $k$ , and then  $b$  and  $c$ ,  $h$  and  $k$  change places, we shall have two other values of this quantity. Adding the three values together, dividing the sum by 3, and multiplying by  $\frac{\alpha \beta \gamma}{a b c}$ , we find in virtue of (d.),

$$V = \pi \alpha \beta \gamma \int \left\{ \frac{2}{3} + \frac{\frac{1}{3}a^2 - g^2}{a^2 + x} + \frac{\frac{1}{3}b^2 - h^2}{b^2 + x} + \frac{\frac{1}{3}c^2 - k^2}{c^2 + x} \right\} \frac{dx}{\Delta}. \quad (4.)$$

As this expression is complicated, we will find  $V$  by another method. From the formulæ already given, we easily perceive that

$$\left. \begin{aligned}
 \iiint \frac{(k-z) \, dx \, dy \, dz}{R^3} &= \iiint dR \sin u \cos u \, du \, dv \\
 &= \iint R \sin u \cos u \, du \, dv = 2 \iint \frac{B}{A} \sin u \cos u \, du \, dv \\
 &= \frac{4k}{c^2} \iint \frac{\sin u \cos^2 u \, du \, dv}{p+q \cos 2v} = \frac{4\pi k}{c^2} \int \frac{\sin u \cos^2 u \, du}{\sqrt{p^2 - q^2}} \\
 &= 4\pi a b k \int_0^{\frac{\pi}{2}} \frac{\cos^2 u \sin u \, du}{\sqrt{(c^2 \sin^2 u + a^2 \cos^2 u)(c^2 \sin^2 u + b^2 \cos^2 u)}}
 \end{aligned} \right\} (5.)$$

And if we transform this by making  $\sin^2 u = \frac{x}{\sqrt{c^2 + x}}$ , it will become

$$2\pi abc k \int \frac{dx}{c^2+x} \frac{1}{\sqrt{(a^2+x)(b^2+x)(c^2+x)}}$$

We have therefore in the general case

$$\left. \begin{aligned} -\frac{dV}{dg} &= \iiint_{R^3} \frac{(g-x) dx dy dz}{R^3} = 2\pi\alpha\epsilon\gamma g \int \frac{dx}{(a^2+x)\Delta} \\ -\frac{dV}{dh} &= 2\pi\alpha\epsilon\gamma h \int \frac{dx}{(b^2+x)\Delta}, \\ -\frac{dV}{dk} &= 2\pi\alpha\epsilon\gamma k \int \frac{dx}{(c^2+x)\Delta}. \end{aligned} \right\} (6.)$$

If we multiply the first of these by  $dg$ , the second by  $dh$ , and the third by  $dk$ , and add them, we may certainly make the integral of the result

$$V = \pi\alpha\epsilon\gamma \int \left\{ \lambda + 1 - \frac{g^2}{a^2+x} - \frac{h^2}{b^2+x} - \frac{k^2}{c^2+x} \right\} \frac{dx}{\Delta},$$

whether  $g$ ,  $h$ , and  $k$  be implicitly contained in (6.) or not, if we suppose  $\lambda$  a function of  $g$ ,  $h$ ,  $k$ . Let us take the partial differentials of it relative to  $g$ ,  $h$  and  $k$ , and equal them to their values given in (6.), we shall have, leaving out quantities which destroy one another,

$$\begin{aligned} \frac{d}{dg} \int \frac{\lambda dx}{\Delta} &= \int \frac{d}{dc^2} \left\{ \left( 1 - \frac{g^2}{a^2+x} - \frac{h^2}{b^2+x} - \frac{k^2}{c^2+x} \right) \frac{1}{\Delta} \right\} \frac{dc^2}{dg} dx \\ &= \int \frac{d}{dx} \left\{ \left( 1 - \frac{g^2}{a^2+x} - \frac{h^2}{b^2+x} - \frac{k^2}{c^2+x} \right) \frac{1}{\Delta} \right\} \frac{dc^2}{dg} dx \\ &= \frac{1}{abc} \left( \frac{g^2}{a^2} + \frac{h^2}{b^2} + \frac{k^2}{c^2} - 1 \right) \frac{dc^2}{dg} = 0. \end{aligned}$$

In like manner,

$$\frac{d}{dh} \int \frac{\lambda dx}{\Delta} = 0, \quad \frac{d}{dk} \int \frac{\lambda dx}{\Delta} = 0.$$

Therefore  $\int \frac{\lambda dx}{\Delta}$  is a quantity independent of  $g$ ,  $h$ ,  $k$ , and also of  $c$ . It must be remembered in this process that  $a^2 = c^2 + a^2 - \gamma^2$ ,  $b^2 = c^2 + \epsilon^2 - \gamma^2$ . But we cannot determine  $\lambda$  by making  $g$ ,  $h$ ,  $k$  and  $c$  infinite, for then both sides of the above equation will vanish independently of any particular value of this quantity. Nor is it easy to see how we can determine it from hence. But we see immediately from (6.) that

$$\begin{aligned} \pi\alpha\epsilon\gamma \int \left\{ \frac{a^2}{a^2+x} + \frac{b^2}{b^2+x} + \frac{c^2}{c^2+x} \right\} \frac{dx}{\Delta} \\ = \frac{1}{2} \iiint_{R^3} \left\{ \frac{a^2}{g}(g-x) + \frac{b^2}{h}(h-y) + \frac{c^2}{k}(k-z) \right\} \frac{dx dy dz}{R^3} \end{aligned}$$



$$\begin{aligned}
 &= \frac{\alpha \epsilon \gamma}{2abc} \iint \left( \frac{a^2}{g} \sin u \sin v + \frac{b^2}{h} \sin u \cos v + \frac{c^2}{k} \cos u \right) dR \sin u du dv \\
 &= \frac{\alpha \epsilon \gamma}{abc} \iint \left( \frac{a^2}{g} \sin u \sin v + \frac{b^2}{h} \sin u \cos v + \frac{c^2}{k} \cos u \right) \frac{B}{A} \sin u du dv.
 \end{aligned}$$

Taking account of such terms only as give value in the final result, this reduces to

$$\begin{aligned}
 \iint \frac{\sin u du dv}{A} &= 2 \iint \frac{\sin u du dv}{p + q \cos 2v} = 2 \pi \int \frac{\sin u du}{\sqrt{p^2 - q^2}} \\
 &= 2 abc^2 \pi \int_0^{\frac{\pi}{2}} \frac{\sin u du}{\sqrt{(c^2 \sin^2 u + b^2 \cos^2 u)(c^2 \sin^2 u + a^2 \cos^2 u)}} \\
 &= abc \pi \int \frac{dx}{\sqrt{(a^2 + x)(b^2 + x)(c^2 + x)}}.
 \end{aligned}$$

Multiply this by the factor  $\frac{\alpha \epsilon \gamma}{abc}$ , which for convenience has

been left out of the last steps, and we have  $\pi \alpha \epsilon \gamma \int \frac{dx}{\Delta}$  for the result. By means of this result, (4.) becomes immediately

$$V = \pi \alpha \epsilon \gamma \int \left\{ 1 - \frac{g^2}{a^2 + x} - \frac{h^2}{b^2 + x} - \frac{k^2}{c^2 + x} \right\} \frac{dx}{\Delta}. \quad (7.)$$

It seems not a little strange that this result should not by the direct method have come out at once. I might give many more examples in application of the theory given at the beginning of this paper; but the above may suffice, the subject being well understood.

Gunthwaite Hall, January 27, 1846.

LXI. *On the Storm-Paths of the Eastern Portion of the North American Continent.* By WILLIAM RADCLIFF BIRT.

*To the Editors of the Philosophical Magazine and Journal.*

GENTLEMEN,

**I**N the opening articles of the first and second numbers of the second series of the American Journal of Science, Mr. Redfield has contributed some further important information to our present stock of knowledge respecting the storm-paths of the eastern portion of the North American continent and its adjacent seas. The chart No. 1, illustrating the articles, exhibits the tracks of sixteen hurricanes; eight of these form an interesting group distinguished by this striking characteristic: the paths are semi-orbital, presenting in each case

the half of an ellipse of greater or less eccentricity; the apices are confined to the thirtieth parallel of latitude, and are more acuminated as they recede from the fiftieth degree of longitude west of Greenwich. The remaining eight depart considerably from this type: four appear in some measure to approximate to it, but most probably with very acuminated apices. A most remarkable one, of a parabolic form, with its apex on the twenty-sixth parallel, was observed in October 1837 (XV.). In October 1842, the north-western portion of an elliptic path with the apex still lower was traced (XIII.). The storm pursuing this path is particularly discussed in the continuation of Mr. Redfield's paper, and its identity with the northers of the Mexican coast insisted on. Mr. Redfield also speaks of other storms that had exhibited the character of northers in the Gulf of Mexico, and afterwards presented all the features of Atlantic storms. Speaking of a storm that occurred in October 1837, he says, "This norther of the Mexican coast had become in due course of progression an Atlantic storm" (see *American Journal of Science*, second series, No. 2, March 1846, p. 166). In October 1844, a storm passed nearly in a direct line from the Gulf of Honduras to Newfoundland (XIV.); and to the gale which is first discussed in the articles before us (XII.), a nearly direct westerly course has been assigned from all the observations that have come to hand.

It is not my intention in the present communication to enter into any examination of the particular gales above enumerated, or to attempt to substantiate or refute either the one or the other of the rival theories which have been offered as an explanation of the phænomena. Those of your readers who are acquainted with Col. Reid's work, are aware that he has most ably discussed the rotatory theory, and for the centripetal theory, I beg to refer to various papers in the *American Journal of Science*. I apprehend that the labours of Redfield, Loomis, and Espy in the United States, Col. Reid in England, and Piddington and Thom in India, have brought the inquiry to that point at which it becomes essential to connect it with some kindred branch of science, in order to see our way clear in resolving the interesting problems that suggest themselves, to strike out a path for working energetically in surmounting the obstacles that still retard our progress in becoming acquainted with the dynamical system of our atmosphere, and in successfully removing the desiderata as they arise.

Among the desiderata of these phænomena, and by far the most important, will be found their origin and final disappearance. Sir John Herschel has suggested, in his Report on

Meteorological Reductions\*, that they may be produced by the crossing of two large atmospheric waves moving in different directions. Some interesting evidence of the existence of such atmospheric waves has been brought forward at late Meetings of the British Association. This evidence rests entirely on the barometric affections of the atmosphere over a large tract of country. In order, therefore, to extend our knowledge of the rotatory gale (Redfield) or the centripetal hurricane (Espy), especially with regard to the desiderata above-mentioned, I apprehend it will not only be important, but absolutely necessary to accompany all the observations of the direction, force, and variation of the wind with barometric readings. These readings, however, must not be confined to the mere period of the passing of the gale; evidence has been adduced of the passage of large atmospheric waves occupying from fifteen to seventeen days between the anterior and posterior troughs, or between successive crests; so that in order to detect the origin of a gale arising from the intersection of two waves, to trace it throughout its destructive course and to observe its final disappearance, it will be essential to discuss the entire system of observations appertaining to both waves, not only in time, but also in space. Our meteorological observations are approaching a degree of uniformity and system that bids fair for uniting these kindred inquiries. I have now before me Prof. Loomis's interesting and ably conceived charts for exhibiting the principal phænomena during the passage of two storms over the United States in February 1842, in which he clearly shows the barometric, thermometric, and anemonal phænomena, and exhibits in a very striking manner the extent of rain, snow, cloud, and blue sky over the whole of the United States twice in the course of each day that the storms prevailed. The barometric phænomena are shown by *lines of equal pressure*; and I apprehend that these lines of equal pressure indicate, especially in one of the two cases, that *two waves* passed over the United States; that as the posterior slope of one wave passed off, the anterior slope of a wave of a different system approached; and that in the point of intersection the storm raged. Should the entire barometric observations taken over the United States on that occasion, February 1 to 4, 1842, support the theory of atmospheric waves, I apprehend Sir John Herschel's suggestions will be partly realized.

I cannot close this notice without adverting to a most important desideratum in this interesting inquiry. Mr. Red-

\* Report of the British Association for the Advancement of Science, 1843, p. 100.

field's storm-paths are confined to the westward of the fiftieth meridian, and he is only able to give one-half of the ellipse in any case. Nearly the whole breadth of the Atlantic is a perfect blank in any storm-map. Surely several captains may be found who will be happy to make observations on their passages out and homeward, and transmit them to head quarters, either in London or some principal city in America; the only additional information required to that furnished by their logs, will be the altitudes of the barometer at given hours of the day and night, with its attached thermometer. The American steamers might furnish important and valuable information.

I have the honour to be,

Gentlemen,

Your very obedient Servant,

Cambridge Heath, April 22, 1846.

W. R. BIRT.

LXII. *On the first introduction of the words Tangent and Secant.* By Prof. DE MORGAN\*.

**A**BOUT the meaning and origin of the word *sine*, there is now no discussion. The words *tangent* and *secant*, though clear as to their meaning, have an origin which is not mentioned by historians. Nobody, in fact, knows where they came from; very few people care.

As it may appear surprising, and perhaps even doubtful, that the first inventor of names now so universally recognised should be unknown, I will begin by stating, that neither Weidler, Heilbronner, Montucla, nor Delambre, mentions the work in which the words first appear. Montucla does not allude to the question of the invention of these terms. Delambre does it, as follows, in a manner which shows that his impression on the subject was not founded on anything precise.

In speaking (*Astr. Moy.*, p. 437) of Vieta's mode of designating the trigonometrical functions, he informs us that Vieta, after stating that no elegant name had been given to what we now call the table of tangents, proposed to continue the use of the term *tabula fecunda*, which had been given by Regiomontanus. "Ce qui," continues Delambre, "n'est pourtant pas plus élégant que le mot *tangente* qu'il réproûve; quant aux *secantes*, il veut qu'on les appelle *hypoténuses des féconds*. Ces dénominations n'étaient pas faites pour être accueillies; on s'est décidé pour ce qui était plus naturel, plus commode, et même plus élégant, quoiqu'en dise Viète." Here it is plain that he thinks the word *tangent* to have been one of those which Vieta spoke of, when he threw his imputation of want

\* Communicated by the Author.

of elegance over all which existed. On looking at the paragraph of Vieta's *Liber Inspectionum*, which Delambre was then describing, we find not the smallest allusion to the word tangent, nor to any name for the table, except the old one of *tabula fecunda*. So that Delambre must have proceeded upon a general impression, that the word tangent was in use in Vieta's time. But as he does not quote any authority for this impression, though much given to incidental allusion to one writer in his description of another, it is not necessary to give it any weight in the face of the positive evidence which I shall produce. I say this, because an impression on the mind of Delambre as to a usage deserves more consideration than the same thing in the case of any other mathematical historian. His memory might fail on an isolated fact, or his information might be incorrect on books which he had not seen; but he was occupied at each one time with masses of writers of one period, and came to each author fresh from that author's own contemporaries, and frequently from *very* close reading of them.

After writing the above paragraph, I happened to find a passage in a later writing (the *Responsorum liber octavus*, published in 1593) which might have left the above impression on Delambre's mind. Here Vieta distinctly names and objects to the words *tangent* and *secant*, and proposes to call the former *prosines* or *amsines*, and the latter *transsinuous* lines. Laugh if you will, he says, at the allegory of the Arabs (meaning the use of the word which is correctly Latinized by *sinus*), but either adopt it altogether, or reject it altogether. This passage strengthens the presumption, that when Vieta wrote the *Canon*, &c. he had never heard the words tangent or secant. The same impression on the part of Delambre occurs again in speaking of Pitiscus (*Astr. Mod.* ii. 33), when he says, "Il a eu le bon esprit de n'imiter ni Viète, ni Rhéticus; il a conservé les noms de sinus, de tangentes et de sécantes." But where either Rheticus or Vieta (in 1579) was to have found the last two names, we are never told.

The *Canon Mathematicus* of Vieta, to which the *Liber Inspectionum* above-mentioned is an appendix, was published in 1579. The work in which tangents and secants are first mentioned under those names, was published four years after.

Its author was Thomas Finck, of Flensburg in Denmark, who was successively professor of mathematics, rhetoric, and medicine at Copenhagen, where he died in 1656, at the age of ninety-five (*Alkin, Gorton, and Biogr. Univ.*): there are references to his astronomical observations in Tycho Brahé. The work we speak of was published at Basle, when he was a

student there (and where two years before he had published an ephemeris) at the age of twenty-two. It is "*Thomæ Finkii Flenspurgensis Geometriæ Rotundi libri xiiii. Ad Fridericum Secundum, Serenissimum Daniæ et Norvegiæ Regem, &c. Cum Gratia et Privileg. Cæs. Majest. Basileæ, per Sebastianum Henricpetri:*" quarto. The colophon is "*Basileæ, per Sebastianum Henricpetri, anno salutis humanæ M.D.LXXXIII. Mense Augusto.*" The work itself is well-worthy of description for its contents, independently of its being the production of so new a student. But I am here only concerned with the table of sines, tangents, and secants, and with the audacity of the young gentleman who presumed to alter the established names of the latter two. In page 73, after drawing the circle, &c., he thus proceeds:—"Erit AI tangens datæ peripheriæ. Sic vocare placuit quia sit perpendicularis extremæ diametro. . . . . Geometria ipsa commodum suppeditavit nomen; nec aliunde adferri commodius poterit. Nam quod quidem numerum fœcundum rectam AI vocant, id ii videant quomodo defendant: mihi non probabunt. Damus aliquid peritissimo illi artificio Regiomontano homini Germano: qui primus hujus vocabuli author dicitur: damus etiam aliquid receptæ consuetudini. Verum id non facile damus ut verba ea in usu retineamus quibus elegantiora, breviora, significantiora, veriora habeamus."

At page 76 the secant thus makes its appearance:—"Sic peripheriæ AE secans est OEI, nempe radius OE continuatus in terminum tangentis E cum continuatione EI. Et hoc nomen huic recte accommodatum putamus. Joachimus Rheticus hypotenusam trianguli recti vocat respectu anguli recti ad A cui subtenditur. Verum cum referatur non ad angulum rectum, sed angulum in centro ad O, hoc est arcum AE: an non potius arcus secans dicatur quam recti anguli hypotenusam judicent alii. Maurolycus canonem Rhetici paulo mutatum in Messanensi Menelai editione, nomine etiam mutato edidit: et beneficium vocavit: tanquam recta OI seu numerus hanc definiens beneficium diceretur. At recta OI non magis est benefica, quam recta AI fœcunda." And again (p. 130), "Sequitur canonis hujus triangulorum pars altera quæ vulgo canon fœcundus, nobis canon tangentium dicitur: et canon hypotenusarum Rhetico, nobis canon secantium vocatur."

From these passages it is obvious that Finck gives the words for the first time, and defends them as suggestions of his own. How then does it happen that his right to them is entirely unknown? Did his learned contemporaries dislike to owe their terms to a youth of twenty-two years? or did the south-erns wish to avoid acknowledging the young Dane?

The first after Finck who used the words tangent and secant was the celebrated Jesuit Clavius, whose edition of Theodosius\*, with trigonometry and tables attached, was published, according to Lalande, in 1586. Blundeville, who copied these tables into his *Exercises*, and who is, as far as I can find, the first Englishman who gave complete trigonometrical tables, cites this date. These tables of Clavius are those of Finck, with the use of the terms tangent and secant; but Finck's name is entirely suppressed. Clavius does not mention the name of the condemned Protestant Rheticus: it is not surprising that he should have served the other Protestant, Finck, in the same way. I do not suspect Clavius of wanting to pass the work of another as his own; he mentions Regiomontanus and Purbach freely enough, and excludes none but persons whom a reputable Jesuit could not name, as Protestants and Copernicans. But I proceed to make good my assertion.

The tables of Clavius are to the same radius and interval as those of Finck. The sines are avowedly from Regiomontanus: the latter gave differences to every ten seconds; Clavius does the same. But Finck gave no differences: Clavius gives no differences to his tangents and secants. In the tables of this period, the tangents and secants in the last degree were often very wrong, having hardly one of what we call the decimal places right. Clavius agrees with Finck in every decimal place, and differs from Vieta and what had *then* been published by Rheticus. For instance, we take the tangent and secant of  $89^\circ 50'$ .

	Tan $89^\circ 50'$ .	Sec. $89^\circ 50'$ .	Date.
Correct value .....	343·77371...	343·77516...	
Vieta .....	343·77371...	343·77516...	1579
Rheticus .....	343·7829002	343·7843784	1551
{ Finck.....	343·7829002	343·7843546 }	1583
{ Clavius .....	343·7829002	343·7843546 }	1586

And it is the same throughout: whenever Finck differs in two or three decimal places from Rheticus, so does Clavius in the same manner. And whereas Vieta and Rheticus have the semiquadrantal form, Clavius agrees with Finck in retaining the quadrantal form.

There is a particular reason why Finck should differ from Rheticus in the secants. The former used the reciprocal of the cosine carried to more figures than it would give truly; the latter demonstrated the formula

\* I have before me the tables in the complete edition of Clavius's works, and have never seen the original edition.

$$\sec \theta = \tan \theta + \tan \left( 45^\circ - \frac{\theta}{2} \right),$$

and used it. Clavius gives the same demonstration of the same formula. There is then no doubt that the celebrated tables of Clavius, the first (I believe) introduced into this country, are no other than the tables of Finck, deprotestantized by the substitution of the name of Clavius for that of Finck. If our Elizabethan mathematicians had known this, they would not have let it pass unnoticed.

The tables of Clavius were copied by Lansberg and Magini (1591 and 1592), both of whom omit all mention of Finck, though the second gives a list of the names which his several predecessors gave to the trigonometrical lines. So completely did the last name disappear from history, that it is not mentioned in its proper volume of Delambre (the *Astron. Moyenne*), except in the index, in which it is stated, speaking of the tables of secants, that "Rhéticus l'a étendue d'abord à toutes les minutes; et c'est ainsi que Finckius l'a reproduit en 1583 en citant Rhéticus mort en 1574." This is incorrect; Rheticus published nothing closer than to ten minutes, and Finck, we may suppose, could not have found out Valentine Otho and the manuscripts at the age of twenty-two; he would have made his final secants more correct if he could have done so. When he cites Rheticus, it is for the name which the latter adopted, not for any mode of calculating; and as I have stated, he made his own secants by his own method.

This work of Finck is so clear and concise, and so much above the usual writing of the sixteenth century, that its author ought to rank very high among the secondary authors of that period.

---

Being engaged in an attempt to trace the early progress of trigonometrical tables in England, I annex the results which I have obtained, in the hope that some of your readers may be able to furnish additional information. I am not aware that any one has ever investigated the point.

Thomas Digges and John Dee, in their several works on the new star in Cassiopea (both published in 1573, and in Latin), mention and use sines, but refer to foreign tables; Digges to the ten-minute canon of Rheticus, Dee to Regiomontanus. William Burroughs, in his tract on the Variation (written in 1581), mentions and uses tables of sines, but describes the doctrine of "signes and triangles" as new and strange to English ears. He professes his intention of interpolating the ten-minute canon, and publishing it, if his pur-



pose be not forestalled by other publications. This he did not do; and to the edition of 1614 the editor appended the tables which Ralph Handson had published in his translation of the Trigonometry of Pitiscus (of the first edition of which I have not found the date). The first sines actually published, as far as I can find, were those at the end of Thomas Fale's *Horologiographia*, first published in 1593; they are to minutes, with a radius of 100000. The first complete canon which I can find is that of Blundeville, in his Exercises, first published in 1597. They are taken from Clavius, and are to every minute, with a radius of ten millions. These Exercises went through seven editions at least, and were latterly corrected from Pitiscus. John Speidell, well-known afterwards for his logarithms, published a small table in 1609, to every ten minutes, and to a radius of 1000. Briggs began his calculation of sines about 1600, in ignorance, we may suppose, of the appearance of the *Opus Palatinum* four years before. This is all I have been able to find on the matter.

There is a work expressly on the history of the trigonometrical canon, which is sometimes cited by foreign writers; it is by Frobesius, and was printed at Helmstadt about 1750; but I cannot find any copy of this work in London.

### LXIII. Complete Collection of Kepler's Works.

By Dr. J. LHOTSKY,\*

**P**ROFESSOR Frisch of Stutgardt has recently published a programme of his intended collection of the works of the great German astronomer and philosopher, which, like a splendid luminary, will enliven the dim polygraphy of the present age. Considering Kepler as the real founder of modern astronomy, the collecting of his works (many of them very rare) is a tribute most due to such great merit. Our present epoch seems especially adapted to such an undertaking. Monuments are everywhere raised to the honour and memory of men who have deserved well of humanity or of their country, in one or another department of human knowledge or enterprise. In following up the track marked out by Kepler in astronomical science, a degree of accuracy and perspicuity has been acquired, which is one of the proudest trophies of the human mind. In such a time, it is impossible that the claims and memory of Kepler could be kept in abeyance any longer. He, the modest searcher and deep thinker, ought to obtain his share of attention and general recognition, so long with-

\* Communicated by the Author.

held from him,—whose discoveries have been confirmed in so splendid a way.

Something, it is true, has been done in the way of atonement towards his long-neglected memory; and it is now thirty-seven years since over his grave at Regensburg (his native city), the image of the great man was placed under the cupola of a monument, on the anniversary of his birth-day, amid the roar of cannon. It cannot, however, be said that this slight token of gratitude could suffice to the memory of one whom his coevals left in misery and distress, while occupied with the examination of the very innermost secrets of science; while at the same time the produce of a painful and thorny life, his splendid works, remained unknown and forgotten. It is true, his name is on the lips of every astronomer and philosopher; the three great *Keplerean laws* are yet the main basis of the knowledge of the heavens; still his works moulder in dust and oblivion. On the other hand, it must be acknowledged that the present position of mathematical science is far different from what it was in the seventeenth century; and problems are now solved with facility and speed which then occasioned much labour; and many a deep axiom and saying of Kepler will be more easily appreciated if presented to the reader in a more modern garb. But it is not merely the contents, but even the *form* and style of the immortal astronomer's works, which imparts value to them; and it requires but little attention to become familiar with that form, albeit hidden and enigmatic.

There are two reasons, however, which have hitherto prevented the greater spread of Kepler's works,—their *rarity* and their *external appearance*. There is hardly a library in Europe where *all* the works of Kepler are to be met with, and many where even the most important are wanting. The cause is obvious. In the then condition of typography and publishing, only a few copies could be printed, and of those many were lost in conveying them about the country and by other accidents. Thus, for instance, Kepler's *Harmonie* and *Astronomia Nova* are so scarce, that only the largest libraries can boast of their possession. The next cause of the neglect of our author's works, is the wretched type, bad paper, and the improper size of many of them. The figures, moreover, are so badly designed, and the letters thereon so indistinct, that they cannot be read without difficulty. These reasons will be deemed sufficient for making a *collection of Kepler's works*. It was the late Prof. Pfaff at Eslingen, who, in 1810, first conceived this design, which, however, did not come to maturity. Still, constant communications with this gentleman

kept the conviction of the importance of such an undertaking alive, and a further investigation of the great work increased the interest for the old astronomer and philosopher. The present editor, however, Prof. Frisch, did not long enjoy the assistance of either Pfaff, or of the great philologist, Prof. Kopp, both being carried off by premature death. The labour was subsequently much aided by the head librarians of Stuttgart and Tübingen, who defrayed the preparatory expenses of the undertaking. Of especial use also was the library of Reütlingen, which contains a very complete collection of the mathematical works of the sixteenth century. All this must of necessity have led to the inquiry after the original MSS. of Kepler, a notice of which is to be found in Murr's *Journal für Kunstgeschichte und allgemeine Literatur*, vols. iii. and xvii. It is to the following effect:—Kepler's son Ludwig, who died in 1663, had preserved the MSS. of his father with the intention of publishing a selection from them. In fact, there appeared in 1634, through his endeavours, a work which, however, the father had prepared for the press, viz. *Somnium sive de Astronomiâ lunari*. All the rest remained unused, most probably because Ludwig, who was a physician, could not understand the problem of his great progenitor. From him the MSS. went to the celebrated Selenographer Hevel, and thence to his son-in-law, the common-councilman Lange at Dantzic. Of Lange they were bought in 1707 by the Leipzig mathematician, Hansch, for 100 florins. They consisted of twenty-two folio volumes, which, besides the drafts of several works already printed (for instance, the *Harmonie*, the Rudolphine Tables, &c.), contained the correspondence of Kepler with many distinguished personages, several astronomical works merely begun, and a host of miscellaneous notices. Hansch intended to publish these MSS. in a splendid form, but was only able to begin this undertaking (too costly in the form he had projected it), and the *Epistolæ ad Keplerum scriptæ, insertis ad easdem responsionibus Keplerianis*, which was patronized by the emperor Charles VI., was the only result he ever achieved. The MSS. thus published, are to be found in the Imperial Library of Vienna. As Hansch fell into poverty, he was obliged to pledge the MSS.; and as he could not redeem them, one Etringer, of Frankfort on the Maine, redeemed them for 128 florins. Thence they came (probably by inheritance) to a Mrs. Trümmer at Frankfort, where they remained unknown until 1770, when Christopher de Murr, a man deserving well of literature, called attention to them. For the purpose of recovering them from oblivion, he addressed himself to several astronomers, as Mayer, Bernouilli,

Kästner, &c.; but besides laudatory commendations on his undertaking, they could not afford any substantial aid. Consequently Murr sent the catalogue of the MSS. to St. Petersburg, and solicited Euler for his intercession. On the recommendation of the latter and other savants, they were purchased in 1774 for the St. Petersburg Imperial Academy. The academicians Euler, Krafft, and Lexell received orders to peruse the MSS. and to select those worthy of publication. Lexell began the revision of a nearly completed work of Kepler's, on the motion of the moon, entitled *Hipparchus*; but there it ended, and neither the work nor the promised completion ever saw the light. These MSS. have ever since reposed, as Prof. Krafft writes, "an ornament of the Petersburg Library,"—useless, unknown. Prof. Frisch took great pains to obtain these MSS. Introduced by Baron de Regendorff, Russian Minister at Stutgardt, and Prof. Schelling, he addressed himself to the imperial government, and received the assurance that the use of them would be granted to him; and from the known scientific munificence of the Petersburg cabinet, it is to be hoped that his further request for the loan of the MSS. will be shortly granted.

As to the plan on which the works are to be edited, Prof. Frisch makes the following statement:—The original text will remain unchanged, except where palpable error has crept in, as it is intended that Kepler shall appear throughout in his truest form. The notes will be as few and concise as possible, in order not to increase the bulk of the work; they will treat either on historical points, or explain difficult passages. As most of the works of Kepler are in Latin, the adoption of that language for the notes has been deemed expedient. The introduction which will precede the whole is to contain a survey of the condition of mathematical and natural science in the century preceding the life of Kepler, and to this will be attached his biography, mostly relating to his scientific labours. But as it is possible that some of the MSS. at St. Petersburg may contain materials for the elucidation of this subject, the compiling of it will be deferred until it shall be ascertained whether permission will be granted for the use of them.

The works will, as far as possible, be printed in the order in which Kepler composed and published them. But those relating to chronology will be put together, and those consisting mostly of numerals, as the Ephemerides, the Rudolphine Tables, and the work on Logarithms, will form the last part of the collection. Prof. Frisch concludes his programme by calling upon all friends of science to aid him in an under-

taking the difficulty of which he is perfectly aware of. Every communication made to him will be received with thanks. The owners of MSS. of Kepler especially might greatly assist the undertaking by the loan of them, or the transmission of accurate copies. Even of the printed works of Kepler, Prof. Frisch has not been able to see the following:—

1. Almanack of the Year 1594\*.
2. *De Fundamentis Astrologiæ*, Prague 1602.
3. *Epistola ad rerum cœlestium amatores de Salis deliquio*, Prague 1605.
4. *Dissertatio cum nuncio sidereo, &c.*, Florence 1610, 4to. [The edition of Prague 1610, 4to, and Frankfort 1611, 8vo, are at hand.]
5. *Responsio ad Epistolam Bontschii*, Sagan 1609.

In conclusion, Prof. Frisch requests the loan, or offers the purchase of any of these works of Kepler.

A considerable time having elapsed since the programme of the Stutgardt Professor was published, the aim of which undoubtedly was to excite attention in, and to obtain aid also from, *this country*, which has associated itself in every useful undertaking all over the world, I have now been induced to take up this subject, as my studies at Prague had led me often in the very track of the life of Kepler, who remained eleven years in that city. But the attention of an English public cannot and ought not to be called for without searching for some *connecting link* between the activity of Kepler and those of English savants. A brief search, indeed, showed me that such existed, and I have been able to complete and correct the programme of Prof. Frisch. If we refer to the names of the various possessors of Kepler's MSS., we find that it was Hevel who obtained them from Kepler's son, and we find the number of volumes or fasciculi to have been twenty-two. It was not to be supposed that the Stutgardt Professor should have known the long array of volumes composing the Philosophical Transactions of the Royal Society, else he would have found a letter of that very Hevel addressed to the Society (or their publisher) on these very MSS., in which they are called the famous Kepler MSS. (vide Phil. Trans. vol. ix. No. 102, p. 27, anno 1674). The title of this remarkable communication is as follows:—“An Extract of M. Hevelius's Letter, lately written to the publisher, concerning the famous Kepler's MSS., together with some Observations of his about the

\* This is the first thing Kepler printed at Grätz. It is very probable that a copy of this work may exist safe in some neglected collection of rubbish.

Use of Telescopic Sights in Astronomical Observations." In looking over this letter, we find that the number of fasciculi in Hevel's possession was not twenty-two, as stated (or at least implied) by Prof. Frisch, but twenty-nine fasciculi. One thing, moreover, is certain, that Hevel must have possessed all the MSS.; deriving them from a source where certainly they had received the highest possible attention. In perusing, therefore, the catalogue of these MSS., as given by Hevel in his letter to the Royal Society, we shall find that Prof. Frisch's hopes or expectations as to what the St. Petersburg documents may contain, can be at once answered, because Hevel could never have had *less* than what is at St. Petersburg *now*; though he might have had more, which, however, would make the case worse. A very short time, in fine, will now set the matter at rest. In regard to Kepler's life, Hevel says as follows:—"At Kepleri vitam studio conscriptam non invenio; interim plurima notatu dignissima, vitam ejus spectantia passim notavi, ex quibus vita ejus possit haud obscure depingi. Quæ verò in specie ex scriptis ejus penes me habeo, catalogus hicce indicabit" (Ibid.). Whatever fates these MSS. might have subsequently undergone, Hevel's Catalogue must ever be considered the most complete. The extract from Hevel's letters to the Royal Society (as printed in the Phil. Trans.) does not imply any especial offer or request. But there exists among the MSS. of the *British Museum*, another document relating to these MSS., and this is an *autograph* letter of Hansch to the Royal Society; it is dated Vienna, November 20, 1734. There Hansch speaks of twenty-two fasciculi, and as this was written *after* his *Epistolæ ad Keplerum* were published, it may be presumed that seven might have merged (whether entirely or partially is not known) in this undertaking. Hansch's letter contains *also* a list of the Kepler fasciculi, but they seem to have been re-arranged, as the contents of most, as given by Hevel and Hansch, do not correspond. It is, moreover, curious that Prof. Frisch says that the *Epistolæ ad Keplerum* were all that Hansch published, while the letter in the British Museum says, "Præter Epistolæ, quæ in folio charta augusta prodierunt, et Librum singularem de Calendario Gregorio quem Ratisbone 1726—in folio pariter typis imprimendum curavi—reliqua MSS. REGIAM desiderant munificentiam." Hansch's letter is entirely lachrymose and supplicatory; and it is a pity to perceive that these MSS. had something ominous in them, as not only their author, but even several of their subsequent owners fell into deep distress.

Having been so far successful in my research, I resolved to

Rev. J. Challis on the Aberration of Light.

see whether some of the works of Kepler, which Prof. Frisch could not discover in Germany, might not be found in our libraries, which certainly are surpassed by the richness of *especial* departments of those of the *respective* countries; and none would expect, for instance, to find more Austrian Incunabula in English libraries than there are in Vienna, &c. But taking the biographical opulence at a fair average, the balance will not be unfavourable for this small and insulated empire. The very first glance I cast in the Catalogue of the British Museum (even in its present transitory state) was encouraging, as I found No. 5 of Prof. Frisch's *Desiderata*. The full title of this little rarity is as follows:—“*Joannis Kepleri Mathematici ad Epistolam Clarissimi Viri D. Jacobi Bartschii Laubani Medicinæ Candidati Præfixam Ephemeridi in anno 1629 Responsio: de Computatioe et Editione Ephemeridum, Typis Saginensibus 1629.*” It is a small 4to pamphlet of only eleven pages, printed on paper and with a type of the then current publications of the day. The conclusion is so characteristic of the man, that we shall translate it:—“But while the storm is raging, and the shipwreck threatens public affairs, nothing remains to us but to let the anchor of our innocuous studies go down to the profound of eternity! Given at Sagan in Silesia, with our own types, anno 1628.” It is known that Kepler had been in some relation with the great Wallenstein, and the place of printing is one of the possessions of the great warrior, he having been Duke of Sagan. The name of the duke is also mentioned in the contents of the work.

London, April 15, 1846.

---

LXIV. On the Aberration of Light, in Reply to Mr. Stokes.

By the Rev. J. CHALLIS, M.A., Plumian Professor of Astronomy in the University of Cambridge.

To the Editors of the *Philosophical Magazine and Journal*.

GENTLEMEN,

I HAD reason to expect, when I made my last communication on the Aberration of Light, that I should not have occasion to trouble you again on this subject. Mr. Stokes's remarks in the April Number compel me to say a few words more.

I can assure Mr. Stokes that I take the aberration of light in its usual acceptation, and I have no doubt that he does also. The difference between us is not in the thing explained, but in the *principles* of our explanations. My explanation, which is very simple and brief, being entirely contained in

*Phil. Mag. S. 3. Vol. 28. No. 188. May 1846.*      2 E

page 91 of the February Number of the Philosophical Magazine, does not even suppose the existence of an æther. On the contrary, Mr. Stokes's rests both on the hypothesis of an æther and on a gratuitous and very particular supposition respecting its motion. By Mr. Stokes's admission, I have shown on my principles, that if to the earth's way, as measured by an astronomical instrument, be added in the same plane an angle equal to the product of the ratio of the earth's velocity to the velocity of light and the sine of the earth's way, we obtain the direction in which light from a star progresses just before it enters the eye. By measures taken with astronomical instruments, it is found that if to the same angle in the same plane be added the product of  $20''\cdot42$  and the sine of the earth's way, the mean place of the star is obtained. [The numerical quantity is that adopted in the British Association Catalogue of Stars.] Now it happens that the ratio of the earth's velocity to the velocity of light is known independently of the above-mentioned measures, by observations of the eclipses of Jupiter's satellites. Delambre states (*Abrégé d'Astronomie*, p. 493), that by very exact and extensive researches on the satellites of Jupiter, he found for this ratio  $20''\cdot25$ . The close approximation of these numerical values justifies me in concluding that the light from the star enters the eye, *quam proxime*, in the direction of a line drawn to the eye from the star's mean place; or, in Mr. Stokes's notation, that  $s_2$  coincides very nearly with  $s$ . Mr. Stokes appears to be dissatisfied because this inference is not deduced by theory alone. I conceive that it is not the less certain because it is deduced from *facts*; and as Mr. Stokes does not contend that it is not *true*, I need say no more on this point.

The "confession" which Mr. Stokes says that I made, I am ready to make again. I allow that, anterior to the above comparison with the result of astronomical measures, it could not be anticipated that aberration would be wholly accounted for by the motion of the earth and the finite velocity of light, without reference to any theory of light. The comparison shows that it *is* so accounted for, and the inevitable consequence is, that any explanation which rests on a hypothetical motion of the æther, must be *fictitious*.

I really think that I have now said quite enough in defence of a very unexceptionable piece of reasoning, and if Mr. Stokes should have anything further to urge, I must decline answering it.

I am, Gentlemen,

Your obedient Servant,

Cambridge Observatory,  
April 11, 1846.

J. CHALLIS



LXV. *On the Finite Solution of Equations.*

By JAMES COCKLE, M.A., *Cantab.*; *Special Pleader*\*.

[The subject concluded from p. 191.]

15. **L**ET  $\lambda', \lambda'', \dots, \mu$  be  $n$  unequal integers, then it might be shown † that  $x^\mu$  equals

$$p_0 + p_{\lambda'} x^{\lambda'} + p_{\lambda''} x^{\lambda''} + \&c.; \dots \dots \dots \text{(ae.)}$$

and, hence, that ‡  $\Lambda''' x^{\lambda'''} + \Lambda^{iv} x^{\lambda^{iv}}$  may be reduced to the same form (ae.). Consequently its second and third terms will amalgamate, respectively, with the first and second terms of the right-hand side of (a.) (thus becoming unavailable), and its only effective part is

$$p_0 + p_{\lambda^v} x^{\lambda^v} + \&c. \dots \dots \dots \text{(af.)}$$

If, therefore, the number of terms in (af.) be  $< 2$ , we shall have (*sup.* p. 132),

$$v'' = a''_2 p_{\lambda''}, \quad v' = a'_1 p_{\lambda'} + a''_1 p_{\lambda''}. \dots \dots \dots \text{(ag.)}$$

16. In general, then, the transformation (b.) of that page can be effected for equations of the **FOURTH** degree without the necessity of fulfilling (ag.); but in *critical* § cases we are limited to the **FIFTH** and higher degrees, since  $p_0$  disappears. On this account biquadratics cannot be reduced to a binomial form, as we might otherwise have inferred ||, for in such case we have, ultimately, to satisfy two homogeneous equations between two quantities of the form  $\Lambda + p_\lambda$ .

17. So, beyond all doubt, the transformation (o.) of p. 190 can, in general, be effected for equations of the **SIXTH** degree, without satisfying (ag.) by means of *one* cubic, two quadratics, and five base equations. But in critical cases we meet with the same obstacle as that mentioned in the last paragraph, and are limited to the **SEVENTH** and higher degrees; so that the solutions of equations of the fifth and sixth degrees present distinct difficulties ¶. If they are absolutely insoluble, may we not hope, from a consideration of the modes in which they evade different proposed methods of solution, to arrive at a more elementary demonstration of the fact than has yet appeared?

*On the Reduction of certain Functions.*

In those cases, in which the length of the calculations is

\* Communicated by T. S. Davies, Esq., F.R.S. and F.S.A.

† *Sup.* p. 191, Note \*. ‡ *Sup.* p. 132. § *Sup.* p. 191, par. 12.

|| *Sup.* p. 133, par. 5.

¶ See Sir W. R. Hamilton's "Inquiry" (cited *sup.* p. 191, Note \*), p. 298, line 25, and p. 317 [9.].

not such as to render the following reductions of merely theoretical interest, we may develop the formulæ for  $\tau$ ,  $\gamma^*$ , and such other symbols as may be requisite, so as to render the operations uniform and comparatively easy. The reductions I allude to, all of which are in theory possible †, are those of

$$f^3(3 \cdot 2^m - 2) \text{ to } h_1^3 + h_2^3 + \dots + h_m^3$$

$$f^4(u_m) \text{ to } h_1^4 + h_2^4 + \dots + h_m^4,$$

$$f^4(v_m) \text{ to } h_1^2 + h_2^2 + \dots + h_m^2,$$

where  $h_r$  is, in general, a linear though not homogeneous function of  $m - r + 1$  undetermined quantities,

$$u_{x+1} = 3 \cdot 2^{2u_x+1} - 2; \quad v_{x+1} = 3 \cdot 2^{2v_x} - 1; \quad u_0 = v_0 = 1;$$

and  $f^a(b)$  denotes the general function of the  $a$ th degree and  $b$ th order ‡, according to the notation which I used at p. 126 of the last volume.

Devereux Court, March 31, 1846.

LXVI. *Experimental Researches in Electricity*.—*Twentieth Series*. By MICHAEL FARADAY, Esq., D.C.L., F.R.S., Fullerian Prof. Chem. Royal Institution, Foreign Associate of the Acad. Sciences, Paris, Cor. Memb. Royal and Imp. Acadd. of Sciences, Petersburgh, Florence, Copenhagen, Berlin, Göttingen, Modena, Stockholm, &c. &c. §

§ 27. *On new magnetic actions, and on the magnetic condition of all matter* ||.

¶ i. *Apparatus required*. ¶ ii. *Action of magnets on heavy glass*. ¶ iii. *Action of magnets on other substances acting magnetically on light*. ¶ iv. *Action of magnets on the metals generally*.

2243. **T**HE contents of the last series of these researches were, I think, sufficient to justify the statement, that a new magnetic condition (*i. e.* one new to our know-

\* *Sup.* p. 190. See also one of my previous papers in the *Phil. Mag.* S. 3. vol. xxvii. pp. 292, 293.

† See *Mathematician*, vol. ii. p. 97. Ex. xlix, for a discussion of the first reduction.

‡ The order being the number of undetermined quantities, the degree the dimensions to which they enter. Might not the term 'simple' be advantageously applied to all equations of the first order?

§ From the *Philosophical Transactions* for 1846, Part I., having been read December 18, 1845.

|| My friend Mr. Wheatstone has this day called my attention to a paper by M. Becquerel, "On the magnetic actions excited in all bodies by the influence of very energetic magnets," read to the Academy of Sciences on the 27th of September 1827, and published in the *Annales de Chimie*, xxxvi. p. 337. It relates to the action of the magnet on a magnetic needle, on

ledge) had been impressed on matter by subjecting it to the action of magnetic and electric forces (2227.); which new condition was made manifest by the powers of action which the matter had acquired over light. The phænomena now to be described are altogether different in their nature; and they prove, not only a magnetic condition of the substances referred to unknown to us before, but also of many others, including a vast number of opaque and metallic bodies, and perhaps all except the magnetic metals and their compounds: and they also, through that condition, present us with the means of undertaking the correlation of magnetic phænomena, and perhaps the construction of a theory of general magnetic action founded on simple fundamental principles.

2244. The whole matter is so new, and the phænomena so varied and general, that I must, with every desire to be brief, describe much which at last will be found to concentrate under simple principles of action. Still, in the present state of our knowledge, such is the only method by which I can make these principles and their results sufficiently manifest.

#### ¶ i. Apparatus required.

2245. The effects to be described require magnetic apparatus of great power, and under perfect command. Both these points are obtained by the use of electro-magnets, which can be raised to a degree of force far beyond that of natural or

soft iron, on the deutoxide and tritoxide of iron, on the tritoxide alone, and on a needle of wood. The author observed, and quotes Coulomb as having also observed, that a needle of wood, under certain conditions, pointed *across* the magnetic curves; and he also states the striking fact that he had found a needle of wood place itself parallel to the wires of a galvanometer. These effects, however, he refers to a degree of magnetism less than that of the tritoxide of iron, but the same in character, for the bodies take the same position. The polarity of steel and iron is stated to be in the direction of the length of the substance, but that of tritoxide of iron, wood and gum-lac, most frequently in the direction of the width, and always when one magnetic pole is employed. "This difference of effect, which establishes a line of demarcation between these two species of phænomena, is due to this, that the magnetism being very feeble in the tritoxide of iron, wood, &c., we may neglect the reaction of the body on itself, and therefore the direct action of the bar ought to overrule it."

As the paper does not refer the phænomena of wood and gum-lac to an elementary *repulsive* action, nor show that they are common to an immense class of bodies, nor distinguish this class, which I have called diamagnetic, from the magnetic class; and, as it makes all magnetic action of one kind, whereas I show that there are two kinds of such action, as distinct from each other as positive and negative electric action are in their way, so I do not think I need alter a word or the date of that which I have written; but am most glad here to acknowledge M. Becquerel's important facts and labours in reference to this subject.—M. F. Dec. 5, 1845.

steel magnets; and further, can be suddenly altogether deprived of power, or made energetic to the highest degree, without the slightest alteration of the arrangement, or of any other circumstance belonging to an experiment.

2246. One of the electro-magnets which I use is that already described under the term Woolwich helix (2192.). The soft iron core belonging to it is twenty-eight inches in length and 2.5 inches in diameter. When thrown into action by ten pair of Grove's plates, either end will sustain one or two half-hundred weights hanging to it. The magnet can be placed either in the vertical or the horizontal position. The iron core is a cylinder with flat ends, but I have had a cone of iron made, two inches in diameter at the base and one inch in height, and this placed at the end of the core, forms a conical termination to it, when required.

2247. Another magnet which I have had made has the horse-shoe form. The bar of iron is forty-six inches in length and 3.75 inches in diameter, and is so bent that the extremities forming the poles are six inches from each other; 522 feet of copper wire 0.17 of an inch in diameter and covered with tape, are wound round the two straight parts of the bar, forming two coils on these parts, each sixteen inches in length, and composed of three layers of wire: the poles are, of course, six inches apart, the ends are planed true, and against these move two short bars of soft iron, 7 inches long and  $2\frac{1}{2}$  by 1 inch thick, which can be adjusted by screws, and held at any distance less than six inches from each other. The ends of these bars form the opposite poles of contrary name; the magnetic field between them can be made of greater or smaller extent, and the intensity of the lines of magnetic force be proportionately varied.

2248. For the suspension of substances between and near the poles of these magnets, I occasionally used a glass jar, with a plate and sliding wire at the top. Six or eight lengths of cocoon silk being equally stretched, were made into one thread and attached, at the upper end, to the sliding rod, and at the lower end to a stirrup of paper, in which anything to be experimented on could be sustained.

2249. Another very useful mode of suspension was to attach one end of a fine thread, six feet long, to an adjustable arm near the ceiling of the room, and terminating at the lower end by a little ring of copper wire; any substance to be suspended could be held in a simple cradle of fine copper wire having eight or ten inches of the wire prolonged upward; this, being bent into a hook at the superior extremity, gave the means of attachment to the ring. The height of the sus-


pended substance could be varied at pleasure, by bending any part of the wire at the instant into the hook form. A glass cylinder placed between the magnetic poles was quite sufficient to keep the suspended substance free from any motion, due to the agitation of the air.

2250. It is necessary, before entering upon an experimental investigation with such an apparatus, to be aware of the effect of any magnetism which the bodies used may possess; the power of the apparatus to make manifest such magnetism is so great, that it is difficult on that account to find writing-paper fit for the stirrup above-mentioned. Before therefore any experiments are instituted, it must be ascertained that the suspending apparatus employed does not point, *i. e.* does not take up a position parallel to the line joining the magnetic poles, by virtue of the magnetic force. When copper suspensions are employed, a peculiar effect is produced (2309.), but when understood, as it will be hereafter, it does not interfere with the results of experiment. The wire should be fine, not magnetic as iron, and the form of the suspending cradle should not be elongated horizontally, but be round or square as to its general dimensions, in that direction.

2251. The substances to be experimented with should be carefully examined, and rejected if not found free from magnetism. Their state is easily ascertained; for, if magnetic, they will either be attracted to the one or the other pole of the great magnet, or else point between them. No examination by smaller magnets, or by a magnetic needle, is sufficient for this purpose.

2252. I shall have such frequent occasion to refer to two chief directions of position across the magnetic field, that to avoid periphrasis, I will here ask leave to use a term or two conditionally.

One of these directions is that from pole to pole, or along the line of magnetic force; I will call it the axial direction: the other is the direction perpendicular to this, and across the line of magnetic force; and for the time, and as respects the space between the poles, I will call it the equatorial direction. Other terms that I may use, I hope will explain themselves.



#### ¶ ii. *Action of magnets on heavy glass.*

2253. The bar of silicated borate of lead, or heavy glass already described as the substance in which magnetic forces were first made effectually to bear on a ray of light (2152:),

and which is two inches long, and about 0.5 of an inch wide and thick, was suspended centrally between the magnetic poles (2247.), and left until the effect of torsion was over. The magnet was then thrown into action by making contact at the voltaic battery: immediately the bar moved, turning round its point of suspension, into a position across the magnetic curve or line of force, and after a few vibrations took up its place of rest there. On being displaced by hand from this position, it returned to it, and this occurred many times in succession.

2254. Either end of the bar indifferently went to either side of the axial line. The determining circumstance was simply inclination of the bar one way or the other to the axial line, at the beginning of the experiment. If a particular or marked end of the bar were on one side of the magnetic, or axial line, when the magnet was rendered active, that end went further outwards, until the bar had taken up the equatorial position.

2255. Neither did any change in the magnetism of the poles, by change in the direction of the electric current, cause any difference in this respect. The bar went by the shortest course to the equatorial position.

2256. The power which urged the bar into this position was so thoroughly under command, that if the bar were swinging it could easily be hastened in its course into this position, or arrested as it was passing from it by seasonable contacts at the voltaic battery.

2257. There are two positions of equilibrium for the bar; one stable, the other unstable. When in the direction of the axis or magnetic line of force, the completion of the electric communication causes no change of place; but if it be the least oblique to this position, then the obliquity increases until the bar arrives at the equatorial position; or if the bar be originally in the equatorial position, then the magnetism causes no further changes, but retains it there (2298. 2299. 2384.).

2258. Here then we have a magnetic bar which points east and west, in relation to north and south poles, *i. e.* points perpendicularly to the lines of magnetic force.

2259. If the bar be adjusted so that its point of suspension, being in the axial line, is not equidistant from the poles, but near to one of them, then the magnetism again makes the bar take up a position perpendicular to the magnetic lines of force; either end of the bar being on the one side of the axial line, or the other, at pleasure. But at the same time there is another effect, for at the moment of completing the electric contact, the centre of gravity of the bar recedes from the pole and remains repelled from it as long as the magnet is retained

excited. On allowing the magnetism to pass away, the bar returns to the place due to it by its gravity.

2260. Precisely the same effect takes place at the other pole of the magnet. Either of them is able to repel the bar, whatever its position may be, and at the same time the bar is made to assume a position, at right angles, to the line of magnetic force.

2261. If the bar be equidistant from the two poles, and in the axial line, then no repulsive effect is or can be observed.

2 62. But preserving the point of suspension in the equatorial line, *i. e.* equidistant from the two poles, and removing it a little on one side or the other of the axial line (2252.), then another effect is brought forth. The bar points as before across the magnetic line of force, but at the same time it recedes from the axial line, increasing its distance from it, and this new position is retained as long as the magnetism continues, and is quitted with its cessation.

2263. Instead of two magnetic poles, a single pole may be used, and that either in a vertical or a horizontal position. The effects are in perfect accordance with those described above; for the bar, when near the pole, is repelled from it in the direction of the line of magnetic force, and at the same time it moves into a position perpendicular to the direction of the magnetic lines passing through it. When the magnet is vertical (2246.) and the bar by its side, this action makes the bar a tangent to the curve of its surface.

2264. To produce these effects, of pointing across the magnetic curves, the form of the heavy glass must be long; a cube or a fragment approaching roundness in form, will not point, but a long piece will. Two or three rounded pieces or cubes, placed side by side in a paper tray, so as to form an oblong accumulation, will also point.

2265. Portions, however, of any form, are *repelled*; so if two pieces be hung up at once in the axial line, one near each pole, they are repelled by their respective poles, and approach, seeming to attract each other. Or if two pieces be hung up in the equatorial line, one on each side of the axis, then they both recede from the axis, seeming to repel each other.

2266. From the little that has been said, it is evident that the bar presents in its motion a complicated result of the force exerted by the magnetic power over the heavy glass, and that when cubes or spheres are employed, a much simpler indication of the effect may be obtained. Accordingly, when a cube was thus used with the two poles, the effect was repulsion or recession from either pole, and also recession from the magnetic axis on either side.

2267. So the indicating particle would move, either along the magnetic curves, or across them; and it would do this either in one direction or the other; the only constant point being, that its tendency was to move from stronger to weaker places of magnetic force.

2268. This appeared much more simply in the case of a single magnetic pole, for then the tendency of the indicating cube or sphere was to move outwards, in the direction of the magnetic lines of force. The appearance was remarkably like a case of weak electric repulsion.

2269. The cause of the pointing of the bar, or any oblong arrangement of the heavy glass, is now evident. It is merely a result of the tendency of the particles to move outwards, or into the positions of weakest magnetic action. The joint exertion of the action of all the particles brings the mass into the position, which, by experiment, is found to belong to it.

2270. When one or two magnetic poles are active at once, the courses described by particles of heavy glass free to move, form a set of lines or curves, which I may have occasion hereafter to refer to; and as I have called air, glass, water, &c. diamagnetics (2149.), so I will distinguish these lines by the term *diamagnetic curves*, both in relation to, and contradiction from, the lines called magnetic curves.

2271. When the bar of heavy glass is immersed in water, alcohol, or æther, contained in a vessel between the poles, all the preceding effects occur; the bar points and the cube recedes exactly in the same manner as in air.

2272. The effects equally occur in vessels of wood, stone, earth, copper, lead, silver, or any of those substances which belong to the diamagnetic class (2149.).

2273. I have obtained the same equatorial direction and motions of the heavy glass bar as those just described, but in a very feeble degree, by the use of a good common steel horse-shoe magnet (2157.). I have not obtained them by the use of the helices (2191. 2192.) without the iron cores.

2274. Here therefore we have magnetic repulsion without polarity, *i. e.* without reference to a particular pole of the magnet, for either pole will repel the substance, and both poles will repel it at once (2262.). The heavy glass, though subject to magnetic action, cannot be considered as magnetic, in the usual acceptance of that term, or as iron, nickel, cobalt, and their compounds. It presents to us, under these circumstances, a magnetic property new to our knowledge; and though the phænomena are very different in their nature and character to those presented by the action of the heavy glass on light (2152.), still they appear to be dependent on, or con-



nected with, the same condition of the glass as made it then effective, and therefore, with those phænomena, prove the reality of this new condition.

¶ iii. *Action of magnets on other substances acting magnetically on light.*


2275. We may now pass from heavy glass to the examination of the other substances, which, when under the power of magnetic or electric forces, are able to affect and rotate a polarized ray (2173.), and may also easily extend the investigation to bodies which, from their irregularity of form, imperfect transparency, or actual opacity, could not be examined by a polarized ray, for here we have no difficulty in the application of the test to all such substances.

2276. The property of being thus repelled and affected by magnetic poles, was soon found not to be peculiar to heavy glass. Borate of lead, flint-glass, and crown-glass set in the same manner equatorially, and were repelled when near to the poles, though not to the same degree as the heavy glass.

2277. Amongst substances which could not be subjected to the examination by light, phosphorus in the form of a cylinder presented the phænomena very well; I think as powerfully as heavy glass, if not more so. A cylinder of sulphur, and a long piece of thick India rubber, neither being magnetic after the ordinary fashion, were well-directed and repelled.

2278. Crystalline bodies were equally obedient, whether taken from the single or double refracting class (2237.). Prisms of quartz, calcareous spar, nitre and sulphate of soda, all pointed well, and were repelled.

2279. I then proceeded to subject a great number of bodies, taken from every class, to the magnetic forces, and will, to illustrate the variety in the nature of the substances, give a comparatively short list of crystalline, amorphous, liquid and organic bodies below. When the bodies were fluids, I inclosed them in thin glass tubes. Flint-glass points equatorially, but if the tube be of very thin glass, this effect is found to be small when the tube is experimented with alone; afterwards, when it is filled with liquid and examined, the effect is such that there is no fear of mistaking that due to the glass for that of the fluid. The tubes must not be closed with cork, sealing-wax, or any ordinary substance taken at random, for these are generally magnetic (2285.). I have usually



so shaped them in the making, and drawn them off at the neck, as to leave the aperture on one side, so that when filled with liquid they required no closing.

2280. Rock crystal.	Glass.
Sulphate of lime.	Litharge.
Sulphate of baryta.	White arsenic.
Sulphate of soda.	Iodine.
Sulphate of potassa.	Phosphorus.
Sulphate of magnesia.	Sulphur.
Alum.	Resin.
Muriate of ammonia.	Spermaceti.
Chloride of lead.	Caffeine.
Chloride of sodium.	Cinchonia.
Nitrate of potassa.	Margaric acid.
Nitrate of lead.	Wax from shell-lac.
Carbonate of soda.	Sealing-wax.
Iceland spar.	Olive-oil.
Acetate of lead.	Oil of turpentine.
Tartrate of potash and antimony.	Jet.
Tartrate of potash and soda.	Caoutchouc.
Tartaric acid.	Sugar.
Citric acid.	Starch.
Water.	Gum-arabic.
Alcohol.	Wood.
Æther.	Ivory.
Nitric acid.	Mutton, dried.
Sulphuric acid.	Beef, fresh.
Muriatic acid.	Beef, dried.
Solutions of various alkaline and earthy salts.	Blood, fresh.
	Blood, dried.
	Leather.
	Apple.
	Bread.

2281. It is curious to see such a list as this of bodies presenting on a sudden this remarkable property, and it is strange to find a piece of wood, or beef, or apple, obedient to or repelled by a magnet. If a man could be suspended, with sufficient delicacy, after the manner of Dufay, and placed in the magnetic field, he would point equatorially; for all the substances of which he is formed, including the blood, possess this property.

2282. The setting equatorially depends upon the form of the body, and the diversity of form presented by the different substances in the list was very great; still the general result, that elongation in one direction was sufficient to make them take up an equatorial position, was established. It was not

difficult to perceive that comparatively large masses would point as readily as small ones, because in larger masses more lines of magnetic force would bear in their action on the body, and this was proved to be the case. Neither was it long before it evidently appeared that the form of a plate or a ring was quite as good as that of a cylinder or a prism; and in practice it was found that plates and flat rings of wood, spermaceti, sulphur, &c., if suspended in the right direction, took up the equatorial position very well. If a plate or ring of heavy glass could be floated in water, so as to be free to move in every direction, and were in that condition subject to magnetic forces diminishing in intensity, it would immediately set itself equatorially, and if its centre coincided with the axis of magnetic power, would remain there; but if its centre were out of this line, it would then, perhaps, gradually pass off from this axis in the plane of the equator, and go out from between the poles.

2283. I do not find that division of the substance has any distinct influence on the effects. A piece of Iceland spar was observed, as to the degree of force with which it set equatorially; it was then broken into six or eight fragments, put into a glass tube and tried again; as well as I could ascertain, the effect was the same. By a second operation, the calcareous spar was reduced into coarse particles; afterwards to a coarse powder, and ultimately to a fine powder: being examined as to the equatorial set each time, I could perceive no difference in the effect, until the very last, when I thought there might be a slight diminution of the tendency, but if so, it was almost insensible. I made the same experiment on silica with the same result, of no diminution of power. In reference to this point I may observe, that starch and other bodies in fine powder exhibited the effect very well.

2284. It would require very nice experiments and great care to ascertain the specific degree of this power of magnetic action possessed by different bodies, and I have made very little progress in that part of the subject. Heavy glass stands above flint-glass, and the latter above plate-glass. Water is beneath all these, and I think alcohol is below water, and æther below alcohol. The borate of lead is I think as high as heavy glass, if not above it, and phosphorus is probably at the head of all the substances just named. I verified the equatorial set of phosphorus between the poles of a common magnet (2273.).

2285. I was much impressed by the fact that blood was not magnetic (2280.), nor any of the specimens tried of red muscular fibre of beef or mutton. This was the more striking,

because, as will be seen hereafter, iron is *always* and in almost *all states* magnetic. But in respect to this point it may be observed, that the ordinary magnetic property of matter and this *new property* are in their effects opposed to each other; and that when this property is strong it may overcome a very slight degree of ordinary magnetic force, just as also a certain amount of the magnetic property may oppose and effectually hide the presence of this force (2422.). It is this circumstance which makes it so necessary to be careful in examining the magnetic condition of the bodies in the first instance (2250.). The following list of a few substances, which were found slightly magnetic, will illustrate this point:—Paper, sealing-wax, china ink, Berlin porcelain, silkworm-gut, asbestos, fluor-spar, red lead, vermilion, peroxide of lead, sulphate of zinc, tourmaline, plumbago, shell-lac, charcoal. In some of these cases the magnetism was generally diffused through the body, in other cases it was limited to a particular part.

2286. Having arrived at this point, I may observe, that we can now have no difficulty in admitting that the phænomena abundantly establish the existence of a magnetic property in matter new to our knowledge. Not the least interesting of the consequences that flow from it, is the manner in which it disposes of the assertion which has sometimes been made, that all bodies are magnetic. Those who hold this view, mean that all bodies are magnetic as iron is, and say that they point between the poles. The new facts give not a mere negative to this statement, but something beyond, namely, an affirmative as to the existence of forces in all ordinary bodies, directly the opposite of those existing in magnetic bodies, for whereas those practically produce attraction, these produce repulsion; those set a body in the axial direction, but these make it take up an equatorial position: and the facts with regard to bodies generally are exactly the reverse of those which the view quoted indicates.

[To be continued.]


## LXVII. *Description of a new Mercurial Trough.*

*By Professor LOUYET of Brussels\*.*

**I**N small laboratories in which one of the chief points to be aimed at is œconomy, in making researches on gases soluble in water, a small porcelain trough is commonly employed capable of containing twenty to twenty-five pounds of mercury. The size of the bell-glass is proportioned to the capacity of the trough; thus only small quantities of gas can be

\* Communicated through Prof. Grove, by the Author.

collected,—quantities often insufficient for the desired experiments. With a view to obviate this inconvenience, I have modified the mercurial trough, so as to be able, without sacrificing œconomy, to collect considerable quantities of gas; and I have thought it may be of use to give a description of this new instrument, for the service of persons engaged in particular researches. This apparatus is formed of a small oblong oak box, to the bottom of which is cemented a glass, which fits exactly over its whole extent. The external surface of this glass is carefully polished and prepared. In the centre of one of its small sides (it is rectangular), is worked a narrow and deep groove parallel to the large sides of the right angle. This opening corresponds to a hollow in the bottom of the box. This being done, I arrange the apparatus in the following way, when I desire to collect a gas over the mercury:—I procure bell-glasses made of emery-stoppered bottles, the bottom of which is removed; the edges of these receivers are prepared and rubbed with emery, and apply accurately to the ground glass, precisely like the receiver of an air-pump. The edges may be very slightly greased, or this precaution may be dispensed with. The receiver is placed on the ground glass, where it is kept fixed with one hand; with the other hand the stopper is removed, and it is entirely filled with mercury; then it is carefully re-stoppered. This being done, a small quantity of mercury is poured into the box, so as to fill the small cavity, and to cover its bottom with a stratum of some millimetres. The receiver may now be moved in all directions, and may be slid until it is over the small cavity, into which the extremity of the curved tube by which the gas is disengaged, is adapted. To one of the angles of the box may be fitted a small pure iron stopcock, by which the mercury is drawn off when the operation is ended. For greater clearness, I subjoin a figure, which represents this new trough of the dimensions which I have adopted.



$AB = 4\frac{1}{2}$  centimetres\*.  $BC = 23$  centimetres.  $CD = 17$  centimetres.  
Depth of the longitudinal groove, taken above the glass plate, = 2 centimetres.

\* The centimetre is = 0.393708 of an English inch.

LXVIII. *Proceedings of Learned Societies.*

## ROYAL SOCIETY.

*Anniversary Meeting, December 1, 1845.*

**T**HE Marquis of Northampton in the Chair.

The noble President stated that the two Royal Medals had been adjudged by the Council to the Astronomer Royal, for his Inquiries into the Tides on the Coast of Ireland; to Mr. Beck, for his Investigation of the Nerves of the Uterus\*; and the Copley Medal to Prof. Schwann, for his valuable work On the Analogies of Vegetables and Animals.

After presenting the Medals, the President proceeded to the biographical notices of some of the deceased members, from which we select the following:—

DR. WILLIAM HEBERDEN, the son of the eminent and accomplished author of the ‘Commentaries on the History and Cure of Diseases,’ was born in London in the year 1767. At the early age of seven, he was sent to school at the Charter House, and appears to

\* The report of the Committee of Physiology on the claims of Mr. Beck’s paper to the award of the Medal, is as follows:—

“The paper of Mr. Beck contains the result of an elaborate anatomical investigation of the Nerves of the Uterus, together with observations on the structure and connexions of the sympathetic nerve.

“By his researches the author has cleared up various points concerning the nerves of the uterus which have hitherto been doubtful or misunderstood. He has determined more precisely than heretofore the source and mode of distribution of these nerves, and the real extent to which the organ is supplied with them. The true nature of the nervous ganglia at the neck of the uterus, and of the plexuses formed by the sympathetic and sacral nerves in the same situation, is also satisfactorily made out, as well as the fact that the branches derived from the sacral nerves are not destined for the uterus, but are distributed to adjacent organs.

“With regard to the sympathetic nerve, it is shown that there are both grey and white separate branches of communication between that nerve and the spinal nerves. This important fact has, it is true, been already pointed out in the recently published work of Tedd and Bowman, but the author of the paper has nevertheless the merit of arriving at it independently, by his own observations. He has further shown that the white and grey constituents of the nerve keep distinct from each other, not only in the so-called trunk of the sympathetic, but also in its primary branches, as far as the visceral ganglia, beyond which the white and grey parts become intermixed in the nerves distributed to the viscera. The precise mode of connexion of the white and grey communicating branches with the spinal nerves is also carefully investigated. These observations appear important as tending to throw light on the constitution of the sympathetic nerve and its relation to the rest of the nervous system.

“The Committee consider the paper of Mr. Beck as a most valuable contribution to the Anatomy of the nervous system, and as affording additional and more precise data for physiological reasoning respecting the nerves to which it refers. On these grounds, as well as on account of the consummate skill and devoted perseverance displayed by the author in his arduous investigations, they have recommended that his paper be rewarded with the Royal Medal.”

have there made rapid progress in the elementary branches of education. His academical studies were pursued in St. John's College, Cambridge, where he highly distinguished himself both by his mathematical and his classical acquirements. Under his father's tuition he applied himself with great diligence to his profession as a pupil of St. George's Hospital, to which, at a subsequent period, he was appointed one of the Physicians. He was elected a Fellow of this Society in the year 1791: and in 1797, became a Fellow of the College of Physicians. He died on the 19th of February, 1845, aged 77.

His contributions to the Philosophical Transactions consist of two papers; the first in 1796, on the influence of cold on the health of the inhabitants of London; in which he shows, in opposition to the popular prejudices then prevalent, that a severe winter is attended with greatly increased mortality. The second paper is entitled "On the heat of July 1825, together with some remarks on sensible cold," in which he points out the causes which influence our sensations of temperature, and more especially the powerful effect of wind in increasing the rate of cooling, and consequently of creating the sensation of cold in the human body, independently of any actual depression in the temperature of the air.

JOHN FREDERIC DANIELL was born in Essex Street, Strand, 12th of March, 1790. His father, George Daniell, Esq., Bencher of the Inner Temple, provided him with a good classical education under his own roof. At an early age he showed fondness for the pursuits of science, and was placed in the sugar refining establishment of a relative, where he introduced important improvements in the manufacture. The pursuits of business, however, were uncongenial to his tastes, and he soon relinquished this occupation. In 1813 he was elected a Fellow of the Royal Society, of which body he continued till the day of his death a zealous and active member.

The services he rendered to more than one branch of science were of no ordinary description. From an early period of his life his mind was directed to the study of meteorology, at a time when it consisted of little more than a vast accumulation of facts and observations.

In the year 1823 he published the first edition of his 'Meteorological Essays,' which constituted a new epoch in the science, and still continues the standard work of reference, the third edition of which he had nearly completed at the time of his death. This was the first attempt to embrace in a general view the scattered facts of the science, and by synthetically applying the known laws which regulate the constitution of gases and vapours, the principles of their equilibrium, and the distribution of heat among them, to give a connected account of the main phenomena of the earth's atmosphere. He insisted on the paramount importance of extreme accuracy in the construction of the instruments employed for such inquiries, and gave directions by which the needful accuracy could with certainty and facility be obtained. By the invention of the hygrometer, which bears his name, he first conferred precision on the means of ascer-

taining the moisture or dryness of the atmosphere, a point of cardinal importance in all investigations of this nature; his instrument still continues that which can be best depended upon for this purpose. With these accurate instruments, he for three years kept a faithful register of the various atmospheric changes; he organized the plan adopted by the Horticultural Society in their annual meteorological reports, a plan which formed the model to the admirable and more extended series of meteorological observations now issued weekly from the Greenwich Observatory under the superintendence of the Astronomer Royal.

In the year 1824 he communicated to the Horticultural Society an essay 'On Artificial Climate,' which appeared in their Transactions for that year. In this paper among other subjects he insisted on the absolute necessity of attention to the moisture of the atmosphere, as well as of that of maintaining in our hot-houses the moisture as well as the temperature of a tropical climate, if we would produce a vegetation of tropical luxuriance. The publication of this essay caused a complete change in the methods adopted for the culture of plants in general, and particularly of those contained in green-houses and hot-houses, which upon the new plans speedily outgrew the houses provided for their reception. The Society immediately awarded him their silver medal to mark their sense of the importance of his views, and now after an experience of more than twenty years, Dr. Lindley, Professor of Botany in University College, not a fortnight before his death, in an article in the *Gardener's Chronicle*, tracing the origin of the improvements in this branch of horticulture, ascribes the rapid advance in the practice of the art, mainly to the sound and original views promulgated in this essay.

For the purpose of making more minute and accurate observations upon variations in the atmospheric pressure, Mr. Daniell proposed to the Royal Society, in 1830, to construct a barometer in which water should be the fluid used instead of mercury. He was in consequence requested to superintend the construction of such an instrument. Great practical difficulties attended the undertaking, but these he happily surmounted, and the instrument now stands in the Hall of the Apartments of the Royal Society; he was engaged in re-adjusting it within a few weeks of his decease. On occasion of the late Antarctic Expedition under the command of Captain Sir James Ross, and the establishment by Government of the Magnetic and Meteorological observations, founded a few years since in different parts of the British Empire, when the Admiralty applied to the Royal Society for instructions as to the nature and extent of the observations to be made, Mr. Daniell was requested by the Committee of Physics of the Royal Society, to draw up the Meteorological portion of these directions. The paper which he then prepared furnished the basis of that part of the Report of the Committee, published in the year 1840, under the sanction of the Royal Society.

But it was not alone to meteorology, and its practical applications, that his labours were confined; his researches upon various chemical subjects were not less numerous or important. More than forty



original papers, including thirteen on meteorology, were communicated by him to various scientific publications; among others he published several memoirs on Crystallization, and its attendant phenomena. Between the years 1830 and 1844, the Transactions of this Society were enriched by twelve papers on important subjects from his pen. He invented a process for making gas from resin for the purposes of illumination, by which the streets of New York are lighted at the present time. For this improvement he received no other acknowledgment than a vote of a few pounds' worth of books. In the year 1830, he described in the Philosophical Transactions, a new instrument for measuring high degrees of heat, such as the temperature of furnaces, and the melting-points of metals. By means of this, his pyrometer, he ascertained numerous facts of great interest both in a scientific and in a practical point of view. For the invention of this instrument, which is still the best for the objects intended, the Royal Society awarded him the Rumford Medal.

After his appointment as Professor of Chemistry in King's College, his researches were turned principally to the phenomena presented by Voltaic Electricity, and they led to the invention of his constant battery; for this the Royal Society conferred upon him the highest honour in their gift, the Copley Medal for the year 1836. The possibility of maintaining powerful and equable currents of electricity for any required period, was established by this invention. The impulse thus given to the progress of electrical research cannot be too highly estimated, and to it must be traced the numerous applications of electricity, to the blasting of rocks, the working of mines, and to submarine operations, and to the arts of electro-plating, gilding, zincing, &c., which have recently acquired such magnitude. His subsequent researches in the same field are contained in the Philosophical Transactions, and were honoured by the Society in the year 1842 by one of the Royal Medals. In 1839 Professor Daniell was placed on the Commission appointed by the Admiralty to inquire into the best method of defending the ships in the Royal Navy from lightning, and the same year the Royal Society honoured him with the office of Foreign Secretary to their body. His "Introduction to Chemical Philosophy," published during the course of this year, contributed still further to increase his reputation, and in 1842 he received from the University of Oxford the honorary degree of D.C.L. In consequence of the rapid corrosion of the copper sheathing of the vessels employed upon the African stations, the Admiralty requested him to examine the damaged sheets of metal and the waters taken up from the localities where the corrosion was the greatest; he detected the cause of this decay, showing that sulphuretted hydrogen was abundantly generated in the ocean at these spots, and succeeded in extracting from the metal plates the sulphur which had occasioned their corrosion. It is a remarkable proof of the variety and extent of Mr. Daniell's acquirements, that he received at different times all the medals in the gift of the Royal Society.

The circumstances which attended the sudden and lamented termination of his valuable life, are known to most of the Fellows of

this Society. On the 13th of March 1845, after delivering his usual lecture at King's College, apparently in perfect health, he attended the Council Meeting of this Society, and shortly after making some observation upon the business of the meeting, was seized with symptoms indicating an attack of apoplexy. Several medical men who were present hastened to his relief, and he was immediately bled; not the slightest benefit, however, attended this measure, and in five minutes he was a corpse. The shock occasioned by this melancholy event, may be easier imagined than described, and cast a widespread gloom over the extensive circle of his friends and acquaintance. As a mark of respect to his memory, the Noble President postponed the ordinary meeting of the Society, which was to have been held that evening. His remains were interred at Norwood, Surrey, where during the last ten years of his life he had resided.

Mr. Daniell survived his wife eleven years, and left a family of two sons and five daughters to deplore his loss. High as were his scientific attainments, he possessed others of a still loftier and more enduring character; to the sterling qualities of a vigorous understanding, and a kind and benevolent heart, he united the humble and unobtrusive piety of a sincere christian.

Mr. GEORGE BASSEVI, an architect of distinguished reputation.

JAQUES DOMINIQUE CASSINI, Comte du Thury, was elected a Foreign Member of this Society in 1789, and at the time of his death had attained the extraordinary age of 97 years: he was the fourth in direct descent of a family which, during a period of nearly two centuries, has been singularly illustrious in the history of the sciences, and more particularly of astronomy. His great grandfather, Jaques Dominique Cassini, one of the greatest astronomers of his age, was born in 1625, and was invited by Louis XIV. from Italy to France, to preside over the magnificent Observatory of Paris, which was built under his directions: his first successor in the direction of this establishment was his son Jean Jaques, an astronomer not less eminent than himself; the second his grandson, more commonly designated as Cassini de Thury, so well known by his great Geodetical Survey and Map of France; and the third his great grandson, the subject of the present notice, who was displaced from it by the troubles of the Revolution, which involved him, at least for a time, in the common proscription of the aristocracy of France. The shock of these sad events seems to have diverted his mind from scientific pursuits, for we find his name connected with no research in astronomy or geodesy during the last half-century.

The Comte de Cassini completed the celebrated map of France which had been begun by his father. He published an account of voyages which he made in 1768 and 1769, for the trial of the marine chronometers of Le Roy, and a memoir "Sur l'influence de l'équinoxe du printemps et du solstice d'été sur les déclinaisons et les variations de l'aiguille aimantée:" he superintended and published an account of the observations which were made in 1789, by a commission appointed for that purpose, for the junction of the Observatories of Paris and Greenwich, with a special reference to

the connection of the Geodetical Survey of France which had been made by his father, with the corresponding Survey of England which was at that time in progress under General Roy: he was the author likewise of "Mémoires pour servir à l'Histoire des Sciences et à celle de l'Observatoire Royal de Paris, suivis de la Vie de Jaques Dominique Cassini, premier du nom."

With him have terminated the honours of the house of Cassini, though he was not the last of his race who distinguished himself in the career of the sciences; his son, Henri Cassini, an upright and enlightened judge in the Cour Royale and the Cour de Cassation, and one of the most learned botanists of his age, fell a victim to the cholera in 1833, and with him died the last stay of the old age of his father: he was the fifth of his family who had been elected a member of the Académie des Sciences. He was a boy at the breaking out of the Revolution, and was compelled, from a regard to his personal safety, to live in the strictest retirement at his father's domain of Thury; a circumstance which turned his attention to the cultivation of Natural History and Botany, and diverted him, as he was accustomed to lament, from those studies and pursuits which formed, as it were, the proper and hereditary honours of his family.

THÉODORE DE SAUSSURE was born in Geneva the 14th of October, 1767. His father, known throughout the civilized world as the geological explorer of the Alps, who first reared observatories on heights almost inaccessible, and who inscribed his name on the eternal snows of the loftiest mountain in Europe, was by profession a physician. Being animated with an ardent love for science, which he cultivated most assiduously, it is not a matter of surprise that after entrusting his son for a short time to a private tutor, he should have undertaken personally his education, so far as to enable him to enter the Academy of Geneva, where young De Saussure soon distinguished himself. Previously to this period, his father had caused him to study medicine, mineralogy, and natural history; and had also inspired him with a taste for experimental chemistry, which he constantly required for the analyses of minerals. By degrees the son became associated with the scientific labours of his father, who records that when he resolved upon attempting the ascent of Mont Blanc, in August 1787, his son, then nineteen years of age, expressed the strongest desire to accompany him; but being apprehensive that he was not sufficiently strong, he was unwillingly obliged to leave him at the Priory at Chamouni, where he made with great care meteorological observations, simultaneously with those carried on at the summit of Mont Blanc. In the month of June of the following year, Théodore de Saussure accompanied his father in the laborious and hazardous expedition to the Col du Géant, where they remained for seventeen days, during which time young De Saussure rose every morning at four o'clock, to commence the meteorological observations, which he continued with unremitting diligence until ten o'clock each night; and so thoroughly did he enter into the scientific pursuits of his parent, that he almost importuned the latter to extend the period of his sojourn amidst those splendid scenes;

which was however effectually prevented by the guides, who, alarmed at the idea of a longer stay amidst those icy heights, destroyed all the remaining provisions, necessarily compelling the De Saussures to descend to Cormayeur.

From this period we find Théodore De Saussure always accompanying his father, who was not slow in availing himself of the great advantages derivable from his son's labours. In 1789, they made the very difficult tour of Monte Rosa, it being the first time that the gold mines of Macugnaga were visited by men of science. It was during this excursion that young De Saussure restored by experiments that confidence in the accuracy of the barometer for measuring heights, which the assertion of Bouguer had tended to weaken. He made seventy experiments at different heights, and in calculating the results, was always careful to make the necessary corrections for temperature and humidity, which Bouguer appears to have neglected. The enthusiasm and physical energy of young De Saussure were almost too great for his father, who was now aged, and weakened by various illnesses, and consequently we often find the latter compelled to resist his son's desire to prolong their arduous excursions.

The storms of the Revolution, more powerful than those of the Alps, at length put an effectual stop to these useful scientific excursions, which had been continued for so many years. Théodore de Saussure, in common with many men of his age, was compelled to leave his country. He visited this country with Alexander Marcet, who many years afterwards became his colleague in the Academy at Geneva. After travelling over England and Scotland, he returned to Geneva, and resolved henceforth to devote his life to scientific pursuits. The taste which he had acquired for chemistry under his father's tuition, had been strengthened in England and in France, where that science was eagerly cultivated; and on his return to Geneva, he determined to select the vegetable kingdom for the field of his researches, and zealously applied himself to discover by experiments the influence of the atmosphere and of soils upon plants, and the various chemical changes which they undergo. With the exception of some few accessory labours, M. de Saussure spent a long life upon this branch of science; and it may be truly said that he has done more to advance the knowledge of vegetable physiology than any other person. It is worthy of remark that he laboured seven years silently, before publishing the results of his investigations. These were comprised in his work entitled "*Recherches Chimiques sur la Végétation*," which appeared in 1804. He subsequently published in the *Annales de Physique et de Chimie*, the results of his investigations into the action of the petals of flowers upon the atmosphere. The great importance which he attached to the nutritive power of carbonic acid upon plants, directed his attention to the proportion of this gas in the atmosphere. In 1816, he published, in the first volume of the *Bibliothèque Universelle*, some researches on this subject, which being greatly extended, formed afterwards a paper which was published in 1830 in the

Mémoires de la Société de Physique et d'Histoire Naturelle de Genève, under the title of "De l'action des huiles sur le gaz oxygène à la température atmosphérique." After examining the action of the green portions of plants, roots and flowers upon the atmosphere, M. de Saussure carried his investigation to the same parts of fruits. The result was a long paper published in 1821, in the Memoirs of the same Society, entitled "Influence des fruits verts sur l'air, avant leur maturité." He shows in this paper, that unripe fruit exercises the same influence as leaves upon the air.

Independently of his vegeto-physiological researches, M. de Saussure published some papers descriptive of minerals in the Journal de Physique. These are entitled, "Analyse du Sappare," "Sur une hydrophane imbibée de cire," "Analyse de la Dolomie," and "Sur le Sappare dur."

M. de Saussure was of a most reserved habit, the result probably of his solitary education: it is recorded of him that he seldom desired to converse with his friends on the scientific subjects occupying his attention; and so far did he carry this reserve, that even his most intimate acquaintances were generally ignorant of the nature of the papers which he proposed reading before the Society of Natural History. The same disposition prevented him from acting as Professor in the Academy of Geneva, though appointed to the Chair of Mineralogy and Geology in the year 1802. It was found impossible to overcome his repugnance to give the usual courses of lectures, though at the same time he gave evidence of his warm interest in the Academy by constantly attending its meetings. In 1814, he was elected a member of the Legislative Council of the Republic of Geneva, but he was too timid to take an active part in the debates of this body. In 1790 he became a member of the Agricultural Section of the Society of Arts, and always continued one of its most zealous supporters. He was a Foreign Member of the French Institute, of the Royal Academies of Naples, Turin and Munich; of the Institute of Fine Arts and Sciences at Amsterdam; of the Linnean Societies of Paris and London; the Wernerian Society of Edinburgh; and was elected a Foreign Member of the Royal Society in 1820. In 1842, M. de Saussure was unanimously elected President of the Scientific Congress, which met that year at Lyons, thus marking the high esteem in which he was held as a man of science. Having preserved throughout life the best health, M. de Saussure died on the 18th of April 1845, at the advanced age of 78, leaving behind him the reputation of a long life passed in severe and patient study, interrupted only when he came before the world with the results of his laborious experiments and researches.

Dr. Roget, reported the following Noblemen and Gentlemen as being duly elected Officers and Council for the ensuing year, viz.—

*President.*—The Marquis of Northampton. *Treasurer.*—George Rennie, Esq., V.P. *Secretaries.*—Peter Mark Roget, M.D., Samuel Hunter Christie, Esq., M.A. *Foreign Secretary.*—Lieut.-Col. Edward Sabine, R.A. *Other Members of the Council.*—John Bostock, M.D.; Sir William Burnett, M.D., K.C.H., V.P.; Charles Daubeny,

M.D.; Bryan Donkin, Esq.; Very Rev. Dean of Ely, V.P.; Thomas Galloway, Esq., M.A.; William Robert Grove, Esq., M.A.; Leonard Horner, Esq., V.P.; Sir J. W. Lubbock, Bart., M.A., V.P.; John Forbes Royle, M.D.; William Sharpey, M.D.; William Henry Smyth, Capt. R.N.; John Taylor, Esq.; Charles Wheatstone, Esq.; Rev. Robert Willis, M.A.; Lord Wrottesley, V.P.

LXIX. *Intelligence and Miscellaneous Articles.*

NOTE BY MR. T. HOPKINS ON HIS PAPER ON THE SEMI-DIURNAL FLUCTUATIONS OF THE BAROMETER.

*To Richard Taylor, Esq.,*

SIR,

IN looking over my paper, inserted in the Philosophical Magazine for March, "On the Causes of the Semi-diurnal Fluctuations of the Barometer," I discovered that a mistake had been made in putting down the figures in the Plymouth table, which express the temperature at 1 P.M. The mistake arose in this way. In the tables, as inserted in the Report of the British Association for 1839, the temperature for 1 P.M. is entered 45·83. This appeared a typographical error, and in order to correct it a 5 was substituted for the 4, leaving the numbers 55·83. This temperature of the dry thermometer was then compared with that of the wet-bulb thermometer, and the influences of temperature and evaporation as thus exhibited were inferred and remarks were made upon them in the paper. But an examination of other parts of the Plymouth report now shows that the figures 45 in the thermometric column should have been transposed, when the temperature at 1 P.M. would have been found only 54·83 instead of 55·83, making the rise of temperature from 10 A.M. to 1 P.M. only 1·99, whilst the force of evaporation is only 0·89. The tabular statement of the rise of temperature and the force of evaporation at the two periods would then be from

5 to 10	{	5·86 of temperature and	}	caused a rise,
		2·18 of evaporation		
10 to 1	{	1·99 of temperature and	}	caused a fall.
		0·89 of evaporation		

Wishing to be correct in my statements, I have to request that you will insert this letter in the next Number of your valuable publication.

I am, Sir,

Your most obedient Servant,

Manchester, March 6, 1846.

THOMAS HOPKINS.

ON SOME NEW COMPOUNDS OF PERCHLORIDE OF TIN.

BY M. LEWY.

The author remarks, that although the perchloride of tin has been the object of numerous researches, the compounds which it forms

with basic chlorides and organic matters have not hitherto attracted much attention.

*Compounds with Water.*—It is well known that when a small quantity of water is added to perchloride of tin, the whole becomes a crystalline mass; on adding more water, the hydrate thus formed dissolves, and yields fresh crystals by slow evaporation; the form of these could not be ascertained on account of their extreme deliquescence. Their formula appeared to be  $\text{SnCl}^2 + 5\text{HO}$ , or to consist of—

Chlorine	.....	40.55
Tin	.....	33.68
Water	.....	25.77
		<hr/>
		100.00

When these crystals are exposed *in vacuo* over sulphuric acid, they lose a certain quantity of water of crystallization, and eventually a hydrate remains containing only two equivalents of water.

*Compounds with the Chlorides.*—It is well known that perchloride of tin possesses properties analogous to those of acids; it combines with basic chlorides to form double chlorides, the greater part of which crystallize very readily; they all contain equal equivalents of perchloride of tin and basic chlorides.

The compounds formed with chloride of potassium and chloride of ammonium are anhydrous. The former contains—

	By analysis.	By calculation.	Equivalents.
Chlorine	.... 52.04	52.01	$\text{Cl}^3$
Tin	..... 28.50	28.79	Sn
Potassium	.. 18.76	19.19	K
	<hr/>	<hr/>	
	99.30	99.99	

*Chloride of Tin and Ammonium.*—This salt consists of—

	By analysis.	By calculation.	Equivalents.
Chlorine	.... 57.33	58.03	$\text{Cl}^3$
Nitrogen	.... 7.70	7.65	N
Hydrogen	.. 7.70	2.19	$\text{H}^4$
Tin	..... 32.30	32.13	Sn
		<hr/>	
		100.00	

The compounds which perchloride of tin forms with the chlorides of sodium, strontium, magnesium, calcium, and barium, all contain water of crystallization; and as far as the author's experiments have yet been carried, these double salts all contain five equivalents of water.

*Double chloride of Sodium and Tin*, when the analysis is corrected by calculation, appears to consist of—

		Equivalents.
Chlorine	..... 45.54	$\text{Cl}^3$
Sodium	..... 9.94	Na
Tin	..... 25.21	Sn
Water	..... 19.30	$\text{Aq}^5$
	<hr/>	
	99.99	

The form of this salt has not been determined, but it appeared to be formed of small prisms.

*Double chloride of Strontium and Tin.*—The corrected analysis gave—

		Equivalents.
Chlorine . . . . .	41·84	Cl <sup>3</sup>
Strontium . . . . .	17·26	Sr
Tin . . . . .	23·17	Sn
Water . . . . .	17·73	Aq <sup>5</sup>
	<hr/>	
	100·00	

This salt has the form of long channeled prisms, the summits of which are not of determinable form.

*Double chloride of Magnesium and Tin.*—The analysis corrected gave—

		Equivalents.
Chlorine . . . . .	47·71	Cl <sup>3</sup>
Magnesium . . . . .	5·66	Mg
Tin . . . . .	26·41	Sn
Water . . . . .	20·21	Aq <sup>5</sup>
	<hr/>	
	99·99	

It appears to crystallize in rhombohedrons of about 125°. This measure is however only an approximation to within one or two degrees; it was impossible to obtain a more accurate one, on account of the extreme deliquescence of the salt.

*Double chloride of Calcium and Tin* gave by corrected analysis,—

		Equivalents.
Chlorine . . . . .	46·17	Cl <sup>3</sup>
Calcium . . . . .	8·69	Ca
Tin . . . . .	25·56	Sn
Water . . . . .	19·56	Aq <sup>5</sup>
	<hr/>	
	99·98	

This salt is still more deliquescent than the preceding; the form of the crystal appears at first to be a cube, but on measuring the angles by the goniometer, one was found of 84° to 86°, and the other of 94° to 96°. It is therefore probable it also crystallizes in rhombohedrons.

*Double chloride of Barium and Tin.*—This gave by corrected analysis,—

		Equivalents.
Chlorine . . . . .	38·13	Cl <sup>3</sup>
Barium . . . . .	24·59	Ba
Tin . . . . .	21·11	Sn
Water . . . . .	16·16	Aq <sup>5</sup>
	<hr/>	
	99·99	

The form of this salt was not determined, but as far as an opinion could be formed, it appeared to consist of small prisms.

*Compounds of Bichloride of Tin and Organic Bodies.*—The author



prepared, as had been some years since been done by M. Kuhlman, compounds of bichloride of tin with sulphuric æther, alcohol, hydrochloric æther, and pyroxylic spirit; and M. Lewy combined it also with oxalic æther, benzoic æther, benzoate of methylene, acetic æther, acetic acid, benzoic acid, oil of bitter almonds, urea, camphor, ethal, &c. The greater part of these compounds formed very fine crystals, but their ready alteration in the air, and even *in vacuo*, as well as the difficulty of purifying them, have hitherto prevented an exact analysis of them; M. Lewy therefore endeavoured to verify the opinion of M. Kuhlman as to the composition of the compounds which he had formed.

*Compound of Perchloride of Tin and Sulphuric Æther.*—This forms crystals of very great beauty; it is obtained, as shown by M. Kuhlman, by mixing the two bodies either in the state of liquids or vapours. The crystals have the form of rhomboidal tables of a brilliant aspect and perfect formation. They are volatile without decomposition, dissolve readily in excess of æther, and decompose in contact with the air. This compound appeared to be formed of—

		Equivalents.
Chlorine . . . . .	34·77	Cl <sup>2</sup>
Tin . . . . .	28·88	Sn
Carbon . . . . .	23·57	C <sup>3</sup>
Hydrogen . . . . .	4·91	H <sup>10</sup>
Oxygen . . . . .	7·86	O <sup>2</sup>
	99·99	

*Compound of Perchloride of Tin and Anhydrous Alcohol.*—This was formed by merely mixing the two liquids. During mixture, the temperature of the substances was always kept below 32° Fahr.; when the combination has taken place, it is to be exposed *in vacuo* to sulphuric acid and potash. After some days the compound appears in the form of small prismatic crystals, which readily dissolve in an excess of alcohol, so that it is easy to re-crystallize them. The crystals must not, however, be kept too long *in vacuo*, as they then alter readily. This compound gave, by corrected analysis,—

		Equivalents.
Chlorine . . . . .	32·74	Cl <sup>3</sup>
Tin . . . . .	36·32	Sn <sup>2</sup>
Carbon . . . . .	14·82	C <sup>3</sup>
Hydrogen . . . . .	3·71	H <sup>12</sup>
Oxygen . . . . .	12·36	O <sup>3</sup>
	99·95	

*Compound of Perchloride of Tin with Oxalic Æther.*—This is prepared in the same manner as the preceding. When small quantities of perchloride of tin are added to oxalic æther, a moment arrives at which a crystalline mass is formed, consisting of small needles grouped round a common centre. These crystals alter readily, and it is best to analyse them as soon as formed. When mixed with water, oxalic æther is reproduced.

Analysis showed that this is a compound of equal equivalents of the perchloride and the æther, or—

		Equivalents.
Chlorine . . . . .	34·94	Cl <sup>2</sup>
Tin . . . . .	29·02	Sn
Carbon . . . . .	17·77	C <sup>6</sup>
Hydrogen . . . . .	2·47	H <sup>3</sup>
Oxygen . . . . .	15·80	O <sup>4</sup>
	100·00	

*Ann. de Ch. et de Phys.*, Mars 1846.

#### ANALYSIS OF TWO SPECIES OF EPIPHYTES, OR AIR PLANTS.

BY JOHN THOMSON, A.M.\*

I. *Commelina Skinneri*.—Until about four months prior to the time this plant was examined, it had roots in some of the pots; but about that time, Mr. Murray, of the Botanic Gardens, cut off all its roots, and left it hanging on the wall to which it had been trained. I had only 353·05 grains of the young shoots to operate on, so that very great precision cannot be expected in the results. After exposing this quantity on a sand-bath to a heat of about 280°, there remained 71·91 grains of the dried plant, so that the difference, which must have been almost wholly water, amounts to 281·14 grains. The dried portion was then burned: it left a residue of 7·14 grains of ashes, which were now subjected to analysis.

After treating the ashes with water to separate the soluble from the insoluble part, and evaporating the two portions to dryness, there were obtained of matters insoluble in water 4·22 grains, and of soluble substances 3·05 grains, the whole amounting to 7·27 grains, there being thus an excess of ·13 grain.

Muriatic acid was then poured on the insoluble portion, when a violent effervescence took place, and only ·77 grain remained undissolved. By fusing this with carbonate of soda, and adding muriatic acid in the ordinary way, there were found to be ·60 grain of silica. The whole quantity dissolved in muriatic acid was now mixed, and ammonia was added. A precipitate fell, which was boiled with caustic soda to remove alumina. What remained was evidently peroxide of iron; it was dried, and found to weigh ·22 grain. The portion dissolved by the caustic soda was precipitated by the addition of muriatic acid, the excess of which was removed by adding carbonate of soda. There were thus found to be ·44 grain of alumina, or phosphate of alumina.

To the washings oxalate of ammonia was added, and after filtering and burning, the precipitate weighed 2·90, which was carbonate of lime.

The next point was to determine the composition of the salts soluble in water. This part of the process was from an accident not completely executed. The only constituents which were deter-

\* Read before the Philosophical Society of Glasgow, December 4, 1844.

mined were the sulphuric acid, the potash, and the soda, the first of which was found, by precipitating with nitrate of barytes, to weigh .92 grain. The potash and soda were separated by means of bichloride of platinum and found to weigh respectively .24 grain and .94 grain. The following then is a statement of the entire results:—

	grs.		grs.		grs.	
Water.....	281.14					
Organic matter	64.77					
Ashes .....	7.14	{	Soluble in water 3.05	{	Sulph. acid . . . . .	.92
					Potash . . . . .	.24
		{	Insoluble in water 4.22	{	Soda . . . . .	.94
					Chlorine, &c. . . . .	.95
					Silica . . . . .	.60
					Perox. of iron . . . . .	.22
					Alumina . . . . .	.44
					Carb. of lime . . . . .	2.90
Entire plant .	353.05		7.27			

100 parts of the plant would contain,—

Water .....	79.64
Organic matter....	18.34
Ashes.....	2.02
	100.00

100 parts of the ashes again would contain approximately—

Soluble salts....	42.72	.....	42.72	
Insoluble .....	59.10	{	Silica .....	8.43
			Peroxide of iron .	3.08
			Alumina, or phos- phate of alumina }	6.16
			Carb. of lime ..	40.62
	101.82		101.01	

II. *Vanilla planifolia*.—The following is the composition of a specimen of the *Vanilla planifolia* which I also examined. Although called an epiphyte, it had roots in some of the pots. It is a very succulent plant with a small round stem, and alternate petiolated, elliptico-lanceolate, polished leaves:—

Water .....	89.06
Organic matter....	9.84
Ashes.....	1.10
	100.00

The ashes were similar in composition to those of the *Commelina Skinneri*. They contained no alumina, and had a perceptible quantity of phosphoric acid. It is probable therefore that the alumina in the first analysis is accidental.

These analyses were conducted under the direction of Dr. R. D. Thomson in the College Laboratory of Glasgow.

ANALYSIS OF *CERADIA FURCATA* RESIN.

BY ROBERT D. THOMSON, M.D.

The plant from which this resin exudes, presents the appearance of coral, and is a native of the coast of Africa, opposite the island of Ichaboe.

The resin possesses an amber colour, and an odour similar to that of olibanum. It partially dissolves in alcohol, and is precipitated by water. Caustic ammonia produces no precipitate in the alcoholic solution. The alcoholic solution possesses a slightly acid reaction, and is not precipitated by nitrate of silver. Specific gravity 1.197, determined by my pupil, Mr. Hugh B. Tennent.

Analysis gave the following results:—

19.9 grains lost by exposure to the temperature of 212° for some days 2.11 grains. During the whole of the period its peculiar odour was emitted. Previous to being subjected to this heat it was pulverized, but it speedily became soft, and collected into a mass. In this state, when burned with oxide of copper and chlorate of potash,

6.24 grains gave 18.33 grains CO<sub>2</sub>,  
and 5.50 ... HO.

This amounts to per cent.—

Carbon .....	80.113
Hydrogen .....	9.793
Oxygen .....	10.094
	<u>100.000</u>

Calculated according to the formula C<sub>10</sub>H<sub>7</sub>O, or C<sub>40</sub>H<sub>28</sub>O<sub>4</sub>, the result would be as follows:—

Carbon .....	80.00
Hydrogen .....	9.33
Oxygen .....	10.67
	<u>100.00</u>


After being heated in the water-bath for some weeks, the resin still continued to emit an odour. It was then pulverized, and again heated somewhat higher, when it speedily gave out fumes, and lost its smell entirely. Its composition was then found to be as follows:—

6.52 grains gave, with oxide of copper and chlorate of potash,—  
15.89 carbonic acid,  
5.02 water,

which are equivalent to—

Carbon .....	66.46
Hydrogen .....	8.55
Oxygen .....	24.99

Calculated according to the formula



its composition will be—

Carbon . . . . .	67·03
Hydrogen . . . . .	8·37
Oxygen . . . . .	24·60

From the *Proceedings of the Philosophical Society of Glasgow*, read February 5, 1845.

METEOROLOGICAL OBSERVATIONS FOR MARCH 1846.

*Chiswick*.—March 1. Overcast. 2. Very fine. 3. Cloudy. 4. Rain. 5. Showery: clear and fine. 6, 7. Very fine. 8. Clear: cloudy: clear. 9. Frosty: fine. 10. Frosty and foggy: fine: very clear. 11. Slight fog: very fine: clear. 12. Foggy. 13. Slight haze. 14. Cloudy and windy. 15. Showery. 16. Cloudy: boisterous: heavy showers. 17. Overcast: clear: slight frost at night. 18. Frosty: overcast: clear and frosty. 19. Frosty: overcast: hazy. 20. Snow early A.M., nearly two inches deep: cloudy: clear and frosty at night. 21. Sharp frost: densely clouded: boisterous, with rain at night. 22. Clear and fine: showery. 23. Rain: cloudy and fine: clear. 24. Cloudy and fine: clear. 25. Fine: overcast: showery. 26. Cloudy and fine. 27. Clear and fine. 28. Hazy. 29. Hazy clouds: fine. 30. Slight haze: cloudy and cold: clear. 31. Dry haze: clear and fine.

Mean temperature of the month . . . . .	43°·43
Mean temperature of March 1845 . . . . .	38·49
Mean temperature of March for the last twenty years . . . . .	42·89
Average amount of rain in March . . . . .	1·36 inch.

*Boston*.—March 1. Foggy. 2, 3. Cloudy. 4. Windy: rain early A.M. 5. Cloudy: rain early A.M. 6. Cloudy: rain P.M. 7—10. Fine. 11, 12. Cloudy. 13. Cloudy: rain at noon. 14. Cloudy: rain A.M. and P.M. 15. Cloudy. 16. Windy: stormy day: rain P.M. 17. Cloudy. 18. Fine. 19. Cloudy: snow early A.M. 20. Cloudy. 21—25. Fine. 26. Cloudy: thunder-storm, with rain P.M. 27, 28. Fine. 29. Fine: rain A.M. 30. Cloudy. 31. Fine.


*Sandwick Manse, Orkney*.—March 1. Bright: cloudy. 2. Clear: cloudy. 3. Showers: clear. 4. Bright: clear. 5. Fine: clear. 6. Clear. 7. Bright: hail-showers. 8. Showers: clear. 9. Damp: drops. 10. Damp: cloudy. 11. Clear: halo. 12. Cloudy: drops. 13. Cloudy: showers. 14. Sleet-showers: showers: sleet. 15. Sleet-showers: cloudy: sleet. 16. Sleet-showers: sleet. 17. Sleet-showers: snow. 18. Snow-showers: snow: cloudy. 19. Snow: clear. 20. Snow: clear: snow: cloudy. 21. Snow-drift: thaw: clear. 22. Cloudy. 23, 24. Bright: clear. 25. Rain: damp: clear. 26. Showers. 27. Showers: clear. 28. Clear. 29. Showers: cloudy. 30. Clear: cloudy. 31. Snow-showers: cloudy.

*Applegarth Manse, Dumfries-shire*.—March 1. Fine till 10: P.M. rain. 2. Heavy showers P.M. 3. Heavy rain all day. 4. Heavy rain all day: flood. 5. Very fine. 6. Showers. 7. Showers: hail: frost. 8. Hoar frost. 9. Slight showers. 10, 11, 12. Fine: fair. 13. Wet A.M. 14. Heavy rain A.M. 15. Rain P.M. 16. Showers: hail: sleet: rain. 17. Hard frost. 18. Frost: snow-showers. 19. Hard frost: clear. 20. Hard frost. 21. Frost: snow: hail: rain: thunder. 22. Rain: hail. 23. Slight drizzle: hail. 24. Showers. 25. Wet A.M.: cleared. 26. Hoar frost: drops. 27. Showers: hail. 28. Hail: rain. 29. Frost: clear and fine. 30. Frost: clear: cloudy. 31. Frost.

Mean temperature of the month . . . . .	42°·2
Mean temperature of March 1845 . . . . .	36·3
Mean temperature of March for 23 years . . . . .	39·0
Mean rain in March for 18 years . . . . .	2·35 inches.

*Meteorological Observations made by Mr. Thompson at the Garden of the Horticultural Society at CHISWICK, near London; by Mr. Veall, at BOSTON; by the Rev. W. Dunbar, at Applegarth Manse, DUMFRIES-SHIRE; and by the Rev. C. Clouston, at Sandwick Manse, ORKNEY.*

Days of Month.	Barometer.				Thermometer.						Wind.				Rain.					
	Chiswick.		Boston		Dumfries-shire.		Orkney Sandwick.		Chiswick.		Boston		Dumfries-shire.		Orkney Sandwick.		Chiswick.	Dumfries-shire.	Orkney Sandwick.	
	Max.	Min.	9 a.m.	9 p.m.	9 a.m.	9 p.m.	9 a.m.	9 p.m.	Max.	Min.	8 a.m.	8 p.m.	Max.	Min.	9 a.m.	9 p.m.				
1846. March.																				
1.	30.056	29.952	29.60	29.75	29.57	29.68	29.50	60	45	43	52	42½	48	48	calm	s.	calm	s.	...	...
2.	29.953	29.928	29.50	29.60	29.42	29.51	29.49	60	41	48	52	45½	46	45	calm	sse.	calm	sse.	...	...
3.	29.869	29.705	29.35	29.30	29.17	29.19	29.11	60	47	50	50	42½	45	46	sw.	s.	sw.	s.	...	...
4.	29.491	29.403	28.89	28.83	29.18	28.79	29.07	52	40	52	51½	46	48	43	sw.	s.	sw.	s.	...	...
5.	29.599	29.471	29.15	29.30	29.37	29.19	29.40	55	34	46.5	51	42½	42½	41	calm	s.	calm	s.	...	...
6.	29.729	29.549	29.17	29.37	29.30	29.39	29.30	53	41	45	49	35½	42	41	calm	s.	calm	s.	...	...
7.	29.777	29.715	29.35	29.48	29.62	29.38	29.60	53	28	44	50	35	45	44	calm	sw.	calm	sw.	...	...
8.	30.013	29.977	29.61	29.84	29.82	29.78	29.87	54	24	41	49	36	40	39	w.	w.	w.	w.	...	...
9.	30.305	30.105	29.75	30.00	30.14	30.00	30.12	56	27	37.5	49	34½	43	45	w.	w.	w.	w.	...	...
10.	30.383	30.368	29.99	30.20	30.11	30.06	30.14	58	29	41	51½	36½	43	47	w.	w.	w.	w.	...	...
11.	30.544	30.407	29.99	30.27	30.41	30.38	30.39	57	28	43	52	44½	46	41	sw.	e.	sw.	e.	...	...
12.	30.600	30.530	30.21	30.41	30.28	30.07	30.26	48	26	45	52	44½	47	47	ne.	calm	calm	s.	...	...
13.	30.434	30.336	30.00	30.09	30.05	29.88	29.84	51	39	43	54½	45	47	41	w.	w.	w.	w.	...	...
14.	30.155	30.001	29.65	29.65	29.76	29.53	29.55	56	47	49	53	43½	42	41	w.	w.	w.	w.	...	...
15.	29.988	29.882	29.55	29.79	29.46	29.71	29.44	57	45	48	50	38	39	37	w.	w.	w.	w.	...	...
16.	29.570	29.267	29.04	29.26	29.78	28.60	28.65	55	31	52	48	44½	43	34	sw.	w.	sw.	w.	...	...
17.	29.508	29.388	29.04	29.25	29.35	29.07	29.53	48	25	34	42	33	34	28	n.	w.	n.	w.	...	...
18.	29.576	29.513	29.25	29.43	29.50	29.56	29.62	47	23	34	36½	28½	29½	26	n.	calm	ne.	s.	...	...
19.	29.573	29.563	29.28	29.50	29.50	29.67	29.70	47	23	30	37	29½	33	29	sw.	calm	ne.	n.	...	...
20.	29.781	29.545	29.31	29.53	29.53	28.96	28.90	46	35	35	40	28	33	36	nw.	calm	ne.	s.	...	...
21.	29.643	29.258	29.30	29.13	28.87	28.96	28.90	46	35	35	40	28	33	36	sw.	calm	sse.	sse.	...	...
22.	29.203	29.177	28.83	28.78	28.88	28.88	28.91	53	29	43.5	47	35½	42	41	sw.	calm	sw.	se.	...	...
23.	29.317	29.169	28.85	28.88	28.95	28.85	28.91	55	37	39	48	35	41½	38	sw.	calm	sw.	ese.	...	...
24.	29.338	29.310	28.95	29.05	29.07	29.19	29.27	53	28	43	49½	39	43	40	s.	w.	se.	e.	...	...
25.	29.387	29.361	29.00	29.11	29.16	29.39	29.39	53	36	45	50	38½	42½	40½	w.	calm	n-s.	sw.	...	...
26.	29.710	29.443	29.07	29.54	29.46	29.31	29.46	54	37	41	50	33	40½	38	w.	calm	sw.	nw.	...	...
27.	29.790	29.734	29.36	29.50	29.48	29.45	29.66	58	27	43	48½	36	42	40½	sw.	calm	w.	nw.	...	...
28.	29.577	29.505	29.22	29.48	29.55	29.73	29.77	52	27	42.5	48	35	42½	37½	w.	calm	ene.	nne.	...	...
29.	30.097	29.807	29.44	29.81	29.99	29.92	30.05	57	35	43	49	33	39	38½	w.	calm	w.	calm	...	...
30.	30.116	29.858	29.77	30.00	29.81	30.07	30.90	54	30	42	51	32	41½	39½	e.	calm	n.	calm	...	...
31.	29.706	29.606	29.40	29.60	29.28	29.68	29.54	62	41	46	50	33½	37½	35	e.	calm	ne.	e.	...	...
Mean.	29.832	29.704	29.38	29.251	29.510	29.510	29.539	53.80	33.06	42.6	48.5	36.9	41.35	39.03	1.09	0.64	3.12	3.47		



THE  
LONDON, EDINBURGH AND DUBLIN  
PHILOSOPHICAL MAGAZINE  
AND  
JOURNAL OF SCIENCE.

---

[THIRD SERIES.]

---

JUNE 1846.

---

LXX. *Researches on the Functions of Plants, with a view of showing that they obey the Physical Laws of Diffusion in the Absorption and Evolution of Gases by their Leaves and Roots.* By D. P. GARDNER, M.D., Member of the Lyceum of Natural History, &c.\*

1. I CONCEIVE a plant to be a porous system, containing an internal mixture of gases, or plant atmosphere, and lying in contact with common air on the one side, and with the gases dissolved in the fluid of the soil on the other. My object in the following remarks is to show that the plant atmosphere is of a fluctuating nature, and depends on the chemical action taking place; and that whatever gases are absorbed or evolved by leaves or roots, depend upon the nature of the internal atmosphere at the time. To place the evidence of these conclusions before the reader, I propose to examine the subject under five heads:—

1st. The epidermis or bounding membrane of plants is porous.

2nd. The constitution of the internal gas of plants.

3rd. The action of roots on the gases of the soil-fluid.

4th. The absorption of gases by plants is a consequence of their porosity.

5th. The action of plants on artificial atmospheres.

---

I. *The Epidermis or Bounding Membrane of Plants is Porous.*

2. The object in this place is to show, that the epidermis is not merely capable of transmitting particular gases, but that it obeys all the laws of a porous system. If this be found true for the bounding membrane, it will necessarily be true for the internal cellular structure.

\* Communicated by the Author.

*Phil. Mag.* S. 3. Vol. 28. No. 189. June 1846.

2 G

3. Experiments were planned for the purpose of determining whether carbonic acid would penetrate into a vessel containing common air through a barrier of vegetable epidermis, and secondly, whether an inclosed mixture of gases of a theoretical composition would solicit the passage of both carbonic acid and oxygen, and at the same time evolve nitrogen.

4. A tube, five inches long and a third of an inch in bore, with a flattened and ground end, was closed by a piece of epidermis obtained from the leaf of the Madeira vine (*Basella lucida*). The tube was then immersed in a mercurial trough and filled to within an inch of the membrane; on suspending it by a wire it did not leak, though there was a pressure of three inches of mercury. Clear lime-water was admitted to displace the mercury, and the arrangement covered by a small bell-jar containing atmospheric air with 10 per cent. carbonic acid. In five hours the lime-water exhibited a distinct pellicle of carbonate of lime. The same result was obtained in more or less time with the epidermis of the cabbage, *Alanthus alata*, *Chenopodium album*, and several species of *Sedum*. Some specimens, as that from the balsam, leaked so fast as not to sustain any mercurial column, whilst others maintained four inches for thirty hours and more.

5. A similar tube was closed with epidermis, and contained an atmosphere of nitrogen 87, oxygen 13 per cent. over mercury, and was covered with a bell-jar as above. The included volume increased during nine hours from 400 to 433 measures, and on analysis consisted of N 76, O 17, CO<sub>2</sub> 7 per cent. Hence the membrane comported itself as a simple porous tissue, allowing nitrogen to pass out and admitting oxygen and carbonic acid. This experiment was also repeated with the foregoing specimens of epidermis, and no absolute variation perceived.

## II. *The Constitution of the Internal Gas of Plants.*

6. The observations hitherto made on the included gases of plants by Davy, Payen, Calvert and Ferrand, and others, cannot be quoted here, because the disturbing effects of light and other causes have not been sufficiently considered. It is not merely the gas of a cavity which is required, but the composition of that which permeates the interior during the vigorous state of the vegetable in sun-light.

7. For the purpose of obtaining this, I transplanted in the May of 1845 a number of plants of *Datura stramonium* and of blue grass (*Poa pratensis*) into tumblers, and allowed them to grow for several weeks before use. Having completed my arrangements for analysing minute quantities of gas by a sliding-



rod eudiometer, I proceeded as follows:—The plants when wanted were obtained in a perfect state by immersing the tumbler in a tub of water, and removing the garden-mould by agitating the fluid; they were thus procured without the slightest mutilation. The plant was then transferred to a convenient pneumatic trough, closed from adherent air and broken under a small receiver. This was done uniformly at 11 o'clock A.M. and as quickly as possible; the gas was immediately analysed.

8. Binoxide of nitrogen was the only substance which could be used to estimate the oxygen, and if properly prepared, is, in my experience, equal to the most intricate apparatus. Thus, in twenty-five analyses of air, made as test experiments of the excellence of my measures, there was obtained a mean of 20·83 per cent. oxygen, from which there never was a variation of 0·2 per cent.; this result closely coincides with that obtained in the elaborate researches of Dumas and Boussingault, *i. e.* 21·8 per cent.

9. Six analyses of the internal gas of *Datura* gave a mean of N 87·5, O 12·5 per cent. without any carbonic acid.

Four analyses of the gas from grass gave a mean of N 86·1, O 13·9 per cent. without carbonic acid. This result closely approximates to the measure published by Dr. Draper in the *Philosophical Magazine* for the gas drawn by the air-pump from grass.

The mean of all the observations is N 86·75, O 13·25 per cent.; and this I assume as the *normal* or plant atmosphere of the green parts at 11 to 12 o'clock A.M., during vigorous existence in the presence of bright diffused light in summer.

10. It is distinctly to be understood that this conditionally normal atmosphere is perpetually changing, and is true only for the time and place. In the preparatory examination of this subject, I obtained measures of the internal gas consisting of N 84·6, O 13·0, CO<sub>2</sub> 2·4 per cent., but overlooked the circumstances. Messrs. Calvert and Ferrand (*Ann. de Ch., &c.*, Aout 1844) found that carbonic acid was always present at night, and give as the composition of the gas from the hollow stems of *Phytolacca decandra* at night, N 76·4, O 20·6, CO<sub>2</sub> 0·3 per cent.

### III. *The Action of Roots on the Gases of the Soil-Fluid.*

11. There are no observations on the action of roots known to me, except those of DeCandolle (*Phys. Veg.* t. i. p. 248), who asserts that uninjured roots exhale no gas, either in light or darkness. Most physiologists infer, that whatever gases exist in the soil-fluid are absorbed therewith; but this is an

unphilosophical view, for it leaves out of consideration the capacity of the sap to absorb them and its condition as to gaseous saturation. In making experiments on the subject, it is also necessary to consider the functions of the plant.

12. On the 25th of June 1844, I commenced a series of observations to determine the action of uninjured roots of *Datura* and blue grass on the gas dissolved in pump-water, which accurately represents the soil-fluid. The plants were obtained as detailed in section 7; they were placed in vessels resembling a bird-fountain, which were capable of being replenished with water to compensate for the evaporation of the leaves, and also of collecting any gas passing from the roots. Three sets of experiments were made: A, the roots and leaves were placed in darkness; B, both portions were exposed to bright diffused light; C, the leaves were illuminated, but the roots in darkness.

13. On the evening of the 25th of June, two sets of plants were arranged according to these plans. The *Daturas*, B, yielded the next morning at 11, a gas the composition of which was N 96.6, O 3.4 per cent.; these two plants were then placed in a dark cupboard for thirty-six hours and evolved no gas whatever; on again exposing them to light, they produced a mixture of N 96.2, O 3.8 per cent. as the mean of six analyses. The grass plants, B, gave off but little gas, and only enough was collected for two measures, which yielded a mean of N 96, O 4 per cent.

The plants C conducted themselves in the same way as B; the *Daturas* gave gas for six analyses, the mean of which was N 96.5, O 3.5 per cent.

The plants A, placed in darkness, gave no gas whatever, although they were attended to for five days.

14. We conclude that roots appear to evolve gas unequally in quantity; that the action of light on the leaves is essential to this phænomenon; and thirdly, it, the exposure of the root, does not seem to have any effect on the result. I do not believe that the gas is evolved from the interior of the plant, but that the roots disturb the equilibrium of the mixture in the water, so that all the carbonic acid is withdrawn and most of the oxygen, leaving behind the sparingly soluble nitrogen, which acquires the elastic condition. That this gaseous disturbance was not a mechanical effect of light and heat, I satisfied myself by observations at the time; and the results of Prof. Morren (*Ann. de Chimie, &c.*, Sept. 1844) show that the sun's light liberates carbonic acid and nitrogen, accumulating oxygen in the water, which is opposed to the effects here observed.

15. The gas of the pump-water was N 48, O 22, CO<sub>2</sub> 30 per cent., therefore the roots absorbed carbonic acid and oxygen in the same way as a porous system containing the normal plant atmosphere, N 86·75, O 13·25 per cent.; this continued during daylight, or during the activity of the vegetable, but in darkness all the gas of the water is taken up without any selection.

IV. *The Absorption of Gases by Plants is a consequence of their Porosity.*

16. We are now in possession of sufficient data to state the case. A porous system lies between two media and contains a certain mixture of gases; the gases of the three systems are

	The air.	The plant-gas.	The water-gas.
Carbonic acid . . . . .	0·05	0·00	30·00
Oxygen . . . . .	20·80	13·25	22·00
Nitrogen . . . . .	79·15	86·75	48·00
	<hr/> 100·00	<hr/> 100·00	<hr/> 100·00

If the plant-gas were confined within an extremely delicate caoutchouc bag and surrounded by either atmosphere, it would soon be disturbed by penetration, nitrogen would be evolved, and carbonic acid and oxygen absorbed. The rapidity of the interchange would depend on the gas and the nature of the membrane.

17. That the action of roots on the water-gas is coincident with this theoretical view, I have attempted to show in the last division of the subject. The experiments detailed in article 5 were also made with this object. It must be remembered that the operation of the plant atmosphere upon the gases of the soil is not direct, as in the leaves, but through the intervention of the sap, which contains a mixture of gases dependent upon those of the interior of the plant, but has a greater capacity for carbonic acid and oxygen than for nitrogen.

18. In the case of leaves, the physical theory of porosity is more strikingly made out, because the internal gas is here in contact with atmospheric air, the epidermis only lying between. The movements witnessed in the experiments of art. 5, represent those of a vigorous plant in sun-light; carbonic acid and oxygen are absorbed and nitrogen evolved. It is not uniformly admitted that oxygen is absorbed and nitrogen evolved in plants, but an investigation of this point leaves us under the conviction that the negative is untenable. Saussure, DeCandolle, Palmer, Daubeny, Draper and others have witnessed the evolution of nitrogen. The continued absorption of oxygen is confirmed by Saussure, Davy, Gough, Achard,

Scheele, Cruickshank and others. Gough's interesting observation, that plants grown in darkness do not become green in light unless oxygen be present, is accounted for by the fact that chlorophyll is an oxidized product.

19. The movements occurring during light owe their continuance to its action. The gases absorbed are destined to equilibrate the internal atmosphere, but cannot effect this object so long as carbon and oxygen are fixed by the plant. Hence the current becomes continuous during daylight, the oxygen and carbon being removed faster than they penetrate.

20. But during darkness the stream is arrested, carbonic acid is no longer drawn from the air, but often evolved, as observed by Ingenhousz and Saussure. Oxygen is for a time required to satisfy chemical affinities, and afterwards the internal gas resembles atmospheric air; 20·6 per cent. oxygen is found, and amounts of nitrogen and carbonic acid differing with the nature of the soil-fluid, the latter gas sometimes rising to 3 per cent., when it is evolved; and at others not attaining a proportion much higher than that of the air, when it is not thrown off during night, as shown by Mr. Pepys.

21. It is the decomposition of the carbonic acid within the plant by the early sun-beams which creates the absorption from without by deranging the internal atmosphere. Carbonic acid being decomposed, a temporary excess of oxygen is produced, which causes a portion to pass outwards; but in preparing the plant for examination, any excess of this body seems to have been removed, so that it is probable that in the living organism oxygen gas is a much more important element than is usually admitted by physiologists.

#### V. *The Action of Plants on Artificial Atmospheres.*

22. To show beyond a doubt that the penetration of gases into plants is a physical and not vital process, we adduce the effects of artificial atmospheres. In these experiments the gases given out and absorbed are not of a definite mixture, but depend altogether on diffusion.

23. M. Marcet placed the same fungi in atmospheres of common air, oxygen, and nitrogen; and after eight to ten hours they changed 100 measures of—

	Air.	Oxygen.	Nitrogen.
into Oxygen . . .	2·0	31·3	0·0
Nitrogen . . .	77·0	24·0	96·1
Carbonic acid .	21·0	44·7	3·9
	<hr style="width: 50%; margin: 0 auto;"/> 100·0	<hr style="width: 50%; margin: 0 auto;"/> 100·0	<hr style="width: 50%; margin: 0 auto;"/> 100·0

without alteration of volume (*Ann. de Chimie, &c.*, t. lviii.

p. 407). It is evident the normal atmosphere of these fungi contained an excess of nitrogen and carbonic acid. Th. de Saussure found that seeds germinating in air absorbed nitrogen, but when placed in a mixture of N 50, O 50, they no longer did so (*Mem. de la Soc. de Genève*, t. vi. p. 545).

24. By overlooking the laws of penetration, DeCandolle, Saussure, Ingenhousz, and Plenck are thrown into contradictory positions in their experiments on the action of the green parts of plants on artificial atmospheres. Thus DeCandolle (*Phys. Veg.* t. i. p. 133), "Les parties verts laissent moins de gas oxigène dans le gas hydrogène que dans le gas azote; elles ne paraissent, contre l'assertion d'Ingenhousz, absorber ni l'un ni l'autre. Il parait aussi certain, malgré l'assertion de Plenck, qu'elles n'exhalent point de gas azote, sauf dans quelques cas, par les corolles."

25. In the summer of 1844 I tested this question by placing some specimens of the *Conferva mucosa* in pump-water and in carbonated water, and allowing them to act for several days on the same water, removing each day the gas generated during light; the plants were therefore subjected in their natural medium to different mixtures of gases dissolved in the fluid. The result was, that the plants in pump-water gave in six hours a gas consisting of O 73, N 27 per cent.; in twenty-four hours, O 53, N 47 per cent.; in forty-eight hours, O 18·6, N 81·4 per cent. In carbonated water, in six hours, O 68 per cent.; in twenty-four hours, O 63 per cent.; in forty-eight hours, O 12, N 88 per cent.; in seventy-two hours, O 3·5, N 96·5 per cent. And these plants continued healthy and acted as before when fresh water was added.

26. *Conclusion.*—From the preceding evidence I infer that plants constitute a simple porous system. The advantages resulting from this philosophical view of vegetation, both in assimilating facts hitherto insulated and in criticising experimental arrangements in vegetable physiology, constitute its chief recommendation. For illustration, we adduce two general laws springing from this theory:—1st. No hypothesis nor argument can be based on the composition of the gases expired by plants without the strictest regard to the disturbing influence of light, the gases of the soil-fluid. 2ndly. No experiments on the action of plants in sun-light can be accepted in determining the functions of leaves unless made in atmospheric air.

27. Finally, I beg to present the following summary of conclusions as fairly deducible from the preceding experiments:—

1. The epidermis of plants, so far as experiments have been made, is porous, and permits the passage of gases according to physical laws.

2. The roots, during the existence of chemical changes in plants, absorb such gases from the soil-fluids as will indirectly satisfy the requisitions of the internal atmosphere.

3. The internal gas of plants fluctuates with the forces which operate on the plant; during the active state of the green vegetable, it resembles a mixture of nitrogen 86·75, oxygen 13·25 per cent., but at night contains more oxygen and a proportion of carbonic acid.

4. The porosity of the entire plant is fully established by its action on artificial atmospheres.

Therefore the physical structure of plants is that of a porous system subject to the laws of diffusion of gases, and endowed with no vitality other than the power of forming cytoblasts and arranging cellules after a definite type.

### LXXI. On the relation of Ozone to Hyponitric Acid.

By Dr. C. F. SCHŒNBEIN\*.

THE chemical effects produced by atmospheric air charged with hyponitric acid ( $\text{NO}_4$ ) are so very like those caused by ozonized air, that some chemists are inclined to consider hyponitric acid as identical with ozone. Both decompose iodide of potassium, transform the yellow prussiate of potash into the red one, decompose sulphuretted hydrogen, colour blue the resin of guaiacum, destroy organic colouring matters, polarize negatively platinum, &c.

In spite of the similarity of chemical properties exhibited by ozone and hyponitric acid, those substances are in many other respects so entirely different that their being identical cannot be thought of: thus ozone is produced under circumstances in which an essential constituent part of hyponitric acid is absent, namely nitrogen. The analogy existing between the chemical action of the two bodies mentioned appears however so striking, that we can hardly help suspecting some connexion to exist between them, and with the view of ascertaining that connexion I have of late made many experiments, the account of which will form the substance of this paper.

The results obtained from these researches are, in my opinion, such as to speak strongly in favour of my conjecture that there exists a compound composed of  $\text{NO}_2 + \text{HO}_2$ . It

\* Communicated by the Chemical Society; having been read November 3, 1845.

appears to me probable that besides hydrate of nitric acid a peroxide of nitrogen and hydrogen is formed when water acts upon hyponitric acid. Agreeably to my view, the presence of  $\text{NO}_2 + \text{HO}_2$ , in the acid mixture, is the cause why the latter decomposes iodide of potassium, &c.

I think it cannot be denied that aqueous vapour will act upon vaporous hyponitric acid as liquid water does upon the liquid acid. Such being the case, and supposing that in the latter case two compounds of the formulas  $\text{NO}_4 + \text{HO}_2$  and  $\text{NO}_2 + \text{HO}_2$  are produced, it must then be admitted also that the same compounds are formed when the vapours of hyponitric acid are mixed with moist atmospheric air. Now if the peroxide of nitrogen and hydrogen (the first of these compounds) should happen to be a volatile substance, it follows further that in a bottle containing moist air, to which vapour of hyponitric acid had been added, an atmosphere must be produced containing some  $\text{NO}_2 + \text{HO}_2$ , and possessing the property of causing the reactions above mentioned. The peroxide of hydrogen contained in  $\text{NO}_2 + \text{HO}_2$  would be the oxidizing agent, decomposing *e. g.* iodide of potassium, &c.

For various reasons I am inclined to consider ozone as  $\text{HO}_2$ , *i. e.* a compound isomeric with Thenard's peroxide of hydrogen. Now if this conjecture should be true, and if there should exist a compound of  $\text{NO}_2 + \text{HO}_2$ , moist atmospheric air being charged with vapour of hyponitric acid would owe its reactions to the presence of ozone. How far such a conjecture is founded upon facts, the experiments I am about to detail will show.

If a piece of carbonate of ammonia, having been strongly ozonized by the means of phosphorus, is suspended in atmospheric air until the latter be so charged with ammoniacal vapours as rapidly to restore the blue colour of reddened litmus paper, that atmosphere continues to enjoy bleaching powers, decomposes iodide of potassium, colours blue the paste of starch containing that salt, transforms the yellow prussiate of potash into the red one, colours the resin of guaiacum blue, discharges the colour of sulphuret of lead; continues, in one word, to possess all the properties belonging to ozone. Hence it follows that ozone is capable of co-existing with the vapours of carbonate of ammonia without suffering decomposition; and I have ascertained that pure ammonia also does not perceptibly destroy ozone. Now if there exists  $\text{NO}_2 + \text{HO}_2$ , we may presume from its constitution that it will likewise be able to co-exist with the vapours of the carbonate of ammonia,  $\text{NO}_2$  being of itself inactive towards ammonia.

Hyponitric acid, or red fuming nitric acid, was gradually mixed up with so much water as to obtain a colourless liquid.

The bottom of a spacious bottle was covered with this mixture, and then a large piece of carbonate of ammonia suspended within the vessel. After the atmosphere, standing over the acid liquid, had assumed the power of colouring rapidly blue a strip of reddened litmus paper, it continued to possess the following properties:—

1. Strips of paper charged with paste of starch containing some iodide of potassium were coloured blue.

2. Strips of paper drenched with an alcoholic solution of guaiacum assumed a blue colour.

3. Strips of paper coloured blue by a solution of indigo turned white.

4. Strips of paper to which sulphuret of lead had been attached, by means of nitrate of lead and sulphuretted hydrogen, gradually turned white.

5. Strips of paper charged with a solution of the common prussiate of potash became deeply yellow.

6. Crystals of the yellow prussiate, after having been suspended for twenty-four hours within this atmosphere, were covered with a crust of the red sesqui-ferrocyanuret of potassium.


From the facts just stated, it appears that the atmosphere in question acts exactly in the same way as ozonized air does, and from the circumstances under which those reactions took place, it follows that the latter could not proceed from free hyponitric or nitrous acid, these acids not being able to co-exist in a state of isolation with the vapours of carbonate of ammonia. We must therefore conclude from these facts, that there was a principle present, in the atmosphere mentioned, which acted after the manner of ozone, and conducted itself, in spite of the presence of ammoniacal vapour, as a highly oxidizing agent.

But if neither free hyponitric nor nitrous acid were the cause of the reactions mentioned, nor nitrite of ammonia, what then is the substance to which the oxidizing powers are to be ascribed? I can answer that question only by supposing that the peroxide of nitrogen and hydrogen is that agent. Before passing to another subject, I take the liberty to mention a circumstance which seems to bear upon the matter in question, and merit some attention. On breathing strongly ozonized air three or four times, a disagreeable and strangling sensation will be experienced near the throat and in the chest. This sensation is very similar to that caused by inhaling air which has stood for some time over a mixture of hyponitric acid and water, and this is the case even if the air happens to be charged with ammoniacal vapours. We observe also in such an atmosphere a peculiar and disagreeable odour,



which is similar to that of common aquafortis, and slightly analogous to the smell of chlorine. It is likely that the odour mentioned belongs to the vapour of our supposed peroxide of nitrogen and hydrogen, and that it is *that* compound which, if inhaled, causes the sensation before mentioned. With regard to the subject under discussion, it seems to me that the way in which the mixture, formed from hyponitric acid and water, acts upon a solution of the yellow prussiate of potash, offers peculiar interest.


My experiments having demonstrated that the salt just mentioned (be it solid or dissolved in water) is readily transformed into the red cyanuret by ozone, and considering the acid mixture before alluded to as an aqueous solution of nitric acid and peroxide of nitrogen and hydrogen, what must happen if that mixture be put together with a solution of the yellow prussiate of potash? Supposing 1 equiv. of the said peroxide and 1 equiv. of nitric acid in the acid mixture to act upon 2 equiv. of the yellow cyanuret, we must obtain from such a reaction 1 equiv. of nitrate of potash, 1 equiv. of the red sesqui-ferrocyanuret of potassium, 1 equiv. of deutoxide of nitrogen, and 2 equiv. of water, for



If a glass tube, open at one end, be half-filled with our acid mixture and half with a dilute solution of the yellow prussiate, on mixing the liquids together a lively disengagement of gas takes place; and if the open end of the tube be put into a vessel holding water, a colourless gas will fill the upper part of the tube. On adding oxygen or atmospheric air to the gas disengaged under the circumstances mentioned, the latter will turn brownish red and exhibit all the properties of deutoxide of nitrogen. As soon as the acid mixture comes in contact with the nearly colourless solution of the common prussiate of potash, the latter turns deeply yellow, and it is very easy to ascertain that the coloured fluid contains nitrate of potash, sesqui-ferrocyanuret of potassium, and no trace of the yellow prussiate, provided a sufficient quantity of the acid mixture had been employed. The reactions indicated are therefore such as they ought to be, if, according to our supposition, the acid mixture contains nitric acid and peroxide of nitrogen and hydrogen. I need hardly say that the disengagement of deutoxide of nitrogen and the transformation of the yellow cyanuret into the red one, which take place under the circumstances mentioned, cannot originate in the nitric acid contained in the acid mixture, for it is well known that dilute pure nitric

acid does not give rise to the reactions described. The cause of those effects must therefore be sought in another substance, and both hyponitric and nitrous acid not being able to co-exist with free water, we are not allowed to consider either the one or the other of those acids as that cause. As I have already remarked, the reactions in question can, according to my opinion, only be accounted for in a satisfactory manner by admitting the presence of  $\text{NO}_2 + \text{HO}_2$  in the acid mixture so often mentioned. My experiments have further shown that iodide of potassium, be it solid or dissolved in water, is readily decomposed by ozone, iodine being eliminated. In putting our acid mixture to a solution of the salt mentioned, deutoxide of nitrogen is abundantly disengaged, iodine precipitated, and nitrate of potash formed. As pure nitric acid containing as much water as the said acid mixture, does not act upon the solution of iodide of potassium, it cannot be the nitric acid of our mixture that causes those phænomena, nor can they, from reasons already stated, proceed from hyponitric or nitrous acids.

If we admit that there is, besides nitric acid, some  $\text{NO}_2 + \text{HO}_2$  present in the acid mixture, I think we may easily account for those reactions.  $\text{HO}_2$  oxidizes the potassium of the iodide, nitric acid unites with the base thus formed, and  $\text{NO}_2$  is set free.



The facts I am about to state are most likely also connected with the subject under discussion.

Largely diluted pure nitric acid, not colouring in the least paste of starch containing chemically pure iodide of potassium, when put for a short time in contact with a number of metals, as zinc, iron, lead, copper, mercury, silver, &c., acquires the property—1. To colour deeply blue the paste mentioned. 2. To transform the yellow cyanuret into the red one. 3. To colour blue the resin of guaiacum. 4. To decompose sulphuretted hydrogen, &c.

Tin is an exception to the rule, for however long dilute nitric acid may have been in contact with that metal, it does not cause the reactions indicated. On the contrary, an acid having the properties mentioned, loses them when mixed in proper quantities with dilute nitric acid which has been in contact with tin. The latter acid has also the power of discharging the colour of paste of starch rendered blue by iodine. From the facts stated, it seems to follow, that when dilute nitric acid is acted upon by oxidable metals, the same oxidizing

agent is produced that forms on mixing hyponitric acid with water; and it appears also that tin engenders with dilute nitric acid a deoxidizing matter, *i. e.* a nitrate of protoxide of tin.

LXXII. *On the Composition of the Fire-Damp of the Newcastle Coal Mines.* By THOMAS GRAHAM, Esq., F.R.S.\*

SOME years ago I examined the gas of these mines, with the same result as Dr. Henry, Davy and Dr. Turner had previously obtained, namely, that it contains no other combustible ingredient than light carburetted hydrogen. But the analysis of the gas of the coal mines in Germany, subsequently published, showing the presence of other gases, particularly of olefiant gas, has rendered a new examination of the gas of the English mines desirable. The gases were,—1, from a seam named the Five-Quarter seam, in the Gateshead colliery, where the gas is collected as it issues, and used for lighting the mine; 2, the gas of Hebburn colliery, which issues from a bore let down into the Bensham seam—a seam of coal which is highly charged with gas, and has been the cause of many accidents; and 3, gas from Killingworth colliery, in the neighbourhood of Jarrow, where the last great explosion occurred. This last gas issues from a fissure in a stratum of sandstone, and has been kept uninterruptedly burning, as the means of lighting the horse-road in the mine, for upwards of ten years, without any sensible diminution in its quantity. The gases were collected personally by my friend Mr. J. Hutchinson, with every requisite precaution to ensure their purity, and prevent admixture of atmospheric air.

The usual eudiometrical process of firing the gases with oxygen was sufficient to prove that they all consisted of light carburetted hydrogen, with the exception of a few per cent. The results were as follows:—

*Gateshead Gas.*—Specific gravity 0·5802.

Carburetted hydrogen . . . . .	94·2
Nitrogen . . . . .	4·5
Oxygen . . . . .	1·3
	100·0

The density of such a mixture is, by calculation, 0·5813.

*Killingworth Gas.*—Specific gravity 0·6306.

Carburetted hydrogen . . . . .	82·5
Nitrogen . . . . .	16·5
Oxygen . . . . .	1·0
	100·0

\* Communicated by the Chemical Society; having been read November 3, 1845.

The theoretical density of this gas, deduced from its composition, is 0.6308.

The Hebburn gas was of specific gravity 0.6327.

Seventy-nine measures of the Killingworth gas, mixed with an equal volume of chlorine, left in the dark for eighteen hours, and afterwards washed with alkali, were reduced to 75 measures; from which the presence of 4 measures of olefiant gas might be inferred. But in a comparative experiment made at the same time on 25.3 measures of pure gas of the acetates, mixed with an equal volume of chlorine, a contraction occurred of 1.3 measure; that is, in exactly the same proportion as with the fire-damp.

It was observed that phosphorus remains strongly luminous in these gases, mixed with a little air, while the addition to them of one-four-hundredth part of olefiant gas, or even a smaller proportion of the volatile hydrocarbon vapours, destroyed this property. Olefiant gas itself, and all the allied hydrocarbons, were thus excluded.

Another property of pure light carburetted hydrogen, observed by myself, enabled me to exclude other combustible gases, namely, that the former gas is capable of entirely resisting the oxidating action of platinum black, and yet permits other gases to be oxidated which are mixed with it even in the smallest proportion, such as carbonic oxide and hydrogen, the first slowly and the last very rapidly; air or oxygen gas being, of course, also present in the mixture. Now platinum black had not the smallest action on a mixture of the gas from the mines with air. No moisture appeared or sensible contraction, and no trace of carbonic acid could be discovered after a protracted contact of twenty-four hours; while, with the addition of one per cent. of hydrogen, the first effects were conspicuously evident in three minutes, and with the same proportion of carbonic oxide, the gas became capable of affecting lime-water in half an hour. These experiments were repeated upon each of the three specimens of fire-damp.

Potassium fused in the fire-damp did not become covered with the green fusible compound of carbonic oxide, nor occasion any contraction. Indeed, however carefully the heat was applied to the potassium by means of an oil-bath, a slight permanent expansion always ensued. The same thing occurred in pure gas of the acetates. It appeared that potassium could not be heated above 300° Fahr. in pure carburetted hydrogen, without causing a decomposition and the evolution of free hydrogen gas.

The gas was also inodorous, and clearly contained no appreciable quantity of any other combustible gas than light

carburetted hydrogen. The only additional matters present were nitrogen and oxygen; the specimen collected in the most favourable circumstances for the exclusion of atmospheric air, namely, that from the Bensham seam, still containing 0·6 per cent. of oxygen. The gases also contained no carbonic acid.

It is worthy of observation, that nothing oxidable at the temperature of the air is found in a volatile state associated with the perfect coal of the Newcastle beds. The remarkable absence of oxidability in light carburetted hydrogen appears to have preserved that alone of all the combustible gases originally evolved in the formation of coal, and which are still found accompanying the imperfect lignite coal of Germany, of which the gas has been examined. This fact is of geological interest, as it proves that an almost indefinitely protracted oxidating action of the air must be taken into account in the formation of coal; air finding a gradual access through the thickest beds of superimposed strata, whether these strata be in a dry state or humid.

In regard to measures for preventing the explosion of the gas in coal mines, and of mitigating the effects of such accidents, I confine myself to two suggestions. The first has reference to the length of time which the fire-damp, from its lightness, continues near the roof, without mixing uniformly with the air circulating through the workings. It was found that a glass jar, of six inches in length and one inch in diameter, filled with fire-damp, and left open with its mouth downwards, continued to retain an explosive mixture for twenty minutes. Now it is very desirable that the fire-damp should be mingled as soon as possible with the whole circulating stream of air, as beyond a certain degree of dilution it ceases to be explosive. Mr. Buddle has stated, "that immediately to the leeward of a blower, though for a considerable way the current may be highly explosive, it often happens that after it has travelled a greater distance in the air-course, it becomes perfectly blended and mixed with the air, so that we can go into it with candles; hence, before we had the use of the Davy lamp, we intentionally made 'long runs,' for the purpose of mixing the air." It is recommended that means be taken to promote an early intermixture of the fire-damp and air; the smallest force is sufficient for this purpose; as a downward velocity of a few inches in the second will bring the light gas from the roof to the floor. The circulating stream might be agitated most easily by a light portable wheel, with vanes, turned by a boy, and so placed as to impel the air in the direction of the ventilation, and not to impede the draft. The gas at the roof undoubtedly often acts as an explosive train,

conveying the combustion to a great distance through the mine, while its continuity would be broken by such mixing, and an explosion, when it occurred, be confined within narrower limits.

Secondly, no effective means exist for succouring the miners after the occurrence of an explosion, although a large proportion of the deaths is not occasioned by fire, or injuries from the force of the explosion, but from suffocation by the after-damp, or carbonic acid gas, which diffuses itself afterwards through all parts of the mine. It is suggested that a cast-iron pipe, from eight to twelve inches in diameter, be permanently fixed in every shaft, with blowing apparatus, above, by which air could be thrown down, and the shaft itself immediately ventilated after the occurrence of an explosion. It is also desirable that, by means of fixed or flexible tubes, this auxiliary circulation should be further extended, and carried as far as practicable into the workings.

LXXIII. *Observations on the Resin of the Xanthorœa hastilis, or Yellow Gum-resin of New Holland.* By JOHN STENHOUSE, Esq., Ph.D.\*

**T**HIS remarkable resin, which is known in commerce as the yellow gum or acaroid resin of Botany Bay, exudes from the *Xanthorœa hastilis*, a tree which grows abundantly in New Holland, especially in the neighbourhood of Sidney. This resin was first described in Governor Phillips's Voyage to New South Wales in 1788. Mr. Phillips states that it was employed by the natives and first settlers as a medicine in cases of diarrhœa. The resin as it occurs in commerce sometimes forms masses of considerable size, but as it is very brittle, although tolerably hard, it usually arrives in the state of a coarse powder. Its colour is a deep yellow, with a slightly reddish shade, considerably resembling gamboge, but darker and less pleasing. The colour of its powder is greenish yellow. When chewed it does not dissolve or stick to the teeth, but tastes slightly astringent and aromatic like storax or benzoin. Its smell is very agreeable and balsamic. When gently heated it melts, and when strongly heated it burns with a strong smoky flame, and emits a fragrant odour resembling balsam of Tolu. The resin contains a trace of an essential oil, to which much of its agreeable smell is probably owing. This oil passes into the receiver when the resin is

\* Communicated by the Chemical Society; having been read November 17, 1845.

distilled with a mixture of carbonate of soda and water, but its quantity is so small that I was unable to examine it more closely. The resin is insoluble in water, but dissolves readily both in alcohol and in æther, especially in the former. Its solution in alcohol has a brownish yellow colour; the addition of water precipitates it as a dark yellow mass, but it does not crystallize out of its alcoholic solution when left to spontaneous evaporation, but remains as a varnish. When digested with strong alkaline lyes, it readily dissolves and forms a brownish red solution; and when the alkali is neutralized with muriatic acid, the resin is precipitated considerably altered as a dark brownish brittle mass. On concentrating the solution out of which the resin has been precipitated, and allowing it to cool, a quantity of impure reddish crystals resembling benzoic acid are gradually deposited. It requires repeated and long-continued digestions with the strongest alkaline lyes to remove the whole of this crystalline acid from the resin, which retains it with very great tenacity. The quantity of the acid is by no means great. It is not easily purified, as its crystals are apt to retain a trace of a reddish colouring matter, from which it is very difficult to free them. The easiest way of getting rid of it, is by dissolving the impure crystals in a small quantity of alcohol and then adding water; the greater portion of the colouring matter is retained in solution, while the crystals are precipitated tolerably white. When purified by repeated crystallizations, they become quite colourless. In appearance, taste, and smell they closely resemble benzoic acid. When dried at  $212^{\circ}$  F. and subjected to analysis,—

I. 0.2284 grm. of substance gave 0.6005  $\text{CO}_2$  and 0.113 HO.

II. 0.2955 grm. of substance prepared on a different occasion gave 0.790  $\text{CO}_2$  and 0.1505 HO.

	Found.		Cinnamic acid.	Benzoic acid.
	I.	II.		
C	71.74	72.91	73.35	68.85
H	5.49	5.65	5.32	4.91
O	22.77	21.44	21.33	26.24
	<u>100.00</u>	<u>100.00</u>	<u>100.00</u>	<u>100.00</u>

It is evident from these analyses that the crystalline acid contains nearly the same amount of carbon and hydrogen as cinnamic acid, with some deficiency however in the carbon. I was led therefore to suspect that it consisted essentially of cinnamic acid, with probably a small admixture of benzoic acid, a suspicion which subsequent experiments tended fully to confirm; for on heating a quantity of the crystals with some peroxide

of manganese and sulphuric acid, oil of bitter almonds was immediately evolved, and on boiling a second portion with hypochlorite of lime, the very peculiar chlorinated oil described in a former paper was also abundantly produced, thus clearly indicating the presence of cinnamic acid. A third portion of the crystals was dissolved in alcohol and left to spontaneous evaporation; it yielded after some time the fine rhombic prisms so characteristic of cinnamic acid when it is crystallized out of alcohol, mixed however with some long acicular crystals, having all the appearance of benzoic acid. I think myself warranted to conclude therefore that Botany Bay resin contains cinnamic acid mixed with a very little benzoic, in which respect it resembles balsam of Tolu, which contains both cinnamic and benzoic acids, though fortunately in much greater abundance.

*Action of Nitric Acid on the Resin.*

When the resin is treated with moderately strong nitric acid in the cold, a violent action ensues with the evolution of nitrous fumes. The resin is completely dissolved if the quantity of the nitric acid is considerable. The colour of the solution is dark red, but by boiling it becomes of a bright yellow colour. The liquid should be evaporated to dryness on the water-bath, to get rid of the great excess of nitric acid. The residue forms a mass of fine yellow crystals, consisting chiefly of carbazotic acid, but mixed with some oxalic and a little nitrobenzoic acids. The nitrobenzoic acid is evidently derived from the cinnamic acid in the resin. The carbazotic acid is easily separated from these other acids by converting it into carbazotate of potash, which is easily purified by one or two crystallizations, and then by decomposing the salt with muriatic acid, pure carbazotic acid may be obtained.

0.3823 grm. of the acid, dried at 212° F., gave 0.442 CO<sub>2</sub> and 0.049 HO.

	Found.	Calculated numbers.
Carbon . . .	31.53	31.37
Hydrogen . .	1.42	1.30
Oxygen . . .	67.05	67.33
	100.00	100.00

0.3975 grm. of the potash salt, decomposed by sulphuric acid and then ignited with carbonate of ammonia, left 0.1300 of sulphate of potash = 17.68 per cent. of potash; calculated quantity 17.60.

The silver salt was also formed by boiling the acid with carbonate of silver. It is a very soluble salt, which crystallizes in fine red-coloured needles. 0.8975 grm. of the salt gave 0.372



Cl Ag = 31.22 Ag, or 33.53 per cent. oxide. The calculated numbers are 31.27 per cent. of silver = 33.59 oxide.

The quantity of carbazotic acid which Botany Bay resin yields when treated with nitric acid is so great, and it is so easily purified, that this resin seems likely to prove the best source of that substance. When the resin is subjected to destructive distillation in an iron or copper retort, it yields a very large quantity of a heavy acid oil mixed with a very small quantity of a neutral oil, which is lighter than water. If however the resin has been previously digested with alkaline lyes, so as to remove all the cinnamic and benzoic acids it contains, the heavy oil is obtained as before, but none of the light essential oil. The acid oil is readily soluble in potash and soda lyes; in its smell and properties it resembles creosote; when it is digested with nitric acid, it is wholly converted into carbazotic acid, and when a slip of fir-wood is dipt in it, and then moistened with either muriatic or nitric acid, the deep blue colour passing quickly into brown, so characteristic of hydrate of phenyle, is immediately produced, with which substance the oil appears completely identical. The light oil above mentioned, the quantity of which is extremely small, is separated from the hydrate of phenyle by saturating it with an alkali and distilling the mixture in a glass retort with a gentle heat. In smell and properties it resembles benzine, and is most probably a mixture of benzine and cinnamene; unfortunately the quantity obtained was so small, that I was unable to subject it to more particular examination.

#### LXXIV. *On the Constitution of Matter.*

*By H. SLOGGETT, Esq.*

*To Richard Taylor, Esq.*

SIR,

HAVING observed in your Journal for December 1845 some remarks on Prof. Faraday's speculation on the constitution of matter by Mr. Laming, wherein he attempts to show, that by a peculiar way of considering the theory of atoms the conducting and insulating powers of bodies appear more intelligible than on any other doctrine, I have been induced to send you a few ideas of mine on the subject, with a hope that you may not consider them unworthy of insertion.

The test of the truth of any hypothesis, is its accordance with all known facts; and any discrepancy, even a single one, between a theory and experiment, is, if not cleared up, fatal to its validity. The one-fluid theory, in electricity though pre-

ferable to its rival in perhaps all other respects, has hitherto been incapable of being generally received, on account of its giving unsatisfactory results when submitted to mathematical analysis. In endeavouring to obviate this difficulty, I conceived the object completely attained, by a supposition somewhat analogous to Mr. Laming's, though essentially different in respect to the assumption of solid atoms. As we know matter only by its properties, it certainly seems more rational to call those properties themselves matter, than to invent an imaginary substance with inseparably attached attributes.

In Mr. Laming's theory I can see no vindication of the theory of solid atoms, because it is not necessary for them to be admitted. He might consider the term "atom" to signify nothing more than "centre of attraction." With this qualification I agree with him, that different atoms are naturally associated with different quantities of electricity; arising, however, from different degrees of power in the atoms, or centres of attraction. An objection apparently arises, *in limine*, to his supposition of incomplete external strata. How are they to remain incomplete when placed in circumstances adapted to supply them with as much electricity as would be necessary to complete them? Or in other words, why should they not retain the electricity which they have once received? I have mentioned this objection because it appears to be an essential point in Mr. Laming's paper, and because it is in fact the only one, except that before mentioned, so far as it goes, in which his hypothesis differs materially from my own. It will be proper to premise, that by the word atom, I mean nothing more than a centre or combination of centres of attractive or repulsive force; those combinations of centres, when they occur, occupying the same point; implicating, in opposition to the usual notion, that matter may be penetrable. It is necessary that this be remembered, because the word in its common sense involves circumstances incompatible with another meaning.

Philosophers have long considered it established that the atoms of bodies attract each other; and it cannot but be admitted that a repulsive principle between them is just as clearly evidenced. Hence we have just grounds for the assumption that the atoms of bodies are both attractive and repulsive of each other. But this is an inconsistency if an atom be but a single principle, and as there is abundant proof of the existence of an agent distinct from matter in bodies, there are ample reasons for attributing the attracting property to this agent (electricity), and the repulsive power to the matter itself. But it remains to be shown how these are to be united in order to

explain the effects which appertain to the action of bodies in general on each other, as well as those which are produced by the agency of electricity.

We assume, then, that the atoms of matter are mutually repulsive of each other, but attractive of those of electricity; and that the atoms of electricity are in like manner self-repulsive and attractive of those of matter. This hypothesis is not new, it was invented long ago to satisfy the Franklinian theory of electricity; but its application has not, to my knowledge, been successfully made. My object here is briefly to show its consistency when rightly applied.

Suppose the centres of matter far more powerful and less numerous than those of electricity. Each atom of the compound will thus consist of an atom (in the sense before stated) of matter surrounded with an atmosphere (so to speak) of electricity, of variable density, in a somewhat similar manner to the air surrounding the earth. This must have a definite limit at some distance from the centre, where the repulsive power of the whole quantity of electricity surrounding the matter on an atom of electricity equals the attractive power of the matter for the same atom; so that beyond this limit none can exist in connexion with the atom. Accordingly every particle of matter will appropriate to itself a definite quantity of electricity dependent on its inherent power; and when any excess above this quantity occurs in a body, it becomes positively electrified, and negatively electrified when there is a deficiency. This admitted, we may enunciate thus: In all bodies, in their natural state, there are two principles reciprocally combined, mutually attractive but each repulsive of itself. If there be an excess of either principle, in one instance the body in which it may occur becomes positively, and in the other negatively electrified.

This will be observed, in effect, to be expressing the two-fluid theory. A simple illustration will exemplify the similarity.

Suppose a conducting sphere charged positively. All its atoms being duly combined with as much as they can retain, it is evident that the superfluous electricity thus thrown on them must, by its elastic property, fly off from them, subject only to an inferior attractive and repulsive force, it being as it were without the effective range of the central forces. Unless retained on the body by a non-conducting medium, it would necessarily fly off entirely. This both the old theories teach us. But how will the case stand when the excitement is negative? This is a question which the partisans of the vitreous and resinous hypotheses were accustomed triumphantly to ask. Indeed I have never seen it answered; arising from

no assigned position being given to electricity in its combination with matter. This being done, the solution is easy, and if it be done satisfactorily, the two-fluid theory must be estimated as nothing but a superfluity. In a sphere, the combined atoms of matter concentrate their power in its centre. Thus the exterior atoms have their atmospheres less strongly attached to them than the interior ones, since the tendency of all the combined atoms to attract electricity to the centre of the sphere is greatest at the surface, varying as the distance from the centre; and because this tendency and the force with which the atom attracts its atmosphere are in opposition. Now it is manifest that a force must act when the resistance to it is least; hence if electricity be abstracted from the sphere it must be from the surface, or rather from the atoms on the surface. If the surface be not covered with a non-conducting medium, the atoms will necessarily supply their deficiency from the contiguous conducting ones, and thus cause an equilibrium. It has however yet to be shown how negatively electrified bodies repel each other; indeed electrical attraction and repulsion generally must be explained before the validity of the theory can be assented to; but as both the received theories do this in a nearly similar manner to the present one, it need not now be entered upon. My object has been briefly to prove the sufficiency of one electric agent to elucidate what it has been considered possible to do only by two, and by so doing to furnish a basis for the explanation of the whole series of electrical phænomena. Statical electricity has alone furnished exercise for legitimate theory; and although the identity of it and voltaic electricity has never been doubted, the connexion between their effects has received no solution. The relation between the atoms of bodies and statical electrical excitement, here suggested, seems to furnish a clue by which dynamical electricity may become more intelligible. By dynamical electricity, I mean voltaic effects generally. This may perhaps more clearly appear by considering the influences to which the particles might be conceived to be subject. In order to this, we must have some standard by which we may compare different atoms or centres of matter; so we will assume an unit for that purpose. Not that an atom must consist of one unit only, but that different atoms may consist of different definite units.

Let the repulsion between one unit of matter and another be called  $R$ , the repulsion between units of electricity  $r$ , the attraction between units of electricity and matter  $a$ ; let the respective quantities of matter in two different atoms be  $M$  and  $m$ , and the respective quantities of electricity  $E$  and  $e$ .

Then the repulsion which these atoms exercise on each other will be represented by

$$M m R + E e r . . . . . (A.)$$

And the attraction will be

$$M e a + m E a . . . . . (B.)$$

Imagine a case of equilibrium, then

$$M m r + E e r = M e a + m E a.$$

Now all those quantities are variable; and it is easy to perceive that the attraction and repulsion will vary with varying values of either of the quantities. At present we wish to know the effect of increase or diminution of the quantity of electricity on an atom.

Now suppose the attraction and repulsion equal, and we get

$$M e a - M m R = E e r - E m a.$$

(1)
(2)

Diminish E then (2) becomes less than (1), and consequently B becomes less than A; that is, the repulsion becomes in excess. This is on the assumption of *ma* being greater than *er*. Increase E, and in like manner the attraction becomes in excess. In a similar manner might it be shown that an increase or diminution of *e* would cause a corresponding attraction or repulsion. Still more so, then, must this occur when both E and *e* increase or diminish together. This is an evident reason for the repulsion existing between bodies negatively electrified. But its more important feature is the view it would give of voltaic excitement; indicating that the current is simply the appropriation of the electricity holding the elements of the liquid compound in combination, to the formation of the new compound which is essentially always in process. It will not be necessary now to trace any further the effects of this mode of considering the constitution of matter. I have said thus much merely as a preliminary necessary for its reception. Whether similar views may have been entertained before, I know not, but they have been my own for a considerable period past, and in their elucidation of all the facts on which I have tried them, there seems to be greater consistency than on many other suppositions which I have met with. It may be worth while to allude to the manner in which the property of conduction is treated by this theory: it would appear that no body is a perfect conductor; the relation between conductors and non-conductors being merely a question of time and velocity; for each atom of a body exercising an attraction and repulsion on free electricity in it, the facility of transmission will depend on the ratio and intensity of those

forces, these again depending on the quantities of electricity and matter in the atoms. This has been merely hinted at, not for explanation, but to show that the theory gives the property referred to, to inherent powers in bodies themselves, and not to the space in which they are situated, and by which they are surrounded. My reasons for considering the atoms of bodies as mere centres of force, have not been given, as they are connected with other subjects on which you may not be able to afford space for entering.

I am, Sir,

Killigrew St., Falmouth,  
Jan. 19, 1846.

Yours, &c.,

HENRY SLOGGETT.

LXXV. *Experiments and Observations on the Mechanical Powers of Electro-Magnetism, Steam, and Horses.* By the Rev. WILLIAM SCORESBY, D.D., F.R.SS. L. and E., Corr. Memb. Inst. Fr., &c., and JAMES P. JOULE, Secretary of the Literary and Philosophical Society of Manchester, Mem. Chem. Soc., &c.\*

AT the last Meeting of the British Association, Dr. Scoresby described a magnetic apparatus of very great power, and gave an account of some experiments he had made with a view to test its capabilities for exciting electrical currents. The coils employed in those experiments were hastily constructed, and by no means calculated to produce a maximum effect. We agreed, therefore, to construct and try more efficient ones on the first opportunity.

Two kinds of revolving armature occurred to us as worthy of trial. One of them consisted of a hollow tube of drawn iron, 24 inches long,  $1\frac{5}{8}$ th inch in diameter, and  $\frac{3}{10}$ ths of an inch thick in the metal, bent into the shape of the letter U. It had a saw-cut along its entire length, in order to prevent the circulation of electrical currents in the substance of the iron. Each of the legs of this armature was wound with 274 feet of covered copper wire,  $\frac{1}{10}$ th of an inch in diameter. The other armature consisted of two bars of iron, each 20 inches long, 4 inches broad, and  $\frac{3}{8}$ ths of an inch thick. These bars were bent edgeways into the form of a semicircle, and then fastened together with the interposition of a piece of calico in order to prevent currents in the iron as much as possible. Each leg of this armature was furnished with two coils of covered copper wire  $\frac{1}{10}$ th of an inch thick. The two coils that were nearest the iron were each 276 feet long; and each of the other two coils was 296 feet long.

\* Communicated by the Authors.

Having placed the two straight steel magnets (each of which was 4 feet 4 inches long, 4 to 5 inches square, and had poles of  $7\frac{1}{2}$  square inches surface) side by side, in a horizontal position, and with two of their poles connected by a suitable armature, we placed the *hollow electro-magnetic armature* on the axis of a revolving apparatus, in such a position that the poles of the armature could revolve at the distance of about  $\frac{1}{4}$ th of an inch from the poles of the steel magnets. The coils were arranged for quantity, and connected by means of a proper "commutator" with platinum plates (each exposing an active surface of 5 or 6 square inches) immersed in a dilute solution of sulphuric acid. The maximum amount of decomposition was effected when the armature revolved 500 times per minute. At this velocity  $\frac{3}{4}$ ths of a cubic inch of the mixed gases were collected per minute.

Having removed the hollow armature, we now fastened the *flat semicircular armature* upon the axis. When this armature, with its four coils arranged for quantity, was rotated at the rate of 500 revolutions per minute, we collected as much as 1.4 cubic inch of the mixed gases per minute. With the same velocity of rotation, two inches of steel wire,  $\frac{1}{10}$ th of an inch thick, were raised to a bright red heat; and one inch of the same kind of wire was fused.

Great as the above effects undoubtedly are, in comparison with previously recorded results, we expect to be able to augment them very much by causing the armatures to revolve opposite the *true poles* of the magnets, and not, as heretofore, opposite their ends. It is proper also to observe, that on account of the imperfect hardness of many of the steel bars\*, the magnets did not possess one quarter of the power due to Dr. Scoresby's principle of construction. We have not, however, hitherto cared to reconstruct the apparatus, because our principal object in the present research was to make experiments with the machine working as an engine, for which purpose the magnets were quite powerful enough.

The battery employed for working the machine as an engine, consisted of three cells of Daniell's constant arrangement. In each cell the copper element exposed an active surface of two

\* The bars of which the magnetic apparatus was constructed were of various lengths, but of otherwise uniform dimensions, viz.  $1\frac{1}{2}$  inch broad and  $\frac{1}{4}$ th of an inch thick. The thickness and mass were found too great for effective hardening, at least for obtaining a degree of hardness capable of sustaining the severity of the magnetic test. Economy and facility of arrangement were the reasons for adopting this construction, rather than the more certain and effective one of *hard thin plates*, described by Dr. Scoresby in his "Magnetical Investigations."

square feet, and the amalgamated zinc plate a surface of  $\frac{2}{3}$  of a square foot. A pretty correct galvanometer, consisting of a circle of thick copper wire and a magnetic needle 3 inches long, was employed for measuring the currents of electricity which were transmitted by the battery through the revolving armatures. The tangents of the deflections of the magnetic needle, corrected by a small equation, indicated the absolute quantities of transmitted electricity. The quantity of zinc consumed in the battery was deduced from the deflections of the needle; the data of the calculation being derived from previous experiments on the quantity of mixed gases evolved from acidulated water by a current capable of producing a given deflection of the needle.

Our first experiments were made with the flat semicircular revolving armature, its four coils being arranged for quantity. The deflection of the needle before the engine was allowed to start amounted to  $64^\circ$ , which indicated a current of 2232, calling the current corresponding to  $45^\circ$ , 1000. The engine, being then allowed to start, presently attained a velocity of 140 revolutions per minute. The needle was then observed to stand steadily at  $43^\circ$ , indicating a current of 920. The consumption of zinc in the battery was estimated to be at the rate of 205 grs. per hour.

Although we were not able to apply as exact a dynamometer as we could have wished, we were nevertheless enabled to arrive at a pretty correct estimation of the power developed, by ascertaining the weight which, when thrown over a wheel connected with the engine, was sufficient to keep it in uniform motion. In this way we found that the force developed in the above experiment was equal to raise 21,100 lbs. to the height of a foot per hour.

On making a second experiment with the same revolving armature and battery, we obtained the following results:— Current before the engine was allowed to start, 2232; current when the armature was rotating at the rate of 180 revolutions per minute, 850; consumption of zinc per hour, 190 grains; force given out per hour, 17,820 lbs. raised a foot.

Mr. J. P. Joule has already proved that the heat evolved by voltaic and magneto-electrical currents is, *ceteris paribus*, proportional to the square of their intensity\*; and that the power of the electro-magnetic engine is obtained at the expense of the heat due to the chemical reactions of the voltaic battery by which it is worked. He has also shown, that if the whole of the heat developed by the consumption of a grain of zinc in a Daniell's battery could be converted into useful

\* Phil. Mag., vol. xviii. p. 308, and vol. xix. p. 260.



mechanical power, it would be equal to raise a weight of 158 lbs. to the height of a foot\*. Hence, if we designate the current when the engine is *at rest* by  $a$ , and the current when the engine is *in motion* by  $b$ , the heat evolved by the circuit in a given time, will, in the two instances, be as  $a^2$  to  $b^2$ . But the quantities of zinc consumed being as  $a$  to  $b$ , the heat, per a given consumption of zinc, will be as  $a$  to  $b$ , or directly as the currents;  $a - b$  will therefore represent the quantity of heat converted by the engine into useful mechanical effect. Therefore, putting  $x$  for the mechanical effect in lbs. raised a foot high per the consumption of a grain of zinc, we have

$$x = \frac{158(a - b)}{a}.$$

From the above equation it is evident that the economical duty will be a maximum when  $b$  vanishes or becomes infinitely small in comparison with  $a$ . In this case  $x = 158$ , while the power of the engine will become infinitely small with regard to work performed in a given time. We must, however, observe that the equation can only be strictly correct when the current  $b$  is uniform, which it never can be exactly, in consequence of the resistance of the magnetic induction against the voltaic current varying in the different positions of the revolving electro-magnetic armature. Hence the current  $b$  is always, to a certain extent, of a *pulsatory* character, which has the effect of causing it to develop more heat than an *uniform* current of the same quantity. From this circumstance, as well as from the unavoidable existence of some slight currents in the substance of the iron of the revolving armature, the actual economical effect will always be somewhat below the duty indicated by our formula.

Applying the formula to our first experiment, we have for the *theoretical* economical effect,

$$\frac{158(2232 - 920)}{2232} = 92.9,$$

while the *actual* economical effect was

$$\frac{21100}{205} = 102.9.$$

In our second experiment, the theoretical economical effect will be

$$\frac{158(2232 - 850)}{2232} = 97.8,$$

\* *Phil. Mag.*, vol. xxiii. p. 441.

and the actual duty,

$$\frac{17820}{190} = 93.8.$$

Taking the mean of the two experiments, we have for the theoretical duty 95.3, and for the actual performance 98.3. Here, therefore, in apparent contradiction to what we have just said, the actual exceeds the theoretical duty. This circumstance is however partly explained by the fact that the solution of sulphuric acid employed in charging the battery had been mixed immediately before the experiments were made, and was in consequence considerably heated; for Daniell has shown that the intensity of his battery increases with its temperature, and it is evident that an increase of the intensity or electromotive force of the cells of the battery must be productive of an increased economical effect.

The next two experiments were made with the hollow revolving armature, its two coils being arranged for quantity. In these and the subsequent experiments, the battery was charged with a *cold* solution.

*Experiment 3.*—Current when the engine was kept at rest, 1381; current when the armature was revolving 80 times per minute, 850; consumption of zinc, 190 grains per hour; power developed, 8800 lbs. raised a foot high per hour. From these data, the theoretical duty will be

$$\frac{158(1381 - 850)}{1381} = 60.7,$$

and the actual duty will be

$$\frac{8800}{190} = 46.3.$$

*Experiment 4.*—Current before the engine was allowed to start, 1381; current when the engine was revolving 102 times per minute, 678; consumption of zinc, 151 grains per hour; power developed, 9000 lbs. raised a foot per hour. Hence for the theoretical duty we have,

$$\frac{158(1381 - 678)}{1381} = 80.4,$$

and for the actual duty,

$$\frac{9000}{151} = 59.6.$$

Lastly, we made two experiments in which the engine was fitted up with two straight electro-magnets fastened parallel to the axis. Each of these straight electro-magnets consisted of a piece of drawn iron tube, 12 inches long,  $1\frac{5}{8}$ th inch in

diameter, and  $\frac{3}{16}$ ths of an inch thick, cut longitudinally to prevent the circulation of electrical currents in the iron, and furnished with a coil of 210 feet of covered copper wire  $\frac{1}{10}$ th of an inch thick. A steel magnet consisting of a considerable number of bars was fitted up in order to excite those ends of the straight electro-magnets which were distant from the large steel magnets. The coils were arranged for quantity.

*Experiment 5.*—Current when the engine was kept still, 2081; current when the armature was revolving 114 times per minute, 1300; consumption of zinc, 291 grains per hour; power developed, 10030 lbs. raised a foot per hour. Hence the theoretical duty will be

$$\frac{158 (2081 - 1300)}{2081} = 59.3,$$

and the actual duty,

$$\frac{10030}{291} = 34.5.$$

*Experiment 6.*—Current before starting, 2035; current when revolving 192 times per minute, 1000; consumption of zinc, 223 grains per hour; power developed, 12,672 lbs. raised a foot per hour. In this case the theoretical duty will be

$$\frac{158 (2035 - 1000)}{2035} = 80.3;$$

the actual performance will be

$$\frac{12672}{223} = 56.8.$$

The mean of the six experiments gives a theoretical duty of 78.5, and an actual duty of 65.6. But, making allowance for the hot solution employed in the first two experiments, we may state that the actual was in general about  $\frac{2}{3}$ ths of the theoretical duty.

Upon the whole we feel ourselves justified in fixing the maximum available duty of an electro-magnetic engine worked by a Daniell's battery at 80 lbs. raised a foot high for each grain of zinc consumed\*, or, in other words, at about half the theoretical maximum of duty.

Before we leave this part of the subject, we may state that the above experiments fully bear out the idea expressed by

\* Dr. Botto states that 45 lbs. of zinc consumed in a Grove's battery are sufficient to work a one-horse power electro-magnetic engine for 24 hours. The intensity of Daniell's battery being  $\frac{2}{3}$ ths of that of Grove, it follows that 75 lbs. of zinc would have been consumed had Dr. Botto employed a Daniell's battery,—a result not widely different from our own.

Dr. Scoresby in his "Magnetical Investigations," that steel magnets on his construction may be employed in the stationary part of the electro-magnetic engine with much greater advantage than electro-magnets. We have already adverted to the imperfect construction of the magnetic apparatus employed in the above experiments; had we employed one of equal weight, but constructed of thin plates of hardened steel, and furnished with armatures and batteries in proportion, we think it highly probable that a power equal to that of one horse might have been attained, the whole weight of the apparatus being considerably under half a ton.

Having thus determined the capabilities of electro-magnetism as a first mover of machinery, it will be interesting and instructive to compare it with two other sources of power, viz. steam and horses.

1. A grain of coal produces, by combustion, sufficient heat to raise the temperature of a lb. of water  $1^{\circ}\cdot634$ . In other words, we may say that the *vis viva* developed by the combustion of a grain of coal is equal to raise a weight of 1335 lbs. to the height of one foot. Now the best Cornish steam-engines raise 143 lbs. per grain of coal; whence it appears that the steam-engine in its most improved state is not able to develop much more than  $\frac{1}{10}$ th of the *vis viva* due to the combustion of coal into useful power, the remaining  $\frac{9}{10}$ ths being given off in the form of heat.

2. A horse, when its power is advantageously applied, is able to raise a weight of 24,000,000 lbs. to the height of one foot per day. In the same time (24 hours) he will consume 12 lbs. of hay and 12 lbs. of corn\*. He is therefore able to raise 143 lbs. by the consumption of one grain of the mixed food. From our own experiments on the combustion of a mixture of hay and corn in oxygen gas, we find that each grain of food, consisting of equal parts of undried hay and corn, is able to give  $0^{\circ}\cdot682$  to a lb. of water, a quantity of heat equivalent to the raising of a weight of 557 lbs. to the height of a foot. Whence it appears, that one quarter of the whole amount of *vis viva* generated by the combustion of food in the

\* We have been kindly informed by Mr. J. V. Gibson of Manchester, an eminent veterinary surgeon, that 14 lbs. of hay and 10 lbs. of corn is the average provender requisite to support a horse of average size, so as to enable him to work daily without any depreciation of his physical condition. We have however equalized the quantities of hay and corn, on account of the experiments on combustion having been made with a mixture containing equal portions.

animal frame, is capable of being applied in producing a useful mechanical effect,—the remaining three-quarters being required in order to keep up the animal heat, &c.

Prof. Magnus of Berlin, has endeavoured to prove that the oxygen which an animal inspires does not combine chemically with the blood, but is merely *absorbed* by it\*. The blood thus charged with oxygen arrives in the capillary vessels, where the oxygen effects a chemical combination with *certain substances*, converting them into carbonic acid and water. The carbonic acid, instead of oxygen, is then absorbed by the blood, and thus reaches the lungs to be removed by contact with the atmosphere. Adopting this view, it becomes exceedingly probable that the *whole* of the *vis viva* due to the oxidation or combustion of the “certain substances” mentioned by Magnus is developed by the muscles. The muscles, by their motion, can communicate *vis viva* to external objects; and, by their friction within the body, can develop heat in various quantities according to circumstances, so as to maintain the animal at a uniform temperature. If these theoretic views be correct, they would lead to the interesting conclusion (which is the same as that announced by Matteucci from other considerations) that the animal frame, though destined to fulfill so many other ends, is, as an engine, more perfect in the œconomy of *vis viva* than the best of human contrivances.

LXXVI. *Experimental Researches in Electricity*.—*Twentieth Series*. By MICHAEL FARADAY, Esq., D.C.L., F.R.S., *Fullerian Prof.*, &c. &c.

[Concluded from p. 406.]

¶ iv. *Action of magnets on metals generally.*

2287. **T**HE metals, as a class, stand amongst bodies having a high and distinct interest in relation both to magnetic and electric forces, and might at first well be expected to present some peculiar phænomena, in relation to the striking property found to be possessed in common by so large a number of substances, so varied in their general characters. As yet no distinction associated with conduction or non-conduction, transparent or opaque, solid or liquid, crystalline or amorphous, whole or broken, has presented itself; whether the metals, distinct as they are as a class, would fall into the great generalization, or whether at last a separation would occur, was to me a point of the highest interest.

2288. That the metals, iron, nickel and cobalt, would stand in a distinct class, appeared almost undoubted; and it will be,

\* [See *Phil. Mag.* S. 3. vol. xxvii. p. 561.]

I think, for the advantage of the inquiry, that I should consider them in a section apart by themselves. Further, if any other metals appeared to be magnetic, as these are, it would be right and expedient to include them in the same class.

2289. My first point, therefore, was to examine the metals for any indication of ordinary magnetism. Such an examination cannot be carried on by magnets anything short in power of those to be used in the further investigation; and in proof of this point I found many specimens of the metals, which appeared to be perfectly free from magnetism when in the presence of a magnetic needle, or a strong horse-shoe magnet (2157.), that yet gave abundant indications when suspended near to one or both poles of the magnets described (2246.).

2290 My test of magnetism was this. If a bar of the metal to be examined, about two inches long, was suspended (2249.) in the magnetic field, and being at first oblique to the axial line, was upon the supervention of the magnetic forces drawn into the axial position instead of being driven into the equatorial line, or remaining in some oblique direction, then I considered it magnetic. Or, if being near one magnetic pole, it was attracted by the pole, instead of being repelled, then I concluded it was magnetic. It is evident that the test is not strict, because, as before pointed out (2285.), a body may have a slight degree of magnetic force, and yet the power of the new property be so great as to neutralize or surpass it. In the first case, it might seem neither to have the one property nor the other; in the second case, it might appear free from magnetism, and possessing the special property in a *small* degree.

2291. I obtained the following metals, so that when examined as above, they did not appear to be magnetic; and in fact if magnetic, were so to an amount so small as not to destroy the results of the other force, or to stop the progress of the inquiry.

Antimony.	Lead.
Bismuth.	Mercury.
Cadmium.	Silver.
Copper.	Tin.
Gold.	Zinc.

2292. The following metals were, and are as yet to me, magnetic, and therefore companions of iron, nickel and cobalt:—

Platinum.	Titanium.
Palladium.	

2293. Whether all these metals are magnetic, in consequence of the presence of a little iron, nickel, or cobalt in them,

or whether any of them are really so of themselves, I do not undertake to decide at present; nor do I mean to say that the metals of the former list are free. I have been much struck by the apparent freedom from iron of almost all the specimens of zinc, copper, antimony and bismuth, which I have examined; and it appears to me very likely that some metals, as arsenic, &c., may have much power in quelling and suppressing the magnetic properties of any portion of iron in them, whilst other metals, as silver or platinum, may have little or no power in this respect.

2294. Resuming the consideration of the influence excited by the magnetic force over those metals which are not magnetic after the manner of iron (2291.), I may state that there are two sets of effects produced which require to be carefully distinguished. One of these depends upon induced magneto-electric currents, and shall be resumed hereafter (2309.). The other includes effects of the same nature as those produced with heavy glass and many other bodies (2276.).

2295. All the non-magnetic metals are subject to the magnetic power, and produce the same general effects as the large class of bodies already described. The force which they then manifest, they possess in different degrees. Antimony and bismuth show it well, and bismuth appears to be especially fitted for the purpose. It excels heavy glass, or borate of lead, and perhaps phosphorus; and a small bar or cylinder of it about two inches long, and from 0.25 to 0.5 of an inch in width, is as well fitted to show the various peculiar phenomena as anything I have yet submitted to examination.

2296. To speak accurately, the bismuth bar which I employed was two inches long, 0.33 of an inch wide, and 0.2 of an inch thick. When this bar was suspended in the magnetic field, between the two poles, and subject to the magnetic force, it pointed freely in the equatorial direction, as the heavy glass did (2253.), and if disturbed from that position returned *freely* to it. This latter point, though perfectly in accordance with the former phenomena, is in such striking contrast with the phenomena presented by copper and some other of the metals (2309.), as to require particular notice here.

2297. The comparative sensibility of bismuth causes several movements to take place under various circumstances, which being complicated in their nature, require careful analysis and explanation. The chief of these, with their causes, I will proceed to point out.

2298. If the cylinder electro-magnet (2246.) be placed vertically so as to present one pole upwards, that pole will exist in the upper end of an iron cylinder, having a flat horizontal

face  $2\frac{1}{2}$  inches in diameter. A small indicating sphere (2266.) of bismuth hung over the centre of this face and close to it, does not move by the magnetism. If the ball be carried outwards, half way, for instance, between the centre and the edge, the magnetism makes it move inwards, or towards the axis (prolonged) of the iron cylinder. If carried still further outwards, it still moves inwards under the influence of the magnetism, and such continues to be the case until it is placed just over the edge of the terminal face of the core, where it has no motion at all (here, by another arrangement of the experiment, it is known to tend in what is at present an upward direction from the core). If carried a little further outwards, the magnetism then makes the bismuth ball tend to go outwards or be repelled, and such continues to be the direction of the force in any further position, or down the side of the end of the core.

2299. In fact, the circular edge formed by the intersection of the end of the core with its sides, is virtually the apex of the magnetic pole, to a body placed like the bismuth ball close to it, and it is because the lines of magnetic force issuing from it diverge as it were, and weaken rapidly in all directions from it, that the ball also tends to pass in all directions either inwards or upwards, or outwards from it, and thus produces the motions described. These same effects do not in fact all occur when the ball, being taken to a greater distance from the iron, is placed in magnetic curves, having generally a simpler direction. In order to remove the effect of the edge, an iron cone was placed on the top of the core, converting the flat end into a cone, and then the indicating ball was urged to move upwards, only when over the apex of the cone, and upward and outwards, as it was more or less on one side of it, being always repelled from the pole in that direction, which transferred it most rapidly from strong to weaker points of magnetic force.

2300. To return to the vertical flat pole: when a horizontal bar of bismuth was suspended concentrically and close to the pole, it could take up a position in any direction relative to the axis of the pole, having at the same time a tendency to move upwards or be repelled from it. If its point of suspension was a little excentric, the bar gradually turned, until it was parallel to a line joining its point of suspension with the prolonged axis of the pole, and the centre of gravity moved inwards. When its point of suspension was just outside the edge of the flat circular terminating face, and the bar formed a certain angle with a radial line joining the axis of the core and the point of suspension, then the movements of the bar were un-



certain and wavering. If the angle with the radial line were less than that above, the bar would move into parallelism with the radius and go inwards: if the angle were greater, the bar would move until perpendicular to the radial line and go outwards. If the centre of the bar were still further out than in the last case, or down by the side of the core, the bar would always place itself perpendicular to the radius and go outwards. All these complications of motion are easily resolved into their simple elementary origin, if reference be had to the character of the circular angle bounding the end of the core; to the direction of the magnetic lines of force issuing from it and the other parts of the pole; to the position of the different parts of the bar in these lines; and the ruling principle that each particle tends to go by the nearest course from *strong* to *weaker* points of magnetic force.

2301. The bismuth points well, and is well repelled (2296.) when immersed in water, alcohol, æther, oil, mercury, &c., and also when inclosed within vessels of earth, glass, copper, lead, &c. (2272.), or when screens of 0·75 or 1 inch in thickness of bismuth, copper or lead intervene. Even when a bismuth cube (2266.) was placed in an iron vessel  $2\frac{1}{2}$  inches in diameter and 0·17 of an inch in thickness, it was well and freely repelled by the magnetic pole.

2302. Whether the bismuth be in one piece or in very fine powder, appears to make no difference in the character or in the degree of its magnetic property (2283.).

2303. I made many experiments with masses and bars of bismuth suspended, or otherwise circumstanced, to ascertain whether two pieces had any mutual action on each other, either of attraction or repulsion, whilst jointly under the influence of the magnetic forces, but I could not find any indication of such mutual action: they appeared to be perfectly indifferent one to another, each tending only to go from stronger to weaker points of magnetic power.

2304. Bismuth, in very fine powder, was sprinkled upon paper, laid over the horizontal circular termination of the vertical pole (2246.). If the paper were tapped, the magnet not being excited, nothing particular occurred; but if the magnetic power were on, then the powder retreated in both directions, inwards and outwards, from a circular line just over the edge of the core, leaving the circle clear, and at the same time showing the tendency of the particles of bismuth in all directions from that line (2299.).

2305. When the pole was terminated by a cone (2246.) and the magnet not in action, paper with bismuth powder sprinkled over it being drawn over the point of the cone, gave

no particular result; but when the magnetism was on, such an operation cleared the powder from every point which came over the cone, so that a mark was traced or written out in clear lines running through the powder, and showing every place where the pole had passed.

2306. The bar of bismuth and a bar of antimony was found to set equatorially between the poles of the ordinary horse-shoe magnet.

2307. The following list may serve to give an idea of the apparent order of some metals, as regards their power of producing these new effects, but I cannot be sure that they are perfectly free from the magnetic metals. In addition to that, there are certain other effects produced by the action of magnetism on metals (2309.) which greatly interfere with the results due to the present property.

Bismuth.	Cadmium.
Antimony.	Mercury.
Zinc.	Silver.
Tin.	Copper.

2308. I have a vague impression that the repulsion of bismuth by a magnet has been observed and published several years ago. If so, it will appear that what must then have been considered as a peculiar and isolated effect, was the consequence of a general property, which is now shown to belong to all matter\*.

2309. I now turn to the consideration of some peculiar phenomena which are presented by copper and several of the metals when they are subjected to the action of magnetic forces, and which so tend to mask effects of the kind already described, that if not known to the inquirer they would lead to much confusion and doubt. These I will first describe as to their appearances, and then proceed to consider their origin.

2310. If instead of a bar of bismuth (2296.) a bar of copper of the same size be suspended between the poles (2247.), and

\* M. de la Rive has this day referred me to the *Bibliothèque Universelle* for 1829, tome xl. p. 82, where it will be found that the experiment spoken of above is due to M. la Baillif of Paris. M. la Baillif showed sixteen years ago that both bismuth and antimony repelled the magnetic needle. It is astonishing that such an experiment has remained so long without further results. I rejoice that I am able to insert this reference before the present series of these researches goes to press. Those who read my papers will see here, as on many other occasions, the results of a memory which becomes continually weaker; I only hope that they will be excused, and that omissions and errors of that nature will be considered as involuntary.—M. F. December 30, 1845.

magnetic power be developed whilst the bar is in a position oblique to the axial and equatorial lines, the experimenter will perceive the bar to be affected, but this will not be manifest by any tendency of the bar to go to the equatorial line; on the contrary, it will advance towards the axial position as if it were magnetic. It will not however continue its course until in that position, but, unlike any effect produced by magnetism, will stop short, and making no vibration beyond or about a given point, will remain there coming at once to a dead rest: and this it will do even though the bar by the effect of torsion or momentum was previously moving with a force that would have caused it to make several gyrations. This effect is in striking contrast with that which occurs when antimony, bismuth, heavy glass, or other such bodies are employed, and it is equally removed from an ordinary magnetic effect.

2311. The position which the bar has taken up it retains with a considerable degree of tenacity, provided the magnetic force be continued. If pushed out of it, it does not return into it, but takes up its new position in the same manner, and holds it with the same stiffness; a push however, which would make the bar spin round several times if no magnetism were present, will now not move it through more than  $20^{\circ}$  or  $30^{\circ}$ . This is not the case with bismuth or heavy glass; they vibrate freely in the magnetic field, and always return to the equatorial position.

2312. The position taken up by the bar may be any position. The bar is moved a little at the instant of superinducing the magnetism, but allowing and providing for that, it may be finally fixed in any position required. Even when swinging with considerable power by torsion or momentum, it may be caught and retained in any place the experimenter wishes.

2313. There are two positions in which the bar may be placed at the beginning of the experiment, from which the magnetism does not move it, the equatorial and the axial positions. When the bar is nearly midway between these, it is usually most strongly affected by the first action of the magnet, but the position of most effect varies with the form and dimensions of the magnetic poles and of the bar.

2314. If the centre of suspension of the bar be in the axial line, but near to one of the poles, these movements occur well, and are clear and distinct in their direction: if it be in the equatorial line, but on one side of the axial line, they are modified, but in a manner which will easily be understood hereafter.

2315. Having thus stated the effect of the supervention of

the magnetic force, let us now remark what occurs at the moment of its cessation; for during its continuance there is no change. If, then, after the magnetism has been sustained for two or three seconds, the electric current be stopped, there is instantly a strong action on the bar, which has the appearance of a revulsion (for the bar returns upon the course which it took for a moment when the electric contact was made), but with such force, that whereas the advance might be perhaps  $15^\circ$  or  $20^\circ$ , the revulsion will cause the bar occasionally to move through two or three revolutions.

2316. Heavy glass or bismuth presents no such phenomena as this.

2317. If, whilst the bar is revolving from revulsion the electric current at the magnet be renewed, the bar instantly stops with the former appearances and results (2310.), and then upon removing the magnetic force is affected again, and, of course, now in a contrary direction to the former revulsion.

2318. When the bar is caught by the magnetic force in the axial or equatorial position, there is no revulsion. When inclined to these positions there is; and the places most powerful in this respect appear to be those most favourable to the first brief advance (2313.). If the bar be in a position at which strong revulsion would occur, and whilst the magnetism is continued be moved by hand into the equatorial or axial position, then on taking off the magnetic force there is no revulsion.

2319. If the continuance of the electric current and consequently of the magnetism be for a moment only, the revulsion is very little, and the shorter the continuance of the magnetic force the less is the revulsion. If the magnetic force be continued for two or three seconds and then interrupted and *instantly* renewed, the bar is loosened and caught again by the power before it sensibly changes its place; and now it may be observed that it does not advance on the *renewal* of the force as it would have done had it been acted on by a first contact in that place (2310.); *i. e.* if the bar be in a certain place inclined to the axial position, the first supervention of the magnetic power causes it to advance towards the axial position; but the bar being in the same place and the magnetic power suspended and *instantly* renewed, the second supervention of force does not move the bar as the first did.

2320. When the copper bar is immersed in water, alcohol, or even mercury, the same effects take place as in the air, but the movements are, of course, not to the same extent.

2321. When plates of copper or bismuth, an inch in thick-

ness, intervene between the poles and the copper bar, the same results occur.

2322. If one magnetic pole only be employed the effects occur near it as well as before, provided that pole have a face large in proportion to the bar, as the end of the iron core (2246.): but if the pole be pointed by the use of the conical termination, or if the bar be opposite the edge of the end of the core, then they become greatly enfeebled or disappear altogether; and only the general fact of repulsion remains (2295.).

2323. The peculiar effects which have just been described are perhaps more strikingly shown if the bar of copper be suspended perpendicularly, and then hung opposite and near to the large face of a single magnetic pole, or the pole being placed vertically, as described (2246. 2263.), anywhere near to its side. The bar, it will be remembered, is two inches in length by 0.33 of an inch in width, and 0.2 of an inch in thickness, and as it now will revolve on an axis parallel to its length, the two smaller dimensions are those which are free to move into new positions. In this case the establishment of the magnetic force causes the bar to turn a little in accordance with the effects before described, and the removal of the magnetic force causes a revulsion, which sends the bar spinning round on its axis several times. But at any moment the bar can again be caught and held in a position as before. The tendency on making contact at the battery is to place the longest moving dimension, *i. e.* the width of the bar, parallel to the line joining the centre of action of the magnet and the bar.

2324. The bar, as before (2311.), is extremely sluggish and as if immersed in a dense fluid, as respects rotation on its own axis; but this sluggishness does not affect the bar as a whole, for any pendulum vibration it has continues unaffected. It is very curious to see the bar, jointly vibrating from its point of suspension (2249.) and rotating on its axis, when first affected by the magnetic force, for instantly the latter motion ceases, but the former goes on with undiminished power.

2325. The same effect of sluggishness occurs with a cube or a globe of copper as with the bar, but the phænomena of the first turn and the revulsion cease (2310. 2315.).

2326. The bars of bismuth and heavy glass present no appearance of this kind. The peculiar phænomena produced by copper are as distinct from the actions of these substances as they are from ordinary magnetic actions.

2327. Endeavouring to explain the cause of these effects, it appears to me that they depend upon the excellent con-

ducting power of copper for electric currents, the *gradual* acquisition and loss of magnetic power by the iron core of the electro-magnet, and the production of those induced currents of magneto-electricity which I described in the First Series of these Experimental Researches (55. 109.).

2328. The obstruction to motion on its own axis, when the bar is subjected to the magnetic forces, belongs equally to the form of a sphere or a cube. It belongs to these bodies, however, only when their axes of rotation are perpendicular or oblique to the lines of magnetic force, and not when they are parallel to it; for the horizontal bar, or the vertical bar, or the cube or sphere, rotate with perfect facility when they are suspended *above* the vertical pole (2246.), the rotation and vibration being then equally free, and the same as the corresponding movements of bismuth or heavy glass. The obstruction is at a maximum when the axis of rotation is perpendicular to the lines of magnetic force, and when the bar or cube, &c. is near to the magnet.

2329. Without going much into the particular circumstances, I may say that the effect is fully explained by the electric currents induced in the copper mass. By reference to the Second Series of these Researches (160.), it will be seen that when a globe, subject to the action of lines of magnetic force, is revolving on an axis perpendicular to these lines, an electric current runs round it in a plane parallel to the axis of rotation and to the magnetic lines, producing consequently a magnetic axis in the globe, at right angles to the magnetic curves of the inducing magnet. The magnetic poles of this axis therefore are in that direction which, in conjunction with the chief magnetic pole, tends to draw the globe back against the direction in which it is revolving. Thus, if a piece of copper be revolving before a north magnetic pole, so that the parts nearest the pole move towards the right-hand, then the right-hand side of that copper will have a south magnetic state, and the left-hand side a north magnetic state; and these states will tend to counteract the motion of the copper towards the right-hand: or if it revolve in the contrary direction, then the right-hand side will have a south magnetic state, and the left-hand side a north magnetic state. Whichever way, therefore, the copper tends to revolve on its own axis, the instant it moves, a power is evolved in such a direction as tends to stop its motion and bring it to rest. Being at rest in reference to this direction of motion, then there is no residual or other effect which tends to disturb it, and it remains still.

2330. If the whole mass be moving parallel to itself, and be small in comparison with the face of the magnetic pole opposite to which it is placed, then, though it pass through the magnetic lines of force, and consequently have a tendency to the formation of magneto-electric currents within it, yet as all parts move with equal velocity and in the same direction through similar magnetic lines of force, the tendency to the formation of a current is the same in every part, and there is no actual production of current, and consequently nothing occurs which can in any way interfere with its freedom of motion. Hence the reason that though the rotation of the bar or cube (2324. 2328.) upon its own axis is stopped, its vibration as a pendulum is not affected.

2331. That neither the one nor the other motion is affected when the bar or cube is over the vertical pole (2328.), is simply because in both cases (with the given dimensions of the pole and the moving metal) the lines of particles through which the induced currents tend to move are parallel throughout the whole mass; and therefore, as there is no part by which the return of the current can be carried on, no current can be formed.

2332. Before proceeding to the explanation of the other phenomena, it will be necessary to point out the fact generally understood and acknowledged, I believe, that time is required for the development of magnetism in an iron core by a current of electricity; and also for its fall back again when the current is stopped. One effect of the gradual rise in power was referred to in the last series of these Researches (2170.). This time is probably longer with iron not well annealed than with very good and perfectly annealed iron. The last portions of magnetism which a given current can develop in a certain core of iron, are also apparently acquired more slowly than the first portions; and these portions (or the condition of iron to which they are due) also appear to be lost more slowly than the other portions of the power. If electric contact be made for an instant only, the magnetism developed by the current disappears as instantly on the breaking of the current, as it appeared on its formation; but if contact be continued for three or four seconds, breaking the contact is by no means accompanied by a disappearance of the magnetism with equal rapidity.

2333. In order to trace the peculiar effect of the copper, and its cause, let us consider the condition of the horizontal bar (2310. 2313.) when in the equatorial position, between the two magnetic poles, or before a single pole; the point of suspension being in a line with the axis of the pole and its ex-

citing wire helix. On sending an electric current through the helix, both it and the magnet it produces will conduce to the formation of currents in the copper bar in the contrary direction. This is shown from my former researches (26.), and may be proved, by placing a small or large wire helix-shaped (if it be desired) in the form of the bar, and carrying away the currents produced in it, by wires to a galvanometer at a distance. Such currents being produced in the copper, only continue whilst the magnetism of the core is rising and then cease (18. 39.), but *whilst* they continue, they give a virtual magnetic polarity to that face of the copper bar which is opposite to a certain pole, the polarity being the same in kind as the pole it faces. Thus on the side of the bar facing the north pole of the magnet, a north polarity will be developed; and on that side facing the south pole, a south polarity will be generated.

2334. It is easy to see that if the copper during this time were opposite only one pole, or being between two poles, were nearer to one than the other, this effect would cause its repulsion. Still, it cannot account for the whole amount of the repulsion observed alike with copper as with bismuth (2295.), because the currents are of but momentary duration, and the repulsion due to them would cease with them. They do, however, cause a brief repulsive effort, to which is chiefly due the first part of the peculiar effect.

2335. For if the copper bar, instead of being parallel to the face of the magnetic pole, and therefore at right angles to the resultant of magnetic force, be inclined, forming, for instance, an angle of  $45^{\circ}$  with the face, then the induced currents will move generally in a plane corresponding more or less to that angle, nearly as they do in the examining helix (2333.), if it be inclined in the same manner. This throws the polar axis of the bar of copper on one side, so that the north polarity is not directly opposed to the north pole of the inducing magnet, and hence the action both of this and the other magnetic pole upon the two polarities of the copper will be to send it further round, or to place it edgewise to the poles, or with its breadth parallel to the magnetic resultant passing through it (2323.): the bar therefore receives an impulse, and the angle of it nearest to the magnet appears to be pulled up towards the magnet. This action of course stops the instant the magnetism of the helix core ceases to rise, and then the motion due to this cause ceases, and the copper is simply subject to the action before described (2295.). At the same time that this twist or small portion of a turn round the point of suspension occurs, the centre of gravity of the whole mass is re-



pelled, and thus I believe all the actions up to this condition of things is accounted for.

2336. Then comes the *revulsion* which occurs upon the cessation of the electric current, and the falling of the magnetism in the core. According to the law of magneto-electric induction, the disappearance of the magnetic force will induce brief currents in the copper bar (28.), but in the contrary direction to those induced in the first instance; and therefore the virtual magnetic pole belonging to the copper for the moment, which is nearest the north end of the electromagnet, will be a south pole; and that which is furthest from the same pole of the magnet will be a north pole. Hence will arise an exertion of force on the bar tending to turn it round its centre of suspension in the contrary direction to that which occurred before, and hence the apparent revulsion; for the angle nearest the magnetic pole will recede from it, the broad face (2323.) or length (2315.) of the bar will come round and face towards the magnet, and an action the reverse in every respect of the first action will take place, except that whereas the motion was then only a few degrees, now it may extend to two or three revolutions.

2337. The cause of this difference is very obvious. In the first instance, the bar of copper was moving under influences powerfully tending to retard and stop it (2329.); in the second case these influences are gone, and the bar revolves freely with a force proportionate to the power exerted by the magnet upon the currents induced by its own action.

2338. Even when the copper is of such form as not to give the oblique resultant of magnetic action from the currents induced in it, when, for instance, it is a cube or a sphere, still the effect of the action described above is evident (2325.). When a plate of copper about three-fourths of an inch in thickness, and weighing two pounds, was sustained upon some loose blocks of wood and placed about 0.1 of an inch from the face of the magnetic pole, it was repelled and held off a certain distance upon the making and continuing of electric contact at the battery; and when the battery current was stopped, it returned towards the pole; but the return was much more powerful than that due to gravity alone (as was ascertained by an experiment), the plate being at that moment actually *attracted*, as well as tending by gravitation towards the magnet, so that it gave a strong tap against it.

2339. Such is, I believe, the explanation of the peculiar phenomena presented by copper in the magnetic field; and the reason why they appear with this metal and not with bismuth or heavy glass, is almost certainly to be found in its high

electro-conducting power, which permits the formation of currents in it by inductive forces, that cannot produce the same in a corresponding degree in bismuth, and of course not at all in heavy glass.

2340. Any ordinary magnetism due to metals by virtue of their inherent power, or the presence of small portions of the magnetic metals in them, must oppose the development of the results I have been describing: and hence metals not of absolute purity cannot be compared with each other in this respect. I have, nevertheless, observed the same phænomena in other metals; and as far as regards the sluggishness of rotatory motion, traced it even into bismuth. The following are the metals which have presented the phænomena in a greater or smaller degree:—

Copper.	Mercury.
Silver.	Platinum.
Gold.	Palladium.
Zinc.	Lead.
Cadmium.	Antimony.
Tin.	Bismuth.

2341. The accordance of these phænomena with the beautiful discovery of Arago\*, with the results of the experiments of Herschel and Babbage†, and with my own former inquiries (81.)‡, are very evident. Whether the effect obtained by Ampère, with his copper cylinder and a helix§, was of this nature, I cannot judge, inasmuch as the circumstances of the experiment and the energy of the apparatus are not sufficiently stated; but it probably may have been.

2342. As, because of other duties, three or four weeks may elapse before I shall be able to complete the verification of certain experiments and conclusions, I submit at once these results to the attention of the Royal Society, and will shortly embody the account of the action of magnets on magnetic metals, their action on gases and vapours, and the general considerations in another series of these Researches.

Royal Institution, Nov. 27, 1845.

\* *Annales de Chimie*, xxvii. 363; xxviii. 325; xxxii. 213. I am very glad to refer here to the *Comptes Rendus* of June 9, 1845, where it appears that it was M. Arago who first obtained his peculiar results by the use of electro- as well as common magnets.

† *Philosophical Transactions*, 1825, p. 467.

‡ *Ibid.* 1832. p. 146.

§ *Bibliothèque Universelle*, xxi. p. 48.

LXXVII. *On the Equations applying to Light under the action of Magnetism.* By G. B. AIRY, Esq., *Astronomer Royal.*

*To the Editors of the Philosophical Magazine and Journal.*

GENTLEMEN,

**B**Y the indulgence of Dr. Faraday, I have been able to observe in the most satisfactory way the phænomena of the rotation of the plane of polarization of light passing through boracic glass and other media under the action of magnetic currents passing nearly in the direction of the light. And in particular I have verified the very remarkable fact that, upon passing the light successively in opposite directions while the magnetic adjustments remain the same, the plane of polarization undergoes the same change of position in regard to space, or undergoes opposite changes of position in regard to the expression of "rotation to the right," or "rotation to the left," as referred to the eye of the observer.

On reflecting upon the important fact that this change is not produced except there be an intermediate diaphanous body, it seems impossible not to conceive that the effect on the light is produced mediately by the action of the magnetic forces on the diaphanous body. The object of this communication is to point out what, as I conceive, must be the form of the mathematical equations existing among the movements of the particles of the glass, &c. or its contained æther, in order to explain the phænomena on mechanical laws.

In order to justify my intruding upon you with a suggestion which is exceedingly imperfect, I think it right to state to you my opinion upon the present condition of the optical theory, and upon several steps which, though leading to nothing conclusive, have nevertheless contributed to the real intellectual progress of the science.

On the truth of the undulatory theory, as regards the geometrical representation of light by undulations based upon transversal vibrations, the resolution of which into vibrations at right angles to each other constitutes polarization, I have not the shadow of a doubt. These undulations, whatever may be the way in which they may have been originally created, I conceive to be propagated by mechanical laws applying to the attractive or repulsive forces of the particles of the medium, the assumed æther, or the medium and the æther combined. But I have seen no mechanical theory to which I attach much importance or any unqualified belief. Nevertheless I think that the investigation and publication of these mechanical theories have been advantageous to the science,

by showing that mechanical laws *may be able* to explain effects never before ascribed to mechanical laws. As regards the progress of intellect, it has been very important to show that variation of velocities, as depending on the period of the oscillations, is mechanically possible; it has been very important to show that transversal vibrations are mechanically possible; it has been very important to show that crystalline separation of differently polarized rays is mechanically possible. It is not that I believe completely in any one of the mechanical explanations which have been given, but that *à priori* difficulties have been removed, and that it may now be considered that there is a fair chance of reducing the whole to mechanical explanation.

In some cases the mechanical theory has stopped at the first step, as for instance in the very remarkable equations indicated by Prof. MacCullagh as competent to represent some of the characteristic phænomena of quartz. It was here an important matter to show that there was opened even a possibility of reducing these anomalous facts to mechanical laws.

The suggestion, which it is the object of this paper to lay before you, is of the same kind as that made by Professor MacCullagh.

In order to reduce the rotation of the plane of polarization to laws, I shall follow the example of Fresnel in assuming that plane-polarized light may be considered as compounded of two beams of circularly-polarized light, one right-handed and the other left-handed, and that the rotation of the plane is produced by a difference of the velocities of the two circularly-polarized beams. And this, I take this opportunity to observe, is actually the simplest way of conceiving the change, at least in instances like that of quartz, &c., and like that before us, when the same change is produced whatever be the position of the plane of polarization (a fact which, at my request, Dr. Faraday has very carefully verified). Although the conception of a plane vibration is easier where the plane of vibration has immediate reference to the plane of reflexion, &c., yet the conception of two circular vibrations is easier where the plane of the compound vibration has no reference to any plane in the apparatus, and is in fact perfectly arbitrary.

Now let  $x_1$  be measured in the direction in which the light is supposed to travel in the first experiment;  $x_2$  in the opposite direction, or in the direction in which the light will travel when, the magnetic adjustments remaining the same, the relative positions of the polariser and analyser are reversed; suppose these to be horizontal: let  $y_1$  be measured horizon-

tally towards the right as regards the course of the light in the first experiment,  $y_2$  towards the right as regards the course of the light in the second experiment (or opposite to  $y_1$ );  $z$  vertical, in a direction common to both experiments. Then, for the first experiment, in order to represent the displacement of particles constituting that ray of circularly polarized light, in which every particle describes a circle in the direction, viewed from the origin of light, opposite to that of the hands of a watch, and in which at any one time the position of all the particles, originally in a straight line, has become a right-handed helix (which I will call Ray No. I.); we must take the following expressions:—where  $\tau$  is the period of vibration,  $v'_1$  the velocity of transmission of the wave, and  $Y'_1$  and  $Z'_1$  the displacements in the direction of  $y_1$  and  $z$  respectively,

$$Y'_1 = a \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x_1}{v'_1} \right),$$

$$Z'_1 = a \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x_1}{v'_1} \right).$$

Similarly, to represent, for the first experiment, the displacement of particles constituting the ray circularly polarized in the opposite direction, or so that each particle describes a circle in the same direction, viewed from the origin of light, as the hands of a watch (which I shall call Ray No. II.), we must have

$$Y''_1 = b \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x_1}{v''_1} \right),$$

$$Z''_1 = -b \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x_1}{v''_1} \right).$$

And in the second experiment, to represent the Ray No. I. of that experiment, we must combine

$$Y'_2 = a \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x_2}{v'_2} \right),$$

$$Z'_2 = a \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x_2}{v'_2} \right);$$

and to represent the Ray No. II., we must combine

$$Y''_2 = b \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x_2}{v''_2} \right),$$

$$Z''_2 = -b \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x_2}{v''_2} \right).$$

And the thing which it is very important to observe is, that

the same mechanical equations referred to the same directions in absolute space must apply to all these displacements.

In ordinary crystals or fluids possessing the property of causing rotation of the plane of polarization in the same direction as referred to the eye of the observer, whether the ray be incident on one side or on the other, mechanical equations are to be sought which will produce the result, that in both cases the velocity of Ray No. I. is greater than that of Ray No. II. (or *vice versâ*); so that if  $v'_1$  is greater than  $v''_1$ ,  $v'_2$  will also be greater than  $v''_2$ . But in the glass affected by magnetism, if in the first experiment the velocity of Ray No. I. is greater than that of Ray No. II., then in the second experiment the velocity of Ray No. I. must be less than that of Ray No. II.; or if  $v'_1$  is greater than  $v''_1$ ,  $v'_2$  must be less than  $v''_2$ .

Now the equation which is deduced from every mechanical supposition that accounts for the propagation of undulations, is of the form

$$\frac{d^2 Y}{dt^2} = A \cdot \frac{d^2 Y}{dx^2},$$

$$\frac{d^2 Z}{dt^2} = A \cdot \frac{d^2 Z}{dx^2}.$$

And it seems probable that these equations, with the addition to each of a small term, may explain the difference of velocities of the Rays No. I. and No. II.

It was pointed out by Prof. MacCullagh, that the equations

$$\frac{d^2 Y}{dt^2} = A \cdot \frac{d^2 Y}{dx^2} + B \cdot \frac{d^3 Z}{dx^3},$$

$$\frac{d^2 Z}{dt^2} = A \cdot \frac{d^2 Z}{dx^2} - B \cdot \frac{d^3 Y}{dx^3}$$

would explain this difference. I may remark here, that in the last term of the second side of each equation, any differential coefficient of an odd order would have sufficed to explain the general fact of difference of velocity; but the third order was adopted by Prof. MacCullagh in order to reconcile the expression for difference of velocity in differently-coloured rays with the fact established by experiment.

It is however necessary to inquire whether, if this assumption makes  $v'_1$  greater than  $v''_1$ , it will make  $v'_2$  greater than  $v''_2$ . For this purpose we must convert the various expressions into expressions referred to the same co-ordinates.

Let  $x_1 = x$ ,  $x_2 = -x$ ;  $y_1 = y$ ,  $y_2 = -y$ ;  
in the first experiment let

$$Y'_1 = Y', \quad Y''_1 = Y''; \quad Z'_1 = Z', \quad Z''_1 = Z'':$$

in the second experiment let

$$Y'_2 = -Y', \quad Y''_2 = -Y'', \quad Z'_2 = Z', \quad Z''_2 = Z''.$$

Then,

in the first experiment, for Ray No. I.,

$$Y' = a \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v'_1} \right),$$

$$Z' = a \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v'_1} \right);$$

and Prof. MacCullagh's equations become

$$-\frac{4\pi^2}{\tau^2} a \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v'_1} \right) = -A \frac{4\pi^2}{\tau^2} \left( \frac{1}{v'_1} \right)^2 a \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v'_1} \right)$$

$$+ B \frac{8\pi^3}{\tau^3} \left( \frac{1}{v'_1} \right)^3 a \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v'_1} \right),$$

$$-\frac{4\pi^2}{\tau^2} a \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v'_1} \right) = -A \cdot \frac{4\pi^2}{\tau^2} \cdot \left( \frac{1}{v'_1} \right)^2 a \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v'_1} \right)$$

$$+ B \cdot \frac{8\pi^3}{\tau^3} \cdot \left( \frac{1}{v'_1} \right)^3 a \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v'_1} \right),$$

which agree in giving

$$(v'_1)^2 = \frac{A}{1 + B \frac{2\pi}{\tau} \left( \frac{1}{v'_1} \right)^3}.$$

For Ray No. II.,

$$Y'' = b \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v''_1} \right),$$

$$Z'' = -b \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v''_1} \right).$$

The equations become

$$-\frac{4\pi^2}{\tau^2} b \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v''_1} \right) = -A \cdot \frac{4\pi^2}{\tau^2} \left( \frac{1}{v''_1} \right)^2 b \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v''_1} \right)$$

$$- B \cdot \frac{8\pi^3}{\tau^3} \left( \frac{1}{v''_1} \right)^3 b \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v''_1} \right),$$

$$+\frac{4\pi^2}{\tau^2} b \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v''_1} \right) = +A \cdot \frac{4\pi^2}{\tau^2} \left( \frac{1}{v''_1} \right)^2 b \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v''_1} \right)$$

$$+ B \cdot \frac{8\pi^3}{\tau^3} \left( \frac{1}{v''_1} \right)^3 b \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v''_1} \right),$$

which agree in giving

$$(v''_1)^2 = \frac{A}{1 - B \frac{2\pi}{\tau} \left(\frac{1}{v''_1}\right)^3}.$$

Hence  $v'_1$  is less than  $v''_1$ .

In the second experiment, for Ray No. I.,

$$Y' = -a \cdot \cos \frac{2\pi}{\tau} \left(t + \frac{x}{v'_2}\right),$$

$$Z' = a \cdot \sin \frac{2\pi}{\tau} \left(t + \frac{x}{v'_2}\right).$$

The equations become

$$+ \frac{4\pi^2}{\tau^2} a \cdot \cos \frac{2\pi}{\tau} \left(t + \frac{x}{v'_2}\right) = + A \frac{4\pi^2}{\tau^2} \cdot \left(\frac{1}{v'_2}\right)^2 a \cdot \cos \frac{2\pi}{\tau} \left(t + \frac{x}{v'_2}\right)$$

$$- B \cdot \frac{8\pi^3}{\tau^3} \cdot \left(\frac{1}{v'_2}\right)^3 a \cdot \cos \frac{2\pi}{\tau} \left(t + \frac{x}{v'_2}\right),$$

$$- \frac{4\pi^2}{\tau^2} a \cdot \sin \frac{2\pi}{\tau} \left(t + \frac{x}{v'_2}\right) = - A \cdot \frac{4\pi^2}{\tau^2} \left(\frac{1}{v'_2}\right)^2 a \cdot \sin \frac{2\pi}{\tau} \left(t + \frac{x}{v'_2}\right)$$

$$+ B \cdot \frac{8\pi^3}{\tau^3} \cdot \left(\frac{1}{v'_2}\right)^3 a \cdot \sin \frac{2\pi}{\tau} \left(t + \frac{x}{v'_2}\right),$$

which agree in giving

$$(v'_2)^2 = \frac{A}{1 + B \frac{2\pi}{\tau} \left(\frac{1}{v'_2}\right)^3}.$$

And similarly, for Ray No. II.,

$$(v''_2)^2 = \frac{A}{1 - B \frac{2\pi}{\tau} \left(\frac{1}{v''_2}\right)^3}.$$

Hence  $v'_2$  is less than  $v''_2$ .

Thus in both experiments (that is, whether the light passes from one side or from the other side) the Ray No. II. travels more quickly than the Ray No. I. And therefore, if in each experiment there is incident a plane-polarized ray, consisting of the combination of a Ray No. I. and a Ray No. II., the plane-polarized ray which is formed by their union after emergence will have its plane of polarization turned from the original plane of polarization, in both experiments in the same direction as the hands of a watch, or in both experiments in the direction opposite to that of the hands of a watch, as referred to the eye of a person looking in the direction of the path of the light.

This result agrees with the phænomena of quartz, turpentine, &c.; and therefore Prof. MacCullagh's equations apply



to the explanation of crystalline rotation of the plane of polarization. But it does not agree with the phænomena of glass, &c. under magnetic action; and for this case new equations must be sought.

The equations which I offer as competent to represent this case are,

$$\frac{d^2 Y}{dt^2} = A \cdot \frac{d^2 Y}{dx^2} + C \cdot \frac{dZ}{dt},$$

$$\frac{d^2 Z}{dt^2} = A \cdot \frac{d^2 Z}{dx^2} - C \cdot \frac{dY}{dt},$$

which are to be verified in the same manner as those applying to the phænomena of quartz, &c.

Thus, in the first experiment, for Ray No. I.,

$$Y^I = a \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v_1} \right),$$

$$Z^I = a \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v_1} \right).$$

The equations become

$$-\frac{4\pi^2}{\tau^2} \cdot a \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v_1} \right) = -A \frac{4\pi^2}{\tau^2} \left( \frac{1}{v_1} \right)^2 a \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v_1} \right) + C \frac{2\pi}{\tau} \cdot a \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v_1} \right),$$

$$-\frac{4\pi^2}{\tau^2} a \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v_1} \right) = -A \frac{4\pi^2}{\tau^2} \left( \frac{1}{v_1} \right)^2 a \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v_1} \right) + C \cdot \frac{2\pi}{\tau} \cdot a \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v_1} \right),$$

which agree in giving

$$(v_1)^2 = \frac{A}{1 + \frac{\tau}{2\pi} C}.$$

For Ray No. II.,

$$Y^{II} = b \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v_1} \right),$$

$$Z^{II} = -b \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v_1} \right);$$

and the equations become

$$-\frac{4\pi^2}{\tau^2} \cdot b \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v_1} \right) = -A \frac{4\pi^2}{\tau^2} \left( \frac{1}{v_1} \right)^2 b \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v_1} \right) - C \frac{2\pi}{\tau} b \cdot \cos \frac{2\pi}{\tau} \left( t - \frac{x}{v_1} \right),$$

$$\begin{aligned}
 + \frac{4\pi^2}{\tau^2} b \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v'_1} \right) &= + A \frac{4\pi^2}{\tau^2} \left( \frac{1}{v'_1} \right)^2 b \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v'_1} \right) \\
 + C \frac{2\pi}{\tau} b \cdot \sin \frac{2\pi}{\tau} \left( t - \frac{x}{v'_1} \right), &
 \end{aligned}$$

which agree in giving

$$(v''_1)^2 = \frac{A}{1 - \frac{\tau}{2\pi} C}$$

Hence  $v'_1$  is less than  $v''_1$ .

In the second experiment, for Ray No. I.,

$$Y'' = -a \cdot \cos \frac{2\pi}{\tau} \left( t + \frac{x}{v'_2} \right),$$

$$Z'' = a \cdot \sin \frac{2\pi}{\tau} \left( t + \frac{x}{v'_2} \right);$$

and the equations become

$$+ \frac{4\pi^2}{\tau^2} a \cdot \cos \frac{2\pi}{\tau} \left( t + \frac{x}{v'_2} \right) = + A \frac{4\pi^2}{\tau^2} \left( \frac{1}{v'_2} \right)^2 a \cdot \cos \frac{2\pi}{\tau} \left( t + \frac{x}{v'_2} \right)$$

$$+ C \frac{2\pi}{\tau} a \cdot \cos \frac{2\pi}{\tau} \left( t + \frac{x}{v'_2} \right),$$

$$- \frac{4\pi^2}{\tau^2} a \cdot \sin \frac{2\pi}{\tau} \left( t + \frac{x}{v'_2} \right) = - A \frac{4\pi^2}{\tau^2} \left( \frac{1}{v'_2} \right)^2 a \cdot \sin \frac{2\pi}{\tau} \left( t + \frac{x}{v'_2} \right)$$

$$- C \frac{2\pi}{\tau} a \cdot \sin \frac{2\pi}{\tau} \left( t + \frac{x}{v'_2} \right),$$

which agree in giving

$$(v''_2)^2 = \frac{A}{1 - \frac{\tau}{2\pi} C}$$

Similarly,

$$(v''_2)^2 = \frac{A}{1 + \frac{\tau}{2\pi} C}$$

Hence  $v'_2$  is greater than  $v''_2$ .

Thus if in one experiment the Ray No. II. travels more quickly than the Ray No. I., in the other experiment the Ray No. II. travels more slowly than the Ray No. I. And therefore if in each experiment there is incident a plane-polarized ray consisting of the combination of a Ray No. I. and a Ray No. II., the plane-polarized ray which is formed by their

union after emergence will have its plane of polarization turned from the original plane of polarization, in one experiment in the same direction as the hands of a watch, and in the other experiment in the opposite direction, as referred to the eye of a person looking in the direction of the path of the light.

This result agrees with the phænomena of boracic glass, &c. under the action of magnetic forces.

Instead of making the second term on the right-hand side of the equation depend on  $\frac{dZ}{dt}$ , we might with equal success have adopted  $\frac{d^3Z}{dt^3}$ ,  $\frac{d^3Z}{dx^2 \cdot dt}$ , or any other differential coefficient of an odd order in which the number of differentiations with respect to  $t$  is odd. Different powers of  $\tau$  and  $v$  will be introduced by different selections. In order to determine which of these selections is best adapted to represent the phænomena, it will be necessary to determine the deviation of the plane of polarization for light of different colours.

If  $\frac{dZ}{dt}$  be adopted, the equations suggested by me will amount to this:—"The force upon any particle in the direction of one ordinate depends in part upon its velocity in the direction of the other ordinate." There is no insurmountable difficulty in conceiving that this may be true, although we have at present no mechanical reason *à priori* for believing that it is true.

To remove the possibility of misunderstanding, I will repeat that I offer these equations with the same intention with which Prof. MacCullagh's equations were offered; not as giving a mechanical explanation of the phænomena, but as showing that the phænomena may be explained by equations, which equations appear to be such as might possibly be deduced from some plausible mechanical assumption, although no such assumption has yet been made.

I am, Gentlemen,

Your obedient Servant,

Royal Observatory, Greenwich,  
May 7, 1846.

G. B. AIRY.

LXXVIII. *Letter to Henry Lord Brougham, F.R.S., &c., containing Remarks on certain Statements in his Lives of Black, Watt and Cavendish. By the Rev. WILLIAM VERNON HARCOURT, F.R.S. &c.*

[Continued from p. 131.]

THERE are few things more remarkable in scientific history than the manner in which Newton may be observed to have dealt with the conjectural part of philosophy. He never speaks of hypothetical speculation but in terms implying somewhat of disdain. And yet in all his works, from the announcement to the Royal Society of his first discoveries respecting light to the last revision of the *Optics and Principia*, an hypothesis of the highest generality holds a conspicuous place.

This apparent inconsistency is however easily explained: he doubtless was deeply impressed with the error into which his predecessor Descartes had fallen, in building a system of philosophy on superficial analogies and precarious conjectures, and looked with some dissatisfaction at the pretension of his cotemporary Hook to set aside the inductive analysis of light, on the faith of a conjectural standard of his own. With Newton the imagining hypotheses was but as child's-play compared with the labour and importance of those severe and sure processes, inductive and deductive, to which he had devoted all the efforts of his mind. He held cheap the exercise of that great faculty of imagination from which the inexhaustible riches of his philosophical invention flowed with spontaneous facility. But though he laid no stress on what he called his "guesses," no man's mind seems ever to have been more continually, as it were, *upon the guess*; and no one ever gave so eminent and instructive an example of steady persistence in that conjectural habit of mind. "To show," says Newton, "that I do not take *gravity* for an essential property of bodies, I have added one question concerning its cause, choosing to propose it by way of question because I am not yet satisfied about it for want of experiments\*." After having himself achieved by a vigorous induction the most extensive generalisation to which the human intellect has ever attained, he still saw, in a stronger light than any one, reasons for doubting whether the law at which he had arrived was so simple and conformable to the rest of nature as to preclude our tracing it to some more general cause. The ascertained rule of gravitation he used but as a stepping-stone on which he might safely tread in advancing towards the great end of

\* Advertisement to *Optics*, 1717.

philosophy,—the reduction of all that is implied in the terms space, force, and matter, to the closest relations and the fewest agencies: he regarded this great discovery with no more partiality than he did the more undeveloped principle of molecular cohesion, with respect to which, after stating his general conception of the force, he comes to this conclusion—“there are therefore *agents* in nature able to make the particles of bodies stick together by very strong attractions; and *it is the business of experimental philosophy to find them out*\*.”

The term *attraction*, be it observed, was always employed by Newton in a provisional sense. “How these attractions may be performed,” he says, “I do not here consider: what I call *attraction* may be performed by *impulse*, or by some other means unknown to me; I use that word here to signify only in general *any force by which bodies tend towards each other*, whatsoever be the cause:” thus he was content to express, in any terms that lay at hand, the mathematical law, whilst he kept the efficient cause in reserve, laying down for the order of investigation this rule—“We must learn from the phænomena of nature what bodies attract one another, and what are the laws and properties of the attraction, before we inquire the cause by which the attraction was performed †.”

The cause of gravity, whatever it may be, he conceived must also lie at the foundation of all the other great classes of force which we observe, and till their laws and properties should have been learnt, he knew that it would be premature to attempt any deep inquiry into their causes. Nevertheless he let loose his fancy in more than one excursion into this wide field of speculation; and it is worth our while to mark the manner in which he surveyed it. For he possessed beyond other men that double power of mind which can adapt itself equally to the furthest and nearest limits of vision, and cast a glance as comprehensive over remote objects, as precise and penetrating into those that are within reach.

The widest of the generalisations to which the conjectures of Newton ascended were marked by a character far different from any which appears in the speculations of those who preceded him. Instead of loose or narrow analogies, in forming his ideas of the interior mechanism and materials of the universe, he clothed the phantoms of his philosophical vision with the most certain and general of the properties of matter: for the hooked atoms of Epicurus, the broken fragments, subtle powder, rounded globules, and feathery filaments of Descartes, he substituted the conception of particles embodying invariable powers of inertia, solidity, and hardness, with

\* Optics, Book 3. Qu. 31.

† Ibid.

forces centrifugal, or centripetal, varying with aggregation and distance. Of such particles, grouped in various modes and degrees of condensation, and variously moulded by the hand of the Creator, he thought all material things might be imagined to consist, by such, both the stability of nature and the conservation of motion might be maintained, and from such, all the great classes of phænomena might be derived.

The general name which he gave to the simplest of these particles was *æther*—a term which he used for the substance of one, or more, highly subtle and elastic fluids, capable of being combined and condensed, and taking, in different states of condensation, the form of light and ordinary matter.

His *æther* was not a mathematical or mechanical abstraction, but a material substance, of the actual existence of which, certain otherwise uninterpretable phænomena, especially of light, heat, and electricity, had convinced him, and which he conceived of, as being “much of the same constitution with *air*, but far rarer, subtler, and more elastic”—“not of one uniform matter, but composed, partly of the main phlegmatic body of *æther*, partly of other various *ætherial* spirits, much after the manner that *air* is compounded of the phlegmatic body of *air* intermixt with various vapours and exhalations,”—one of these spirits being the electric, another the magnetic, a third the gravitating principle. The latter he figured to himself as “not of the main body of phlegmatic *æther*, but of something very thinly and subtly diffused through it (perhaps of an unctuous, gummy, tenacious or springy nature\*), and bearing much the same relation to *æther* which the *vital aërial spirit*, requisite for the conservation of flame and vital motions, does to *air* †.”

This was the first speculation of Newton respecting “the cause of the gravitating attraction of the earth.” “For if such an *ætherial spirit*,” he adds, “may be condensed in fermenting or burning bodies, or otherwise coagulated in the pores of the earth and water into some kind of humid active matter, for the common uses of nature (adhering to the sides of those pores after the manner that vapours condense on the side of a vessel), the vast body of the earth, which may be every where to the very centre in perpetual working, may continually condense so much of this spirit as to cause it from above to descend with great celerity for a supply: in which descent it

\* Such expressions as these, used only in the earliest of Newton's speculations, appear to be in the style of the Epicurean school; but his meaning, as is evident from the variety of the terms which he uses, was only to describe in *popular language*, attractive and repulsive force.

† Registry Book of the Royal Society, vol. v. from 1675 to 1679, p. 67.

may bear down with it the bodies it pervades with force proportional to all their parts it acts upon, nature making a circulation by the slow ascent of so much matter out of the bowels of the earth in an ærial form, which for a time constitutes the atmosphere, but being continually buoyed up by the new air, exhalations, and vapours rising under, at length (some part of the vapours which return in rain excepted) vanishes again into the ætherial spaces, and there perhaps in time relents and is attenuated into its first principles. For nature is a perpetual circulatory worker, generating fluids out of solids, and solids out of fluids, fixed things out of volatile, and volatile out of fluid, subtile out of gross, and gross out of subtile, some things to ascend and make the upper terrestrial juices, rivers, and the atmosphere, and by consequence others to descend for a requital to the former. And as the earth, so perhaps may the sun imbibe this spirit copiously, to conserve his shining, and keep the planets from receding further from him: and they that will may also suppose that this spirit affords, or carries with it, the solary fuel and material principle of light, and that the vast ætherial spaces between us and the stars are for a sufficient depository for this food of the sun and planets\*.”

How far in a geometrical and mechanical point of view a supposition which presents to us the problem of an uniform central loss of force in a sphere of “*tenacious or springy*” fluid, urged by a constant pressure, and drawing down or impelling the bodies that float in it with a force proportional to the number of their ultimate particles, can have been contemplated as tending to satisfy the conditions of the law of gravity, I leave to mathematicians to judge. This supposition preceded the public announcement of the law by ten years; but Newton has himself stated that he had deduced that law from Kepler’s some twenty years before he published it†.

He soon, however, in a letter to Boyle in 1678, abandoned this form of hypothesis for one in which he supposes the æther no longer a gradually absorbed, centripetal, atmosphere, but a *stationary* fluid, “which consists of parts, differing from one another in *subtily* by indefinite degrees,” so arranged by the force with which the *pores of matter* repel the *ætherial particles* in proportion to their *magnitude*, “that from the top of the air to the surface of the earth, and again from the surface of the earth to the centre thereof, the æther is insensibly finer and finer;” and in an ætherial atmosphere so constituted he holds that bodies would be propelled towards each other by the as-

\* Registry Book of the Royal Society, vol. v. from 1675 to 1679, p. 68.

† Letter of Newton to Halley, 1686.

sumed greater repulsion of the larger particles of æther from their pores. In this letter he made a comprehensive conjectural effort to reduce the whole system of the laws of nature, whether bearing the aspect of impulse or attraction, under the dominion of *two kinds of repulsive force*, the one of mutual repulsion between the particles of æther, the other of repulsion between the particles of æther and those of ordinary matter.

In the edition of his Optics which he printed nearly forty years afterwards, in 1717, he deliberately delivered, when in full possession of the laws of gravity, another hypothesis on this subject, taking for his fundamental assumption this fact presumed from the phænomena of light, that a subtle and elastic fluid, within bodies and without them, follows some law of density which increases from their centre indefinitely into space, and merely representing the force by which they gravitate as *repulsive*. Further he has not explained himself; and it may perhaps be inferred from his subsequently omitting in an edition of the *Principia* the mention of *gravity* when he enumerates, at the end of that work, the other phænomena of molecular attraction and cohesion, electricity, light, heat, muscular motion, and nervous sensation, which he attributes to the force of "a very subtle spirit," pervading and lurking in dense bodies, but not yet sufficiently manifested by experiments,—that he was dissatisfied with his own conceptions of its gravific action, and had never reduced them into a mathematical form. Thus much however it may be worth while to remark, as deserving perhaps the attention of those who may follow Vince and Playfair in discussing the possible sufficiency of Newton's hypothesis—that in the Optics he alleges reasons for supposing the *elastic force* of ætherial particles to be *inversely proportional* to their *magnitude*\*. This leaves ground to believe that with the supposition of a *density* increasing with the distance he may have combined his former conjecture of an increasing *magnitude* of the particles, and so far an elasticity proportionably diminished; which gives latitude at least to the hypothesis, as making the mutual repulsion of the particles at different distances from the centre, depend on more elements than one.

But the knowledge of the experimental laws of molecular force was not sufficiently advanced to justify any serious attempt at mathematical theory, either on this subject or any other connected with them; nor did he offer these hypotheses as more than cursory hints, and specimens of a generalising and simplifying spirit of conjecture, so far illustrating nature,

\* Optics, ed. 4. book iii. p. 326.



as they embraced, and embodied, real facts and accurate conceptions of phænomena.

The subjects on which in this point of view the above-mentioned hypotheses, taken together with the questions in the Optics, threw the most important light, were the phænomena of colours, and of chemistry. I shall confine myself to his speculations on the latter subject, which lead directly to the question at issue—namely, what were the ideas of philosophers before the time of Black respecting the nature of air, and whether the *unity* of the aërial element was any part of their belief.

Most remarkable, among the *divinations* of Newton, is his *introduction* of the doctrine of chemical affinity in the optical *queries*, where he connects the phænomena of chemistry with those of electricity, as both due to *molecular forces acting at insensible distances*. He enumerates electricity among those “attractions which reach to sensible distances, and so have been observed by vulgar eyes;” he then suggests, that “there may be *others* which reach to so small distances as hitherto escape observation,” and adding that “perhaps *electrical attraction may reach to such small distances, even without being excited by friction\**,” goes on to couple it with the phænomena of *chemical affinity*, as produced by the same species of force. What is this, if it be well weighed, but the principle of all that experience has since brought to light in respect to galvanic and electro-chemical forces? here was the prophet’s eye, anticipating the progress of science and the actual indications of the kind of force which he surmised.

That which follows on the point of chemical affinity itself is equally remarkable. For observe how, guided in this instance by the few obscure phænomena before him, he deals with the molecules which represent this peculiar form of attraction: they are not elementary molecules, nor molecules of equal magnitude, but *compound particles whose force of affinity is in the inverse ratio of their composition*—“the smallest particles cohering by the strongest attractions, and composing bigger particles of weaker virtue, and many of these cohering, and composing bigger particles whose virtue is still weaker, and so on, for divers successions, until the progression end in the biggest particles on which the operations of chemistry, and the colours of natural bodies depend, and which by cohering compose bodies of a sensible magnitude†.” Have we not, in this conception of chemical affinity as depending on the *successive addition of units of force*, the principle of multiple proportions, of which the experimental demonstration was

\* Optics, book iii. p. 351.

† Ibid. p. 370.

reserved for Dalton, whose first views of that important induction were suggested perhaps by these very conjectures of Newton?

In other respects the theory of affinities is hardly laid down by him with more distinctness in this mature work, than in his younger speculations, in the earliest of which he applied it, as an universal property of bodies, to supposed ætherial fluids, and in the next to the factitious airs then recently discovered.

The chemist who remembers the modern observation, that gases (including that *vital aërial spirit* to which Newton compared his æther) are powerfully condensed in the pores of charcoal, on the surface of metals, and in the interior of a ball of spongy platina, cannot fail to be struck with the singular anticipation which the *first* of Newton's *hypotheses* display, of a close connexion between molecular attractions and chemical changes, and a subjection of the most elastic of bodies to both these forces in common. Nor will his admiration be diminished, when he finds the theory of elective and mediating affinities first broached for such a purpose as to explain the dark phænomena of muscular motion, and the material means through which the soul acts on the body, by the supposition of *relative degrees of sociableness and unsociableness* between the brain and muscles on the one hand, and on the other, a conjectural array of ætherial fluids imagined to be even rarer and more elastic than the most subtle and repulsive air\*.

After this, we are not astonished to find the same master mind, in its *second* survey, so laying down the theoretical map of gaseous chemistry, that in truth the chemists who followed, down to the æra of Higgins, Dalton, and Gay-Lussac, did little more than work out by experiment the principles which Newton had assumed.

The application of chemical principles to ætherial matter is contained in a letter to Oldenburg, from which I have already given some quotations, read before the Royal Society in Dec. 1675. This elaborate communication, strange to say, has never been printed, except in the ponderous and seldom opened volumes of Birch's history of that Society, and consequently is scarcely known, even in our own country, to men of science, otherwise than by a few extracts from that part of it which relates to light, published in the Philosophical Transactions by Dr. Young.

The theory of gases, as communicated to Boyle in 1678, you will find in Birch's life of that philosopher, or in Newton's collective works. In his letter to Boyle, after supposing

\* Letter to Oldenburg, Registry of the Royal Society, vol. v.

certain atmospheres of æther to surround the particles of bodies, and describing a pressure of elastic forces, which varying with the distance produces cohesion at small distances, and repulsion at greater, he deduces among other consequences this—"that the particles of vapours, exhalations, and air, do stand at a distance from one another, and recede as far from one another as the pressure of the incumbent atmosphere will let them: for I conceive," he says, "the confused mass of vapour, air, and exhalations, which we call the atmosphere, to be nothing else but the particles of all sorts of bodies of which the earth consists, separated from one another and kept at a distance by the said principle."

He then proceeds to distinguish the *three* different ways which nature has of "transmuting gross compact substances into aërial ones"—vaporisation—volatility—and the liberation of fixed air, and to propose a theory to explain the differences. From the hypothesis, to which I before alluded, of a double repulsive force, producing unequal degrees of ætherial pressure, he deduces different spheres of cohesion and repulsion for different bodies, and their particles, in proportion to their density and size: small particles are easily detached, and easily condensed; and this is the condition of volatile substances, and of liquids—"when the particles of a body are very small, as I suppose," he says, "those of water are, the action of heat may be sufficient to shake them asunder;" and "as fast as the motion of heat can shake them off, those particles, by the said principle, will float up and down at a distance from one another, and from the particles of air, and make that substance we call vapour." "But if the particles be much larger, they then require the greater force of *dissolving menstruums* to separate them." Thus he comes to the chief object of this letter, which was to illustrate the theory of gases—of the substances, that is, then recently discovered to be more *durably fixed*, and more *durably aërial*, than vapours or volatile effluvia. For this purpose, having assumed that the essence of such substances is, that their constituent particles are relatively larger and denser, and therefore, by hypothesis, more elastic than others in the *aërial*, and more cohesive in the *fixed* condition, he brings in the doctrine of chemical affinities, elective and mediate, to liberate them from their close state of cohesion, and force them out of the proximate sphere of compression into the remoter one of repulsion. And thus, as subsidiary to a wild play of philosophical fancy, were those great principles laid down, which experience has subsequently verified, and on which the whole fabric of the chemistry of solids, liquids, and gases, has been built.

In these views the new discovery of the various permanence and condensability of the gases has a conspicuous place: "On the same difference of size," he says, "may depend the more or less permanency of aërial substances in their state of rarefaction." "This may be the reason why the small particles of vapours come easily together and are reduced back into water, unless the heat which keeps them in agitation be so great as to dissipate them as fast as they come together, but the grosser particles of exhalations raised by fermentation keep their aërial form more obstinately, because the æther within is rarer. Nor does the size only, but the density, of the particles also conduce to the permanency of aërial substances: for the excess of density of the æther *without* such particles above that of the æther *within* them is still greater: which has made me sometimes think that the true permanent air may be of a *metallic* original, the particles of no substances being more dense than those of metals. This I think is also favoured by experience: for I remember I once read in the Philosophical Transactions how M. Huygens at Paris found, that the air made by dissolving salt of tartar would in two or three days' time condense and fall down again; but the air made by dissolving a metal continued without condensing or relenting in the least. If you consider then how by the continual fermentations made in the bowels of the earth there are aërial substances raised out of all kinds of bodies, all which together make the atmosphere, you will not perhaps think it absurd, that the most permanent part of the atmosphere, which is the true air, should be constituted of *these*; especially since they are the heaviest of all others, and so must subside to the lower parts of the atmosphere and float upon the surface of the earth, and buoy up the lighter exhalations and vapours to float in greatest plenty above them. Thus I say it ought to be with the metallic exhalations raised in the bowels of the earth by the action of acid menstruums; and thus it is with the true permanent air."

These extracts show that Newton considered the hydrogen gas which Boyle had obtained from iron, and the nitrous gas which Huygens had obtained from copper, as consisting of the ultimate particles of the iron and copper themselves, brought into a state of aërial elasticity; and further, that apprehending his aëtherial hypothesis to be thus strengthened by experimental facts, he proceeded to generalise so boldly, as to conclude that the whole body of the inferior atmosphere may be constituted of various metallic substances, and that the power and persistence of elastic force in different kinds of air may be proportionate to the size and density of their chemical elements.

This supposition, that the most permanent airs are of a metallic origin and nature, representing hydrogen for instance as *ferreous gas*, was set aside by the experiments of Cavendish, which proved that the gas from *iron* is identical with the gas from *zinc*, in specific gravity, in explosive power, in the quantity in which it combines with oxygen, and in the result of the combination: they went also, as far as our experiments reach, to invalidate the general supposition that the repulsive force of the particles of matter is in proportion to their weight and density; since they proved that hydrogen is, both in its *elastic* and in its *fixed* state, the lightest of bodies; unless indeed its high refractive power should be thought a stronger argument for the density, than its low combining weight for the lightness, of its molecules.

Newton seems not to have been aware, that the facts of the condensation of one gas and permanence of another, the observation of which he here ascribes to Huygens, had been established some ten years before by experiments instituted at the Royal Society, in which his correspondent Boyle had assisted—a circumstance however which was notified to the public when Huygens's paper was printed\*. The experiments themselves having, I think, never been published, the interest which we equally take in tracing back the history of science, the curiosity of the experiments, and the celebrity of the experimenters, prompt me to give you some extracts on this subject from the journals of the Society.

From these it appears that on January the 4th, 1664, a year before the publication of the *Micrographia*, Hook exhibited to the Society "experiments to show that air is the universal dissolvent of *sulphureous* [combustible] bodies, and that

\* An account of Huygens's experiments was printed at Paris in 1674, and appears in the *Philosophical Transactions*, No. 119, dated November 22, 1675, under the title of—"some experiments made in the air-pump by M. Papin directed by M. Hugen's." The following extract contains the facts to which Newton referred:—"The experimenter being desirous to see whether these ebullitions did make *new air*, put in the recipient a gage, and observed that when the liquors were mingled, the water in the gage rose very nimbly to the top of the gage; and drawing out the new air he made the gage-water subside again; and by this means it was seen, that all these kinds of ebullition make an air which expands itself like common air. Yet here is something that seems to be very remarkable, which is, that the air made by these ebullitions is *not of the same nature*: for it has been found experimentally, that the air formed by the mixture of *aquafortis* and copper remains always *air*, and always keeps up the water in the glass; but on the contrary, the air which has been made by the mixture of oil of tartar and oil of vitriol is almost all destroyed of itself, in the space of twenty-four hours. All these ebullitions hitherto spoken of are greater in *vacuo* than in the open air; but with lime it is not so."

this dissolution is fire, adding that this was done by a nitrous substance inherent in, and mixed with, the air."

Here was the first distinct conception, and evidence, of the composition of the atmosphere. The French physician Rey had before proved that air enters into fixed combination with solid matter: his proof rested on a capital observation which he quotes from the *Basilica Antimonii* of Hamerus Poppius: this chemist, "placing," says Rey, "a burning glass in the sun's rays, directed their focus on the apex of a cone of antimony, till the whole becomes white, when the calcination is complete. It is a wonderful thing, Poppius added, that although in this calcination the antimony loses much of its substance by the vapours and fumes which exhale copiously, yet so it is, its weight increases instead of diminishing\*." The philosophical acumen of Rey seized on the truth unequivocally shown in this simplified form of calcination, in which he discerned the presence of but two ponderables, and he concluded,—1. That the increase of weight arose from the air being solidified in the antimony; 2. That the two substances combined to a definite degree of saturation; 3. That the increase of weight observed in other metals, whether by calcination or simple exposure to the air, is due to the same cause—conclusions which, if their publicity had been equal to their value, would doubtless have been recorded for the early and distinct enunciation which they contain both of a fundamental principle, and an important though as yet unanalysed fact, as the first step in this branch of science, in consequence as well as time†.

But Rey, though he recognised the ponderable and combining qualities of air, considered it with the other philosophers of his day, as an element simple in essence, though mutable in form: and the first scientific question of the accuracy of this supposition was raised by Boyle in 1654. In the same Essay in which his discovery of the factitious airs was announced, he quoted from Paracelsus the following remarkable passage:—"As the stomach converts meat, and makes part of it useful to the body, rejecting the other part, so *the lungs con-*

\* Essay 25.

† Rey's work was first published in 1630. It contains, besides the speculation here mentioned, a just correction of the view which *the schools* had taken of a fact affirmed in the Physics of Aristotle—that a blown bladder is heavier than an empty one. Rey showed that this is true only if the bladder be blown to such a degree as to *compress* the air, and that the fact, so stated, is a real proof that the air has *absolute weight*. This is perhaps the first correct *published* statement of the weight of air, as an experimental fact. It is evident however from a letter of Baliani, quoted by Venturi, that Galileo had not only taught the same doctrine, but made his experiments on the specific gravity of the air before 1630.

sume part of the air, proscribing the rest;" and having observed upon it, that "though this opinion is not, as some of the same author's, absurd, it should not be barely asserted, but explicated and proved," proceeded to relate, that "that deservedly famous mechanician and chemist Cornelius Drebell contrived for the late learned king James a vessel to go under water," and that on inquiry of Drebell's surviving relatives into the principle of his contrivance, it appeared, that "he conceived it to be not the whole body of the air, but a certain quintessence or spiritual part of it that makes it fit for respiration, which being spent, the remaining grosser body, or carcase, if I may so call it, of the air, is unable to cherish the vital flame residing in the heart; so that, for aught I could gather," says Boyle, "besides the mechanical contrivance of his vessel, he had a chemical liquor which he accounted the chief secret of his navigation; for when from time to time he perceived that the finer and purer part of the air was consumed, or overclogged by the steams and respiration of those that went in his ship, he would, by unstopping a vessel full of this liquor speedily restore to the troubled air such a proportion of vital parts, as would make it again for a good while fit for respiration\*."

The experiments which Hook exhibited to the Royal Society in 1764 afforded, it must be allowed, but precarious grounds for the theory of the composition of the atmosphere which his sagacity advanced: he showed that in vessels containing a limited quantity of air, combustibles burn and waste for a limited time; and change its quality so that it is no longer capable of supporting combustion; and he showed that they undergo no loss of substance when heated without air: he took some live coals and put them under a glass vessel—"whereupon the said cole in a little time went out; but being then taken out, and exposed to the free air, recovered its burning:" sulphur in like manner would not burn when "hermetically sealed," and charcoal heated without air—"was not sensibly diminished;" he added—"that a combustible substance kept red-hot, yea in a fire as hot as to melt copper, would not waste, but as soon as fresh air was admitted did burn away and consume." Boyle proposed that trial should be made whether the extinguished combustible could be re-lighted by the burning-glass, or by red-hot iron; and it was found that it could not be rekindled without the admission of air.

Yet nothing can be more accurate than the theoretical account of combustion given in the *Micrographia*, where, laying

\* New Exp. Physico-mechanical.

down the principle that "the different volatility, or fixedness, of the parts of bodies seems to consist only in this, that the one is of a texture, or has component parts, which will be easily rarefied into the form of air, and that the other hath such as will not without much ado be brought to such a constitution," Hook states that "in the dissolution of *sulphureous* [combustible] bodies, by a substance inherent in, and mixed with, the air, which is like, if not the *very same* with, that fixed in saltpetre, a certain part of the bodies is united and mixed, or *dissolved and turned into*, the air, and made to fly up and down with it, in the same manner as a metalline or other body, dissolved into any *menstruum*, doth follow the motions and progress of that *menstruum* till it be precipitated."

Even the fact, afterwards proved by Cavendish, of the *density* of the gaseous product of this *dissolution*, was predicted by Hook; for in an experiment—"to prove that the substance of a candle or lamp is dissolved by the air, and the greatest part thereof reduced into a *fluid in the form of air*,"—he observes, that "the reason why this mixed body, which *certainly is otherwise heavier than the air*, and so ought to descend, doth notwithstanding ascend, is from the extraordinary rarefaction of the same by the nearness and centrality of the flame and heat, whereby it is made much lighter than the ambient air\*."

"\* Experiment to prove that the substance of a candle or lamp is dissolved by the air, and the greatest part thereof reduced into a fluid in the form of air—showed the Royal Society 22-29 Feb. 1671-2."—Registry of the Royal Society.

"Using a large reflecting glass, or convex refracting one, so placed in respect to my eye that a candle, set at a certain distance beyond the refracting glass, or between the eye and the surface of the reflecting glass, enlightened the whole area of the said glasses in respect to the eye, then continuing to keep the eye in that place where the area of the glasses appeared to be wholly filled with the flame of the candle, I caused another candle to be placed very near the said glasses, between the eye and the glass, or beyond where I used the refracting glass, then looking steadfastly at the flame of the last candle, it was very plain to be perceived, that the flame thereof was encompassed with a stream of liquor, which seemed to issue out of the wick, and to ascend up in a continued current or *jet d'eau*, and to keep itself entire and unmixed with the ambient air, notwithstanding that it was a considerable way carried above the aforesaid flame. 'T was yet further observable that the shining flame was placed in the midst of this *jet d'eau* at the lower end thereof, but that it did not ascend proportionally in height to the height of the *jet d'eau*, that where the tip of the flame ended, there ascended up a small line of an opacous body or smoke, which to a good height above the flame kept the middle of the stream. The manifestation of these phænomena was from the differing refraction of the body of the *jet d'eau* from that of the ambient air, for the flame of the first candle being but small and placed at considerable distance from the refracting and reflecting globe, the smallest variation in the refraction of the



The *production* of volatile salts in combustion, by an analogous process of combination, seems likewise to have been apprehended by him, where he represents "other parts of the combustible," not capable of the aerial form, as nevertheless so "mixing and *uniting* with the parts of the air," as "to make a coagulum or precipitation, as one may call it, which is separated from the air," but being light and volatile is carried up by its motion, till the agitation that kept it rarefied ceases, and it condenses into "a certain salt which may be extracted out of soot:" and the view thus expressed appears from the Registry to have been corroborated at one of these sessions by Boyle, who observed that "vegetables reduced in the open air yield store of volatile salt like that of hartshorn and other animal bodies, whereas in common distillations he had not found them to yield a grain."

Hook produced evidence also before the Society of that sameness of effect, by which he identified the particular ingredient in the air that supports combustion with one of the fixed constituents of nitre. To this purport he "made an experiment with charcoal enclosed in a glass, to which nitre being put, and the hole suddenly stopped up, the fire revived, although no fresh air could get in,"—and another "of gunpowder burning without air."

It is curious to remark that a similar experiment was made some fifty years before by the Cabbalist and Rosicrucian antagonist of Kepler and Mersenne, Fludd; who in proof "that the substance of saltpetre is nothing else but *air* congealed by cold\*," relates that he filled an egg with it, mixed with sulphur and quick lime, and closing the aperture with wax

medium between the first candle and the eye caused the darkness to intermix with the light, so as to exhibit the appearance of the heterogeneous *jet d'eau*. This *jet d'eau* I suppose to be nothing else but the mixture of the air with the parts of the candle which are dissolved into it in the flame. The reason why *this mixed body, which certainly is otherwise heavier than the air*, and so ought to descend, doth notwithstanding ascend with great swiftness, is first from the ascent of the flame in the middle, and next from the extraordinary rarefaction of the same by the nearness and centrality of the flame and heat, whereby it is made much lighter than the ambient air."

\* "Videmus salis petreæ substantiam nihil aliud esse quam *aërem* frigore congelatum, cui si accedit sulphuris aliqua portio, licet exigua, admodum strepitum ingentem edit, fulguraque artificialia emittit."—Utriusque Cosmi Historia, vol. i. tract. 1. lib. 7. cap. 6. De fulmine et tonitru, 1617. "In 2da demonstratione, candela in fundo vasis alicujus aqua repleti affigitur, cujus flamma per orificium phialæ ingrediens, depresso ejus orificio ad angulos rectos cum candela in vasis aquæ, sursum attrahet tantam aquæ proportionem quantam aëris in phiala inclusi consumpsit; aër enim nutrit ignem, et nutriendo consumitur; ac ne vacuum admittatur, aqua, hoc est tertium elementum, locum possidet aëris comesti."—Ibid. tract. 2. part. 1. lib. 3. Reg. 6.

placed the egg under water, where it exploded. Fludd also burnt a candle in a glass vessel over water, and observed that it raised the water in proportion to the quantity of air, enclosed in the vessel, which was consumed and burnt; for "air," he adds, "nourishes fire, and in nourishing consumes it." This sounds like the truth which Hook announced: but Fludd had no distinct idea of *the weight* of air, or of the great principle which led that philosopher to predict, and observe, *ponderable* products from its consumption.

Boyle supported Hook's views by "affirming that gun-powder burns very well in a receiver out of which the air has been extracted," and he afterwards took the pains to experiment with nitre compounded of nitric acid and potash out of contact with the atmosphere "*in vacuo* Boyleano," for the sake of "removing the suspicion that it does not burn without air being supplied by the numerous eruptions of the aerial particles *intercepted* by those that by their coalition make up the nitrous corpuscles\*." Boyle also remarked at this discussion that "*tin* mixed with nitre will kindle it;" to which Hook added, that filings of iron will do the same. This remark was justly deemed of such importance, that the Society ordered the experiment to be tried; and it was found that "filings of tin being cast on nitre, over a fire, made it flame; though it benot known," adds the writer of the Minutes †, "that *sulphur* was ever extracted out of tin; which seems to infer that there are bodies *combustible* which are not *sulphureous*."

The only verification, in the Registry, of the intimation given in the Micrographia that the same principle in the air which supports combustion is concerned "in respiration and the preservation of life," consists in an experiment suggested by Dr. Ent, in which a bird was enclosed with a chaffer of live coals in a receiver; the Society observed the extinction of the fire to be followed by failure of vitality in the bird, which revived on the readmission of air.

After these inquiries Dr. Wilkins proposed (on the 8th of March 1764) "that the following experiment (of Dr. Wren's suggestion) might be made, viz. to put a fermenting liquor in a glass ball to which a stop-cock should be fitted, and to tie a bladder about the top of the stop-cock, by which means a certain air generated by the fermenting liquor would pass into the bladder, and upon the turning of the stop-cock be kept there in the form of air without relapsing into water. This, or the like, to be tried at the next meeting. Mr. Hook mentioned several liquors that by their working upon one another

\* New Experiments touching Flame and Air.

† Oldenburg.

would generate *an air*; viz. oil of tartar and vitriol, spirit of wine and turpentine, &c. Colonel Blunt added that oysters pounded and put into wine would make it ferment."

"On the 15th of March, the experiment of generating air was made in this manner. There was taken a common glass phial with two pipes, and some pounded oyster-shells and aquafortis; and as soon as the aquafortis was by one of the pipes poured in upon the powder, and the hole stopped with a piece of hard cement, the ebullition caused by the corrosion of the shells by the aquafortis did in a very little time blow up the bladder (tied on the other pipe) so as to swell it with air, very plump; which expansion remained till the rising of the Society, when the vessel in that posture was locked up in the box of the watch, to remain there until the next assembly." Dr. Wren made use of this experiment—"to explicate the motion of the muscles by explosion." "There was also taken a bottle containing strong ale that had been bottled awhile; and over the bottle's mouth was tied an ox-bladder out of which the air was squeezed; after which by loosening the cork by degrees the air was blown out into the bladder by the expansion of the fermenting liquor within, and the bladder was almost half-filled with an aërial spirit generated by the working liquor." Mr. Boyle, bearing perhaps in mind his anecdote of Drebell's submarine vessel, suggested that the experiment was capable of improvement for the producing of air under water, and mentioned coral, or oyster-shells, and distilled vinegar, as wholesome substances for that purpose: he moved that an animal might be put into the receiver of his engine and the air exhausted till the creature grew sickly, and that then some new air might be produced in the receiver by a contrivance of making distilled vinegar work upon coral, to see whether by this means the animal could be revived. Dr. Wilkins moved that at the next meeting the air generated by the mixture of aquafortis and the pounded oyster-shells might be blown into a dog's or cat's mouth, to see what would be the effect thereof."

"On the 22nd of March there were two experiments made for the finding out a way to breathe under water, useful for divers. The first was made by putting a bird into a rarefying engine, and with it a glass bottle with distilled vinegar and pounded oyster-shells, which whilst the vinegar is dissolving them affords a stream supposed to be *a kind of new air* fit for respiration. The bottle was also close stopped with a cork, so ordered that by pulling the stop-cock placed on the top of the receiver, the cork might, by turning it, be pulled out without admitting an ingress of the external air into the re-

ceiver at all: then the receiver being accurately cemented to the engine the air was pumped out; whereupon the bird grew sick, and when he was thought near dying, the bottle was unstopped, that the streams and supposed air that had been shut up in it during the operation might have liberty to expand themselves in the receiver for the refreshing and recovering of the animal: but here it succeeded not; in so much that though the bird was taken out of the receiver and exposed to the fresh air, yet it recovered not."

"The other experiment was made with a kitling after the manner of the former, only that instead of distilled vinegar was employed aquafortis, whereof the success was, that the air being drawn out till the cat had done struggling, and was upon the point of expiration, and the bottle being unstopped to emit the streams and supposed air into the receiver, the cat did soon begin to recover, whereupon the animal had fresh air given it, which was again exhausted, to see whether it would revive of itself, without any nitrous exhalation; but after this exhaustion the cat appeared to be dying, whereupon she was after a little while taken out into the open air wherein she revived again."

"It was also moved that a standard might be used to know what quantity of air was generated."

"The glass phial with the swelled bladder, experimented upon at the last meeting and shut up till this day, was produced, and the bladder found evidently shrunk. Ordered to be tried next day with a glass phial whelmed under water, thereby to gather all the bubbles of the air generated by the corrosion."

"On being inquired how it was known that that which was supposed to be air produced by the dissolving of pounded oyster-shells by spirit of nitre, or distilled vinegar, or aquafortis, was true air, and answer being made by the President\*—that *a body rarefied by heat, and condensed by cold, was air*, the bladder was put to the fire where it expanded again as much as formerly and being removed from thence became somewhat flaccid again."

"It being moved that it might be tried whether the streams produced by the operation of distilled vinegar upon the powder of oyster-shells were convenient for respiration, the trial was made, and the bottle wherein that dissolution was performed, carried about to the company for every one to smell to it, and it was found by most of the company incommodious, as it was undiluted."

"It being moved by Mr. Hook that the air-boxes contrived

\* Lord Brouncker.

for diving might be tried by the persons bespoke by Mr. Pepys for diving, it was ordered that this diver should be sent to Mr. Hook to be instructed by him touching the use of the said boxes under water."

"On the 29th of March an experiment was made for the generating of air by putting aquafortis and the powder of oyster-shells in a small glass phial under water, and whelming a large glass filled with water over it to receive the steam to be generated by the corrosion: the success whereof was that the whelmed glass was filled about  $\frac{1}{4}$ th full with an aërial substance—ordered to be set by till the next meeting."

"It was moved that a way might be thought on, of producing an air that might be useful to respire."

"On the 12th of April Mr. Boyle proposed [inter alia] to try whether the eggs of silk worms and snails would be hatched, as also whether seeds would germinate and thrive, all, in an exhausted receiver."

"Dr. Goddard affirmed that plants live as much upon air as the earth."

"Mr. Hook, being called upon to give an account of one of the last days experiments touching the air generated by aquafortis and the powder of oyster-shells, reported that the greatest part of it was returned into liquor."

"The same was ordered to make, the next day, the experiment of generating air with bottled ale, supposed to be wholesome to breathe in, which the air hitherto generated is not\*."

On June the 14th "an account was given of an experiment of the growth of water-cresses in a receiver." Having been kept for a week in an exhausted receiver they showed no growth; the air being admitted "they grew in the same time two or three inches."

These experiments remained unprinted: but a more complete discussion of the same subjects not long afterwards appeared. In 1668, at the early age of 23, Mayow, adopting the theory of Hook, published a tract in which "he delivered his thoughts of the use of respiration, waving those opinions that would have it serve either to cool the heart, or to make the blood pass through the lungs out of the right ventricle of the heart into the left, or to reduce the thicker venal blood into thinner and finer parts, and affirming that there is *something in the air* absolutely necessary to life, *which is conveyed*

\* On the 24th of May in this year (1764) the following record is entered: "The king had been pleased himself to make the observation (on the variation of the needle) at Whitehall, and had found no variation at all, the needle standing in the meridian."

into the blood, which whatever it be being exhausted the rest of the air is made useless, and no more fit for respiration; where yet he doth not exclude this use, that, together with the expelled air, the vapours also steaming out of the blood are thrown out. And inquiring what that may be in the air so necessary to life he conjectures that it is the more subtle and nitrous particles with which the air abounds which are communicated to the blood through the lungs, and this *aërial nitre* he makes so necessary to all life, that even plants themselves do not grow in earth deprived thereof\*.”

In 1673-74 he gave a fuller account of his opinions in another treatise†, the views contained in which exhibit one of the finest examples extant of the success with which a man of philosophical genius, having seized a true principle, may deduce from the observation of a few facts distinctly apprehended a whole train of real and important consequences, long before the principle itself can be deemed to have been proved by demonstrative experiments.

In reproducing the theory of the *Micrographia*, he took no care to give the original author of it the credit which was due; and in his own turn is passed unmentioned by Lavoisier, who did not distinguish this precursor of his own discoveries from the rest of his chemical predecessors, on whom he pronounces this general censure—that “they all allowed themselves to be carried away by the spirit of their age, which contented itself with assertions without proofs, or at least often regarded very slight probabilities as such‡,”—a censure which it is but just to qualify by the reflection, that in experimental philosophy solid proofs are not to be discovered without the preliminary of happy conjectures.

In this tract Mayow expressly says, “Though the particles of air are very minute, and are vulgarly taken for an element of the greatest simplicity, it appears to me necessary to judge them to be a compound§; and he adds,—“it is manifest that the air is deprived of its elastic force by the respiration of animals much in the same manner as by the deflagration of flame.” The latter assertion he made good by experiment: he not only observed, but measured, *the amount of elastic force lost in both these cases*; and he proved that animals, when confined in air which has been already diminished by combustion, *survive but half the time* that they would have lived

\* “An account of two books—*Tractatus duo, prior de Respiratione*, a Joh. Mayow, Oxon. 1668.” *Phil. Trans.* No. 41. p. 833.

† *Tract 5. Med. Phys.* Imprim. Jul. 17, 1673.

‡ *Traité de Chimie.* Discours préliminaire, tome i. p. 16.

§ *Tract. de parte aërea igneaque Spiritus Nutri*, cap. 7. p. 114.

in an equal volume of common air\*. But he advanced little beyond his predecessor in demonstrating the air to be a compound. It is not to be supposed he says that that aërial supporter of combustion is *the whole air*, but only *a part of it*, which is more active and subtle than the rest; since a light enclosed under a glass expires, even whilst the vessel still contains abundance of air: for we cannot believe that the particles of air which *were* in the said glass can be *annihilated*, nor yet *dissipated*; since they cannot pass through the glass." But this reasoning, though probable, is not conclusive; since it was certainly possible that the enclosed air might have been diminished by condensation instead of abstraction, and have become unfit to burn and to be breathed by a total vitiation, instead of a partial loss.

Yet, after all, Mayow's reasoning appears to advantage by the side of Priestley's, or Scheele's, even when in the progress of experiment his nitro-aërial spirit, or fire-air, had been actually divorced from "*its consort*," and when the latter great chemist had approached a complete analysis of the atmosphere. For so difficult did Scheele find it to interpret his own experiments, that when he had in his hands the "*liver of sulphur*" which had produced a given diminution in a given volume of air,—when he had found the specific gravity of the diminished air to be less than that of common air, and the "*fire-air*," which he had succeeded in separating from numerous substances, to have a greater specific gravity, as well as a greater power of supporting combustion,—when by reuniting them he had recomposed an air with all the properties of common air restored,—when he had arrived at the conclusion—"that the air consists of two different kinds of elastic fluids," and that the "*fire-air*" makes between a third and a fourth of the whole bulk,—when coming finally to the ultimate question of the analysis, he failed to find the "*lost air*" in the *liver* of sulphur. Then he gave the reins to his imagination, and embracing the idea, that heat is a compound of "*fire-air*" with an imaginary substance invented by Stahl, concluded that by the action of a double affinity the "*fire-air*" in his experiment had combined with the *phlogiston* of the *liver* of sulphur, and that the compound had passed through the pores of the glass by which it had before been confined. Where

\* *Tract. de parte aërea igneaque Spiritus Nitri*, cap. 7. p. 101. "Comperi aërem per lucernæ deflagrationem in spatium ex parte circiter tricesimâ minus quam antea reductum esse. Postquam fumi lucernæ deflagrantis, quibus cucurbita prædicta repleta est, prorsus evanuerunt, vitrumque intus æque ac prius pellucidum evasit, conatus sum secunda vice lucernam in eadem accendere, radios solares in aliam camphoræ portionem, in vitro eo pariter suspensam, uti prius conjiciendo."

weight disappears, analysis is impossible. So he left the composition of the atmosphere to be demonstrated by those who believed, with Mayow, that elastic fluids cannot penetrate glass, and who took the pains to weigh both the air and the substances by which it was diminished; whilst he went on pursuing the phantom of his imaginative genius to the examination of imponderable essences and the great discovery of the chemical forces of light, and of the distinctions between the heat of contact, and the heat of radiation\*.

But what shall we say to the improvements of Priestley on the principles of Mayow? Priestley—who many months after he is said by you, and others, to have discovered oxygen gas, tells us himself, that he “had no doubt it had all the properties of genuine common air.” On the 1st of August 1774, Priestley with a burning-glass, following the method of Boyle, collected this gas and observed “that a candle burnt in it with a remarkably vigorous flame, but did not give sufficient attention to the circumstance at that time—that the flame of the candle, besides being larger, burnt with more splendour and heat, than in nitrous air exposed to iron or liver of sulphur.” In the October following, “I mentioned,” he says, “my surprise at the air I had got, to M. Lavoisier, but at the same time had no suspicion that it was wholesome, so far was I from knowing what it was that I had really found, and taking for granted that it was nothing more than such kind of air as I had brought nitrous air to be by the processes above-mentioned.” He mentioned it also to all his philosophical acquaintance at Paris and elsewhere, “having no idea at that time to what these remarkable facts would lead.” On the 19th of November however, having agitated it in water, he “found that a candle still burned in it as well as in common air,” though after “the same degree of agitation phlogisticated nitrous air would certainly have extinguished a candle.” “In this ignorance,” he adds, “of its real nature I continued from this time to the 1st of March following.” “But in the course of this month I not only ascertained the nature of this kind of air, though very gradually; but was led by it, as I then thought, to the complete discovery of the constitution of the air we breathe. Till this 1st of March 1775, I had so little suspicion of its being wholesome, that I had not even thought of applying to it the test of nitrous air;” “but it occurred to me at last to make the experiment, and putting one measure of nitrous air to two measures of this air, I found not only that it was diminished, but that it was diminished quite as much as common air, and that the redness of the mixture was likewise equal to

\* Scheele's Experiments on Air and Fire.



that of a similar mixture of nitrous and common air. After this I had no doubt but that the air from *merc. calcinatus* was fit for respiration, and that it had all the other genuine properties of common air. But I did not take notice of what I might have observed if I had not been so fully possessed by the notion of there being no air better than common air, that the redness was really deeper, and the diminution something greater, than common air would have admitted. *I now concluded that all the constituent parts of the air were equally and in their proper proportion imbibed in the preparation of this substance, and also in the process of making red lead\**—a conclusion identical with the ideas of Rey in 1630.

The next step in Priestley's inquiry was the employment of Mayow's mice, which convinced him that this air was *longer* respirable than common air; but his ideas of it were less accurate than Mayow's, for instead of considering it, with him, *a constituent part of nitric acid*, he thought it *a compound of nitric acid and earth*; and in December 1777, "no doubt remained on his mind that atmospheric air, or the thing that we breathe, consists of the nitrous [nitric] acid and earth, with so much phlogiston as is necessary to its elasticity, and likewise so much more as is necessary to bring it from its state of perfect purity to the mean condition in which we find it."

You now see the error into which you have fallen when you represent Priestley as discovering *before Lavoisier* that "this was a gas wholly different from all other gases formerly known," and may perhaps suspect that you are not justified in condemning as "*an unworthy and lamentable proceeding*" on Lavoisier's part, "*the intruding himself into the history of this discovery, knowing that Priestley was the sole discoverer.*" A property of this gas, which under Priestley's observation had led to nothing, in the hands of Lavoisier gave rise to one of the most important investigations in the annals of chemistry; he, it appears from your own admission, had ascertained the relations of this elementary substance to various bases and to the atmosphere, between August 1774 and March 1775, at which date the foregoing extracts show the "*sole author of the discovery*" to have "had no doubt that it had all the genuine properties of *common air*." Whoever may be called the discoverer of oxygen, whether Hook and Mayow, who first inferred its existence in nitre and in air,—or Boyle, who first *disengaged* the elastic gas from *minium*,—or Hales, who *collected* it from the same material,—or Nieuwentyt, who attri-

\* Experiments and Observations on different kinds of Air, vol. ii. p. 113, ed. 1790.

buted its elasticity to "the expansion of the fire particles lodged in the minium, supposing fire to be a particular fluid which maintains its own essence and figure, remaining always fire, though not always burning,"—or Priestley, who observed that it supported combustion,—or Lavoisier, who distinguished it as a gas, *sui generis*, and determined its principal combinations,—if the question be, which of these names deserves the highest place in "*the history of this discovery*," a philosopher I apprehend might be apt to hesitate,—especially perhaps between those which stand *first* in the list, and that which stands *last*.

But you have made a greater mistake in attributing to Priestley the discovery of *nitrogen*\*; and in that mistake have again wronged Cavendish of his due. Had you taken the trouble to read a paper on this subject which I have published from his MSS.†, you would have found that the same philosopher, who exceeded all his cotemporaries in analysing the air with accuracy, was the first who demonstrated it to contain, after burning, a mephitic gas, incapable of supporting combustion, and *distinct from fixed air*: you would have known that, some time before Priestley's publication in the Philosophical Transactions of March 1772, Cavendish communicated to him this paper, containing all the details of an experiment, in which a measured volume of air, confined under water, was passed backwards and forwards through a bent tube filled with powdered charcoal, and heated red-hot—the absorption was found to be definite, and the total loss of volume was ascertained—the fixed air was separated by soap-leys—the volume separated was observed and deducted—the specific gravity of the residual gas was examined, and it was found "rather lighter than common air;" lastly, it was found to extinguish flame, but to extinguish it, by the criterion of the watch, more slowly than fixed air. I know that you would be far from conceding to me that any experiment, however skilfully devised, carries with it its own conclusions: but then you would have known too the very words in which Cavendish conveyed those conclusions to Priestley—"The natural meaning of *mephitic* air is any air which suffocates animals; and this is what Dr. Priestley seems to mean by the word: but in all probability there are *many kinds of air* which possess this property: I am sure there are *two*—namely *fixed air*, and

\* This discovery has been also erroneously assigned to Rutherford, to whom Robison, who derides "the trifling or vague writings of a Nollet, a Ferguson or a Priestley," (Edinburgh Evangel. Physics,) would fain ascribe also a share in the discovery of oxygen.

† Report of the British Association, Append. to Address, p. 63.

common air in which candles have burned, or which has passed through the fire. Air which has passed through a charcoal fire contains a great deal of fixed air which is generated from the charcoal; but it consists principally of common air which has suffered a change in its nature from the fire. As I formerly made an experiment on this subject which seems to contain some new circumstances, I will here set it down\*."

This important communication Priestley scarcely turned to better account than that which he afterwards received from the same skilful friend, of the composition of water; he quotes it indeed explicitly, but most defectively, in his paper in the Phil. Transactions of 1772. "Mr. Cavendish," he says, "favoured me with an account of some experiments of his, in which a quantity of common air was reduced from 180 to 162 oz. measures, by passing through a red-hot iron tube filled with the dust of charcoal: this diminution he ascribed to such a *destruction* of common air as Dr. Hales imagined to be the consequence of burning: Mr. Cavendish also observed that there had been a generation of fixed air in this process, but that it was absorbed by soap-leys: this experiment I also

\* "I transferred some common air out of one receiver through burning charcoal into a second receiver, by means of a bent pipe, the middle of which was filled with powdered charcoal and heated red-hot, both receivers being inverted into vessels of water, and the second receiver being full of water, so that no air could get into it but what came out of the first receiver and passed through the charcoal. The quantity of air driven out of the first receiver was 180 oz. measures, that driven into the second receiver was 190 oz. measures. In order to see whether any of this was fixed air, some soap-leys were mixed with the water in the basin into which the mouth of this second receiver was immersed: it was thereby reduced to 166 oz.; so that 24 oz. measures were absorbed by the soap-leys, all of which we may conclude to be fixed air produced from the charcoal; therefore 14 oz. of common air were absorbed by the fumes of the burning charcoal, agreeable to what Dr. Hales and others have observed, that all burning bodies absorb air. The 166 oz. of air remaining were passed back again in the same manner as before, through fresh, burning charcoal into the other receiver: it then measured 167 oz. and was reduced by soap-leys to 162 oz.; so that this time, only 5 oz. of fixed air were generated from the charcoal, and only 4 oz. of common air absorbed. The reason of this was that since the air was rendered almost unfit for making bodies burn by passing once through the charcoal, not much charcoal could be consumed by it the second time; for charcoal will not burn without the assistance of fresh air, and consequently not much fixed air could be generated, nor much common air absorbed. The specific gravity of this air was found to differ very little from that of common air, of the two it seemed rather lighter. It extinguished flame, and rendered common air unfit for making bodies burn, in the same manner as fixed air, but in a less degree, as a candle which burnt about 80" in pure common air mixed with  $\frac{6}{5}$  of fixed air, burnt about 26" in common air, mixed with the same portion of this burnt air." The gas thus obtained by Cavendish was nitrogen, with perhaps  $\frac{1}{8}$  of carbonic oxide.

repeated, with a small variation of circumstances, and with almost the same result." He takes no notice of the distinction established in Cavendish's paper between "fixed air," and "common air in which candles have burnt or which has passed through the fire;" and so entirely does he misunderstand, or disregard, Cavendish's intimation of the relative levity of the latter when purified from fixed air by caustic potash, as to "conclude, after making several trials, that the air in which candles have burned" (without having been subjected to such purification) "*is rather lighter than common air\**;" whilst with regard to the *lost air*, which the paper communicated to him described as "*absorbed by the fumes of the burning charcoal*," he represents Cavendish as having ascribed that loss to the "*destruction of common air*."

Though Priestley however here proves himself *not* to have been, as you imagine, the discoverer of nitrogen, this indistinct, but fruitful, experimenter gave in the same document three original and pregnant notifications; for he announced in it—1. the effect of vegetables in restoring the respirable quality of the air; 2. the application of the known absorbing power of nitrous gas, as a test of that respirable quality; 3. his observation that candles burn with an enlarged flame in the gas produced by the distillation of nitre†. This observation it is from which those who call him the discoverer of oxygen should date the discovery: for he knew as much of the gas from nitre in 1772 as of that from minium in 1774; and it was the application of nitrous gas here stated, which led, in 1780, in the hands of Cavendish, to the first accurate analysis of the atmosphere, and in 1781 to the solution of the great problem—what becomes of the air lost in the combustion of hydrogen gas?

In scientific value doubtless there can be no comparison between the experimental inductions of Cavendish, or La-

\* "I could not find any considerable difference in the specific gravity of the air in which candles or brimstone had burnt out. I am satisfied however that it is not heavier than common air, which must have been manifest if so great a diminution of the quantity had been owing, as Dr. Hales and others supposed, to the elasticity of the whole mass being impaired. After making several trials for this purpose I concluded that air thus diminished in bulk is rather lighter than common air."—*Phil. Trans.* 1772, p. 164.

† "All the kinds of factitious air on which I have yet made the experiment are highly noxious, except that which is extracted from saltpetre or alum; but *in this even a candle burned just as in common air*. In one quantity which I got from saltpetre a candle not only burned, but *the flame was increased*, and something was heard like a hissing, similar to the decrepitation of nitre in an open fire; this experiment was made when the air was fresh made, and while it contained some particles of nitre which it would probably have deposited afterwards."—*Phil. Trans.* 1772, p. 245.

voisier, and the inferences of those earlier philosophers whose speculations we are engaged in investigating: but when we follow Mayow's deductions from the assumption, on probable grounds, that there exists in the atmosphere a gas which in the act of combining with other bodies produces the phænomena of combustion,—when we observe him concluding the identity of his gas with one of the components of *nitre* from the atmospheric production of that salt, and from the sameness of its effect in enabling substances to burn,—when we further observe him determining it to be fixed in *the acid component*\* of nitre, and supporting this view of the subject by alleging the sameness of the effect of nitric acid and of the burning-glass, in adding to antimony weight and specific medical properties,—when we find him extending these views to other substances, stating with most remarkable accuracy the acidification, in various cases and degrees, of sulphureous and fermenting substances by atmospheric exposure, and hence inferring that this gas is the principle not only of combustion but of acidity †,—when

\* “Jam vero cum pars nitri aërea in spiritu ejus acido existat, non vero in sale fixo, quod reliquam nitri partem constituit, uti supra ostendimus, concludere licet particulas igneo-aëreas nitri, quæ cum parte ejus aërea idem sunt, in *spiritu nitri* reconditas esse.”—*De parte aërea igneaque Spiritus Nitri*, p. 18.

† After stating (*Ibid.* p. 37) that the acid of oil of vitriol is not due to any acid already existing in sulphur, of which he says there are no signs, he adds—“potius putandum est, particulas ignis nitro-aëreas, in longa illa distillatione vitrioli, cum sulphure metallico Colcotharis congregari et effervescere: unde fit quod particulae sulphuris istius salinae, inter particulas igneas mutuo se atterentes interpositae, contundantur et comminuentur, ita ut eadem tandem exacuuntur, et ad fluoris statum perducantur, quæ demum ignis vi in altum delatae, oleum vitrioli componunt, haud multum secus ac spiritum sulphuris (sulphurous acid gas) per deflagrationem ejus fieri supra ostendimus.” “Si vitriolum ad totalem spiritus acidi expulsionem calcinatum, aëri humido aliquamdiu expositum fuerit, idem spiritu acido de novo imprægnabitur. Nempe spiritus nitro-aëreus cum sulphure metallico *Colcotharis* lente congregitur, motuque obscuro cum eodem effervescit; unde fit, quod particulae salinae, aut metallicae sulphuris istius, modo supra dicto, ad florem perducantur. Profecto vix concipi queat qua alia ratione spiritus iste vitriolicus in Colcothare produceretur; neque enim idem in Colcothare mox e distillatione extitit; neque putandum est eum *totaliter* ab aëre prosapiam ducere, ut alibi ostensum est.” “Vitriola e lapide seu potius glebe salino-sulphurea (vulgo Marchasitam vocant) conficiuntur, e qua igni commissa flores sulphuris vulgaris, copia satis ampla, eliciuntur: postquam autem gleba ea aëri, astrisque pluviis, aliquandiu exposita est, et dein, prout ejus fert natura, sponte sua fermentata est, eadem vitriolo ubertim imprægnabitur: nimirum spiritus nitro-aëreus cum sulphure metallico *mar-chasitarum* istarum effervescens, partem earum fixiorem in liquorem acidum convertit, qui mox ab ortu suo particulas metallicas lapidis adoritur, evocatque; tandemque cum iisdem in vitriolum coalescit. Quinetiam *Rubigo ferri*, quæ naturalem vitriolicam obtinet particularum nitro-aërearum cum sulphure ferri *metallico* congrementium actione produci videtur.” “Ani-

by an induction of the like kind we find him showing it to us as the principle by which metals gain weight from the air, and vegetables germinate and grow, and undergo an obscure fermentation [*æstum obscurum*] in their life and their decay\*,—lastly, when we find him ascribing to the same principle the phænomena of respiration, and representing the reduction of this gas from the elastic to the fixed state by its union with the blood, in the lungs and elsewhere, as the cause of its change of colour, its heat and its aptness for stimulating the heart and exciting muscular motion†—in contemplating so

madvertendum est insuper quod non tantum in rebus solidis, sed etiam in liquoribus, sal acidum, sive *achor*, spiritus nitro-ærei actione producatur." "Præterea nescio an non spiritus acidi e lignis ponderosis distillati, simili ratione per ignis operationem inter distillandum fiant." "Illud etiam obiter annotamus, quod spiritus acidi e saccharo, et melle, distillati, haud multum absimili ratione, per actionem spiritus nitro-ærei ignei, fieri videantur." "Liquorum autem fermentatio in eo consistit, quod particulæ nitro-æreæ aut *liquori insitæ*, aut *aliunde advenientes*, cum particulis liquoris salino-sulphureis [basic] effervescunt." "Huc etiam spectat, quod vina, aut cerevisia generosiora, radiis solaribus diu exposita, aut in loco calido detenta, processu temporis in acetum commigrant." "Ex iis quæ dicta sunt haud difficile erit intellectu quomodo spiritus acidus nitri in terra generatur." "Et ita demum ostendere conatus sum, quod salia quæcunque acida a particulis salinis, spiritus nitro-ærei ope, ad fluorem sive fusionem evecitæ producantur." "Quoad differentiam liquorum acidorum, eam a diversitate salium e quibus iidem constituuntur procedere putandum est, uti etiam ex eo, quod salia fixa, nunc magis, nunc vero minus a spiritu nitro-æreo alterantur, exacuenturque; et tamen inter salia acida quæcunque affinitas magna est et similitudo; inque iis omnibus particulæ nitro-æreæ-igneæque veluti in subjecto idoneo hospitantur." "Particulæ terræ salinæ hoc modo ad fluorem evecitæ hospitium idoneum fiunt, in quo particulæ nitro-æreæ recondantur detineanturque: ab iis autem utrisque strictim unitis spiritum nitri, qualis distillatione elicitur constitutum esse arbitror."

\* "In ortu vegetabilium spiritus nitro-æreus in motu et vigore positus, sulphur in statu fixo existens adoritur, quo tandem ad volatilitatem perducto, spiritus nitro-æreus in salinis vinculis incarceratus figitur." "Nostra fert opinio etiam fermentationem ad vegetabilium interitum tendentem a particulis nitro-æreis et salino-sulphureis, se invicem commoventibus, procedere." "Spiritus nitro-æreus a conjuge sua salina violenter abruptus motu suo impetuoso omnia perturbat, mixtique compagem solvit." "Ea quæ spiritum nitro-æreum excludunt res a corruptione vindicant." He instances fruits, flesh and butter, as being preserved from putrefaction, and iron from rust by things which exclude this gas, especially inflammable things, such as oil.

† "Quenadmodum particulæ nitro-æreæ terræ spiracula\* lente subeunt, ibidem cum particulis salino-sulphureis, iis vero immaturis, æstu obscuro congregiuntur, a quo vegetabilium vita dependet—ita particulæ eædem nitro-æreæ magis confertim in cruoris massam pulmonum ministerio introductæ, particulisque ejus salino-sulphureis ad justum vigorem evecitæ quoad minima admixtæ, fermentationem satis insignem, qualis ad vitam animale requisita est, efficiunt."—p. 147. He states that the colour of arterial blood has been shown by Lower, to be owing to the admixture of air with it in the lungs, and lays it down, that the heat of the body is due

just and splendid a generalisation, running parallel to the whole range of chemical induction on all those subjects which occupied the succeeding century, it is impossible not to allow that this young man handed down a bright light to all who followed him\*, and made more of a few facts, than the greater part of the next generation did of many.

Mayow also examined the two kinds of air which Boyle had obtained by the action of the nitric and vitriolic acids on iron, and observed the permanence of the one gas and the partial condensation of the other. To determine whether they resembled common air in containing any of the nitro-aërial *aura*, he added them to air in which a mouse was confined, and inferred that they do not, from their not prolonging the animal's life. He then examined their relative elasticity, and finding in them the same capacities of compression and expansion as in common air, he decided that there exist various elastic fluids, and held with Newton that these, as well as that *aura* which he deemed pre-eminently elastic, and the residual gas from which it is abstracted by respiration, owe their different degrees of elasticity and permanence to elementary differences in their particles, and in the substances from which they are derived †.

The only philosopher, as far as I am aware, who dissented from these views, was the elder Bernoulli, having detailed his own respecting fixed air ‡, “Mayow,” he said, “after

to the combination of these nitro-aërial particles with the blood, and the increased heat in exercise to a greater number being breathed in the same time. In like manner he accounts for febrile heat, for acid in the blood and urine, for the digestion of the food, and for muscular contraction.

\* Mayow's work, besides its publication in England, was at least twice reprinted abroad; a detailed account of it was given in the Philosophical Transactions. It was repeatedly quoted by Hales, whose book was in every chemist's hands, and by other authors: it was therefore sufficiently known to have produced a real influence on the minds of men.

† *De Spir. Nit.* cap. 9. p. 163.—“Utrum aër de novo generari possit?”—In his account of Boyle's gases he says—“*Aura* prædicta haud minori vi elastica quam aër vulgaris donatur prout sequenti experimento mihi compertum est.” “imili ratione experimentum feci, num aër in quo animal, aut lucerna expirassent, æque ac aër inviolatus, vi elastica pollent; et quidem mihi videtur aër iste haud minus quam aër quivis alius se expandere.” “quanquam *aura* ista in qua animal aut lucerna expirarunt vi elastica æque ac aër inviolatus pollet, et tamen eadem particulis nitro-aëreis vitalibusque destituitur.” “Hic etiam referre possumus quod in cap. sup. de *aura* hujusmodi aërisque vulgaris differentia annotavimus, et tamen verisimile est *auræ* istiusmodi cum aëre vulgari magnam affinitatem intercedere, vimque elasticam eorum utrorumque a causa haud multum diversa provenire. Etenim cum ferrum e particulis rigidis, item spiritus corrosivi e particulis nitro-aëreis summe elasticis constant, *aura* ex iis ntrisque invicem fervescentibus conflata ab aëre vulgari haud multum diversa crit.”

‡ “Allata experimenta satis, ni fallor, ostendunt existentiam aëris in cor-  
*Phil. Mag.* S. 3. No. 190. *Suppl.* Vol. 28. 2 M

various experiments concluded, that the substance itself of the globule, from which air was produced in them, is changed into an 'aura,' as water is changed into vapour by heat: but unlike vapour, this *aura* remains *aura*, and as he himself proves by experiment, retains its elastic force. Does there exist then any other body besides air which is fluid and endowed with elastic force? I scarcely believe it. He alleges indeed as a reason for denying to this *aura* the nature of common air, that he has found by experiment that the said *aura* is incapable of supporting life: as if, because it does not support life, therefore it cannot be air! we see our atmospheric air itself in time of pestilence unfitted to support life: has it therefore ceased to be air? it would be absurd to say this. It is not to be denied that in the space E. H. [of a glass tube filled with carbonic gas] other particles besides air find room, separated perhaps by the impetuous motion of the effervescence from the acid liquor, or the solid globule, and carried

poribus, sed et alterum nobis ostendendum est, nimirum quod aër iste sit aëre naturalis consistentiæ densior. Hoc autem sequenti experimento demonstratur. Sumatur vasculum liquore quodam acido semiplenum, ut A. C. D. B., et tubus aliquis vitreus E. F., altera parte E. clausus, altera vero F. apertus, impleatur eodem liquore; hujus vero orificio F. induatur globulus G., de luto, vel creta, in quibus nempe multæ particulæ alkali insunt, confectus; statimque, indice super orificium F. posito, invertatur tubus; et liquor in vasculo contento immergatur orificium F.; amoto digito, mox observabitur magnam effervescentiam excitari, quæ per aliquot horas durabit, donec omnis aër, intra particulas alkali contentus, solutis vinculis quibus coarctabatur, ad superiora ascenderit, et materia subsederit; tum demum animadvertitur, aërem hunc, postquam despumaverit, in suprema parte depresso liquore, magnum spatium E. H. occupare: quandoquidem autem superficies H. liquoris in tubo altior est superficie liquoris in vasculo, erit aër in spatio E. H. contentus aliquantum rarior aëre externo; proinde, ut fiat ejusdem consistentiæ, opus est ut aut tubus altius immergatur, aut plus liquoris affundatur, donec superficies interna coincidat cum superficie exteriori; quo facto, erit quidem spatium E. H. priori paululum contractius, et aër in eo contentus naturalis consistentiæ: nihilo minus tamen adhuc majus erit, duplo, triplo, quadruplo, (pro diversitate materiæ terrestris ex qua globulus conficitur, qua scilicet plus vel minus particularum alkali in se continet,) quam quod tota moles globuli G. occupat; quod certum indicium est aërem istum, cum omnis adhuc in globo continebatur, multo densiorem fuisse quam aër externus est: posito enim globulum constare, ex una parte, materiæ terrestris, et ex una parte, pororum, quibus nempe aër condensatus inest,—vel, quod eodem recidit, spatium quod materia terrestris occupat esse æquale spatio quod aër in poris contentus replet,—si nunc spatium E. H. sit duplum spatii globuli totius, sequitur, aërem in globulo contentum quadruplo densiorem esse quam est aër externus, si triplum sextuplo, si quadruplum octuplo, et sic porro in subdupla ratione; si vero ponatur spatium materiæ terrestris non esse æquale spatio pororum, sed in alia ratione majoris vel minoris inæqualitatis, densitates aëris in globulo æque facile ad calculum revocari possunt.”—Bernoulli, *Dissertatio de Effervescentia et Fermentatione*, No. 1. p. 20. 1690.



up with the air. Nor can we wonder that such an air, filled with miasmata, if breathed by animals, cannot keep them alive, especially when it is obvious that the spirit of nitre, and the globule of iron, used by the distinguished author, abound in many impure and poisonous particles, which if introduced into the system in breathing, may well corrupt the mass of the blood and induce death. If instead of the spirit of nitre he had chanced to use another acid liquor of a more *benign* quality, for instance the spirit of vitriol, and instead of a globule of iron, had taken one of an earthy kind, as in my experiment, the animal doubtless would not have perished, or at least would have lived longer. So that we may collect from this, not that the air, *as air*, destroyed the animal, but only incidentally, so far as it abounds with particles of a different kind and unfit for the support of life. But that we may make certain of one fact—namely that the substance of the globule itself is not changed by the fermentation into air, but that air really pre-existed in the globule, and was therefore *not* generated anew, the following experiment may be tried. Let the weight of an earthy globule, well-dried, be taken with perfect accuracy before the effervescence: then after the effervescence, when all the particles of the globule subside to the bottom, let the whole mass of the globule, which now lies dispersed, be carefully re-collected from the liquor; and let it be well-dried as before: lastly, let the weight also of the dried material be accurately ascertained by the help of the balance: this done, we shall find that the substance of the globule has lost nothing of its weight, or at least scarce a hundredth part, which perhaps exhaled with the air during the effervescence. But according to Mayow, it ought to have lost by far the greatest part of its weight; since it follows from his hypothesis, that the whole body of air occupying the upper part of the tube was taken from the substance of the globule; and so its weight should have been notably diminished, which nevertheless is contrary to experiment.”

In this criticism Bernoulli overlooked the chief fact on which the theory of Mayow rested—the constant diminution of the volume of common air, when breathed or burnt. And his attempt at an experimental refutation of it may serve to convince you of the danger which the greatest men may incur when they venture on deciding chemical questions without a knowledge of chemistry. To give the utmost credit to the alleged result of his experiment we must presume the “acid liquor” employed in it to have been *oil of vitriol*: but any boy in a chemist’s laboratory could have told him that the vitriolated lime which he collected at the end of the experiment

was a different substance from the chalk with which he began it, and that the consequence therefore which he drew from the weight remaining the same was a fallacy. The compounds of sulphur, I perceive, are a stumbling-block even to you \*: to Bernoulli's reputation as an experimental philosopher they are more fatal than he deemed them to animal life; for the wholesomeness of the air from so "benign" an acid as oil of vitriol, and "a globule of the earthy kind, as in his own experiment," was an assumption which the first trial would have discovered to be false. But it is more surprising that the computations, congenial to his own studies, which he proceeded to make, of the amount of condensation of the air, in the pores of the chalk, should not have apprised him that the globule on which he experimented *must have lost weight*, when so great a volume of condensed air was disengaged from it.

At a later period the younger Bernoulli paid so much respect to his father's opinion as to speak of the multiplicity of airs as a doubtful question. "The question," he says, "has long been agitated, whether the factitious elastic *aura* brought out of bodies, is ordinary air, or not; which question I shall not decide. If however the air of gunpowder be taken to be 1000 times denser than natural air, and 10,000 more elastic, then it follows from what precedes, that air compressed by an infinite force cannot be condensed more than 1331 times, and according to the same rule the elasticity of an air four times more dense than the natural air would be to the elasticity of natural air as  $4 + \frac{1}{4}$  to 1. But whether the experiments instituted by others, which make the ratio of these elasticities as 4 to 1, were conducted with sufficient accuracy, and whether the heat of the air, whilst under pressure, remained the same, I know not. It is probable however that the same *aura* which lies latent in the pores of the gunpowder is the cause of the elasticity of elastic bodies, and contractile *villous* materials; for when bodies are reduced by any force to an unnatural form, the elastic *aura* abounding in the little vacuities is compressed, and in giving the form of greatest space to those vacuities brings the body back to its original shape and extent."

In the English school, however, of pneumatic chemistry, and in the chief successor to the views and experiments of Boyle and Hook, of Mayow and Newton, there was no hesitation on this point. You have quoted the opinion of Hales, as representing, *instar omnium*, the general notion among experimental philosophers before the time of Black, that air was a single and simple element; and your inaccuracy on this point is not to be wondered at; because Hales's opinion has

\* Note to the Lives of Cavendish, Watt and Black, vol. ii. p. 511.

been over and over again mis-stated, even by eminent chemical writers. Those however, who are better occupied in making scientific discoveries than in reviewing them, may be excused, if they appear to be often less exactly acquainted with the opinions of others than with their own, so far at least as we can fully acquit them of desiring to exalt their own views, or the views of a particular æra, or a favourite author, by underrating all that has gone before.

The mistake in this case has certainly in great measure arisen from the circumstance, that the inquiries of Hales were directed more to the generic and physical properties of gases, than to their specific and chemical distinctions. He calls "airs generated in effervescences"—"true permanent air"; he has been supposed to mean that they are true *atmospheric* air; his real meaning was—that they are true elastic fluids, and, with the same permanence of constitution, possess the *same elastic force* as common air. This important fact had been before announced by Mayow, but was first ascertained with precision by Hales. "That I might," he says, "with the greater degree of certainty be assured of the degrees of compressibility of these different airs, I divided the capacities of two equal tubes into quarters of cubic inches, by pouring severally those quantities of water into the tubes, and then cutting notches with a file on the sides of the tubes at the several surfaces of the water; by which means I could see, by the ascent of the compressed water in the tubes, that both the factitious and common air were exactly alike compressible in all degrees of compressure, from the beginning till they were loaded with a weight equal to that of three atmospheres, which was the furthest I durst venture for fear of bursting the glass\*." Having made this contribution to our knowledge of the physical properties of the gases, and established that at common pressures and temperatures "with equal weights they are compressed exactly in the same proportion with common air," he went on to examine whether there exists any difference of specific gravity between the air and them; but contenting himself with the single experiment to which I have already referred, where no difference could be detected †, he left to Cavendish the grand discovery of the distinctions of density in elastic fluids; and it may possibly increase your respect for that discovery to remark that his false conclusion led him into much error in computing the weight of ærial substance fixed in various bodies from the volume which they yielded, on the supposition that the density of all airs is the same.

\* Stat. Essays, Append. p. 314.

† Analysis of the Air, Exp. 77.

Hales however rendered essential service to what may be more strictly called the *chemical* philosophy of aerial fluids. I have before noticed that we owe to him the discovery of a fact in gaseous chemistry, the consequence of which it is impossible to overrate—the condensation of atmospheric air by nitrous gas, in such a manner that *the two gases were observed by him to occupy the same space*. He first also determined with *numerical* exactness, and by very ingenious methods, the volume of air absorbed in a variety of chemical processes, and stated in the clearest terms the chemical nature of that *absorption*,—a statement adopted, as I have shown, by Cavendish, and strangely misconstrued by Priestley. “They were changed,” he says, “from a repelling elastic to a fixed state by the strong attraction of other particles which I call *absorbing*.” He taught the chemists of the succeeding generation how to procure almost all the gases which formed the subjects of their investigation; and he taught them also the more important lesson of conducting those investigations by *measure* and *weight*. Some of his experiments led directly to the most important conclusions at which they arrived. It was not for nothing that he observed that the “Sal Tartar” (very highly calcined) with which he essayed to purify the air for respiration had “absorbed one-third of the fuliginous vapours which arose from the burning candle\*,” or that he recorded experiments on phosphorus, in which “2 grains, fired in a large receiver, flamed and filled the retort with white fumes, expanded into a space equal to 60 inches, and absorbed 28 cubic inches of air;” and “when 3 grains were weighed soon after it was burnt, it had lost half a grain of its weight†.”

It is true that he made no advance towards analysing the air: and further, he argued, and argued justly, that “the sudden and fatal effect of noxious vapours, which has hitherto been supposed to be *wholly* owing to the loss and waste of the *vivifying spirit of air*, may not unreasonably be *also* attributed” to other causes, which he enumerates. “If,” he says, “the continuance of the burning of a candle be *wholly* owing to the *vivifying spirit*, then supposing, in the case of a receiver capacious enough for a candle to burn a minute in it, that half the vivifying spirit be drawn out with half the air in 10 seconds of time, the candle should not go out at the end of those 10 seconds, but burn 10 seconds more; which it does not, therefore the burning of the candle is not *wholly* owing to the *vivifying spirit*, but to certain degrees of the air’s elasticity,”—a principle which he goes on to illustrate by the “common

\* Analysis of the Air, edit. 1727, p. 272.

† Ibid. p. 169.

observation, that in very cold frosty weather fires burn most briskly\*.”

But we are by no means to conclude from hence that Hales had any doubt of the plurality of elastic fluids; on that point he quotes, as at once the foundation and the result of all his inquiries, the opinions expressed by Newton in the Optics:—“The illustrious Sir I. Newton,” he begins, “observes, that true permanent air arises by fermentation, or heat, from those bodies which chemists call fixed, whose particles adhere by a strong attraction, and are therefore not separated or rarefied without fermentation, those particles receding from one another with the greatest repulsive force, and being most difficultly brought together which upon contact are most strongly united.” “Dense bodies by fermentation rarefy into *several sorts of air*, and this air by fermentation, and sometimes without it, returns into dense bodies†,” “of the truth of which,” Hales adds, “we have proof from many of the following experiments.” And as he begins, so he ends: for having again repeated the same quotation from Newton, he closes his “*analysis of air*,” by drawing this general inference from all his researches—“Since we find in fact from these experiments that air arises from a great variety of dense bodies both by fire and fermentation, it is probable they may have very different degrees of elasticity in proportion to the different size and density of their particles, and the different forces with which they were thrown off into an elastic state.”

What now, give me leave to ask, becomes of your statement, that “when D’Alembert wrote the article ‘*Air*’ in the Encyclopédie in 1751, he gave the doctrine then universally received, that all the other kinds of air were only impure atmospheric air, and that this fluid alone was permanently elastic?” You tell us elsewhere that D’Alembert disregarded inductive philosophy, and professed himself ignorant of chemistry: and thus I should have accounted for his ignorance on this point, if I had not found on consulting the volume which you quote, that he really expressed *no such opinion* respecting air, and moreover has stated fully the views entertained of it by those, who in his own words, “supposent qu’il peut être produit et engendré, et que ce n’est autre chose que la matière des autres corps, devenue par les changemens qui s’y sont faits, susceptibles d’une élasticité permanente.” D’Alembert says indeed, that *some of the ancients* considered the air as a *simple element*, but remarks with truth, that they did not attach the same sense to that term as ourselves.

I have now completed the sketch which I promised, of the

\* Analysis of the Air, edit. 1727, p. 247.

† Ibid. p. 312.

gradual advance of this branch of science, in the æra beginning with Boyle and ending with Hales, during which the prevalent theory *multiplied the number of gases beyond the truth*, by supposing them as numerous as the substances from which they were obtained. This may be regarded as the æra of the first regular school of inductive science (if we except the less perfectly methodised school of Galileo), instituted by the original members of the Royal Society, for the professed purpose of executing the grand design of Bacon.

We have been lately told by a very able and lively writer, that the sole use and effect of the *Novum Organum* of the great founder of this school, was to bring down philosophers from high but barren aims to the level of common utility,—that “the inductive method has been practised ever since the beginning of the world,”—that “it is not only not true that Bacon invented it, but that it is not true that he was the first who correctly analysed that method and explained its uses,”—and that “he greatly overrated its utility;” we have been further told that the difference between the “*instances*” which make an absurd induction, and those which constitute a sound one, “is not in the *kind* of instances, but in the *number* of instances; that is to say, the difference is not in that part of the process for which Bacon has given precise rules, but in a circumstance for which no precise rule can be given:” and this notion of philosophical induction is illustrated by asking, “Will ten instances do, or fifty, or a hundred? In how many months would the first human beings who settled on the shores of the ocean have been justified in believing that the moon had an influence on the tides? After how many experiments would Jenner have been justified in believing that he had discovered a safeguard against the small-pox? These are questions,” it is added, “to which it would be most desirable to have a precise answer; but unhappily they are questions to which no precise answer can be returned\*.” Certainly, if the force of induction, and the inquisition and demonstration of truth, *does* depend on the “*number* of instances,” and not on “the *kind*,” Bacon has written in vain; but *you* know, my Lord, how it was, that in the hands of one who had better studied “the inductive method,” a vague and local idea, darkened with errors that destroyed its credibility and use, passed through the *mint of a very few decisive experiments* into the *treasury* of accepted truths: that which this author esteems the inductive process had been repeated thousands of times without fruit; but when Jenner, after vaccinating a child, inoculated it, and found that it resisted the virus of

\* Life of Lord Bacon, p. 411.

inoculation, the probability that "he had discovered a safeguard" rose at once, by the force even of a single experiment, to an amount which medical experience could "precisely" assign.

Mr. Macauley considers the credulity of those whom he calls "the dupes of Mesmerism," as due, not so much to neglect of these laws of evidence, as to want of natural sagacity; but the history of science by no means justifies this view of unfounded opinions: the truth is, that all sciences, except the mathematical, had stood for centuries in the same position in which such studies as go by the names of Animal Magnetism and Craniology, appear to stand now,—the position, that is, of collections of alleged facts unscrutinized and unsifted,—of generalisations precariously deduced, and truths, where they contained any, mystified and confused.

This was the state of science when Bacon appeared. The master science of *evidence*, like every other science, requires for its perfection both *rules* and *examples*. Bacon gave the *rules*. It has been observed by one well-qualified to offer an opinion, that "he traced not merely the *outline* but the *ramifications* of science that did not yet exist\*." But the chief, the all-pervading, ever-during benefit,—the force of direction, which he gave to the progress of knowledge, consisted in this—that he *did* "first analyse the inductive method correctly, that he first taught the specific value of every part of its evidence, and *that* with such precision," and such impressiveness, that a great school was founded upon his writings, who have handed down from him the torch of science, and have proceeded during the last two hundred years to practise, and mature, his rules.

Yet we shall do no more than describe a real change in the history of inductive science, if we shall proceed to speak of the experimental school of the æra which commences with Black and Cavendish, as the school of Newton: for the severe reason of the mathematician, grafted on that inductive principle of simplifying, and hedging in, ideas more complex than space and number, till they are divided and narrowed to the point of demonstration, shone forth in Newton's immortal works, and especially in his Optics, with a light as much more powerful than even the luminous lessons of Bacon, as example is more powerful than precept.

In this I believe you will agree with me, that if in our seats of learning the attentive study of *such examples of reasoning* had been made one of the essential requisites of an accomplished and sound education, we should not have seen so

\* Playfair's Dissertation, Encycl. Brit., p. 55.

many educated persons, ignorant of the laws of evidence and unconscious of their own deficiency, become as easy victims as the most ignorant, to wild paradox and blind credulity.

What the Optics were for experimental philosophy in general, that little unpretending *duodecimo* volume, of scarce a hundred pages, which Black published in 1755, on the properties of Magnesia, was to chemistry. It was, as you say, the second instance of a most beautiful example of inductive research; and the method of reasoning pursued in it deserves to be more particularly described, as constituting indeed the highest of all its merits. Not one word is there here of the *sulphureous* principle of the old chemists, or the corresponding *phlogistic* of the new: but there is, observe, *one general established principle*, reigning in the experimenter's thoughts, governing his hand, interpreting every phænomenon as it presents itself, dictating every successive experiment, and bringing forth each consequent discovery in that brief and transparent investigation.

The principle by which it was thus illuminated, was *the principle of elective affinities*,—a principle, first stated as we have seen, and generalised by Newton, experimentally noticed by Mayow, with others of the early chemists, and then recently systematised and *tabulated* by the French chemist Geoffroi. And here if we adopt such expressions as yours, in calling this “an example of *strict inductive investigation*,” let us understand clearly what we mean; let us not forget that the process of what is called the *inductive* method, in its most usual applications to such a science as chemistry, does not differ from that which is called *deductive* in *mechanics*, otherwise than in the degree of our reliance on the generality of the laws to which it is applied: in mechanics we now assume the laws which we have observed, to be applicable to *all matter* whatever; in chemistry, when the *nature of the subject is widely different* from those on which we have experimented, we dare not trust the *certainty* of our generalisations: the firmest believer in ætherial matter would hesitate to presume on Newton's hypothesis of its possessing chemical affinities, as a certain truth; and gas was to Black what *æther* is to us. His reasonings respecting fixed air were in fact all *deductions* from the presumed principle of elective attractions; but as far as regards the chief point of his discovery—the silent transference which he remarked of the gaseous substance that, as Hales had taught him, was fixed in salt of tartar, to calcined magnesia, and again from magnesia alba to caustic lime,—*the principle* which suggested the remark and the experiments, was itself *confirmed and established*, in its extension to gase-



ous substances, by *the result of those experiments*. In every such course of research, whatever offers itself *fortuitously* is observed by an eye which is on the watch for the appearance of the laws, known or assumed, that fill its meditations; and the whole *design* with which each experiment is instituted, is to test the applicability of those laws, and to try the validity, or the accuracy, of principles which have more or less the character of *foregone conclusions*.

This is experimental philosophy: this is the science of observing, interrogating, and interpreting, nature—apart from that faculty of catching *far analogies* on the wings of a lively and just imagination, which constitutes perhaps the highest part of the *genius* of a philosopher, though we should be much in error, if we regarded even this high gift of Heaven as incapable of being improved by rule, example, and use.

Thus it was that Black, under the guidance of the light which a clear conception of the laws of affinity shed over his mind, proved by a short series of experiments so devised as to eliminate, one by one, *all alternative suppositions*, the following points:—1. That magnesia is a distinct substance, having its own laws of combination to distinguish it from other earths—2. that that substance, which is sometimes found in air, and sometimes fixed both in this and other absorbent earths and alkalies, is subject to the laws of chemical composition, decomposition, and transfer—3. that common air does not enter into the same combinations as fixed air;—and lastly, he inferred from the general analogy of the effects of chemical attraction, that unsaturated affinity is the *form*, as Bacon would have termed it, of *causticity*. This brief, simple, and choice specimen of synthetico-analytical research, to that time unexampled in chemistry, he completed and crowned, by denoting the law of double decomposition as dependent on “*the sum of the forces,*” and fixing the place, not of magnesia only, which was as much as he at first contemplated, but of fixed air, side by side with the acids, in its own place in the *table of relative affinities*\*.

\* Essays and Obs. Phys. and Lit., vol. ii. pp. 221–24. The following description, by the French chemist De Lasône, in 1753, of the manner in which an aërial spirit is combined with lime and iron in the waters of Vichy, is worthy of notice, as a curious anticipation of truth since more exactly developed:—

“Toutes ces expériences prouvent évidemment que ces eaux sont alkales, par un principe salin et par une terre absorbante; qu’elles contiennent une matière ferrugineuse; qu’elles contiennent un principe spiritueux, composé non seulement d’un air sur-abondant, comme il s’en trouve dans quelques eaux, mais encore d’une portion de cette terre subtile dont nous venons de parler, jointe au principe huileux du bitume, et volatilisée par cet air, qui vrai-semblablement est le principal agent qui tient cette terre sus-

Black had certainly very little ambition, and apparently little of the activity of an ardent curiosity: for here he rested, after drawing from the facts before him some pregnant inferences, as to the production, for instance, of this fixed air from charcoal, and its diffusion through the atmosphere\*. He did not even measure, or collect, the air extricated in his experiments, still less did he try its density; he did not extend his inquiries at all into its elastic condition: and the consequence was that on *that point* which you take for the *stress* of his discovery, he rather retrograded from the inferences of his predecessors than advanced beyond them: for he went no further in his conclusions than this—"Quick-lime therefore does not attract air when in its most ordinary form, but is capable of being joined to one particular species only, which is dispersed through the atmosphere, *either in the shape of an exceedingly subtle powder*, or more probably in that of an elastic fluid†." He did more, it is true, than discover the chemical affinity of *one* substance only which floats in the air, or is fixed in many earths and alkalies; for that discovery, as it limited the *number* of such substances, so it extended to the rest the probability of a *like chemical constitution*: but whether these substances are or are not *elastic*, Black, like Daniel Bernoulli, declined to decide. The demonstration of this fact—that there exists *more than one species of elastic fluid* permanent at a common temperature and pressure when not acted upon by a condensing attraction—*was reserved for Cavendish*; being the consequence of that determination of its specific gravity of which you speak so slightly.—And here again, you see that in your haste you have denied this great philosopher his due.

And now that we have not only walked together over a part *pendue*, puisque lorsqu'on l'en chasse brusquement en secouant l'eau minérale, la terre se dépose aussi promptement, et qu'au contraire elle ne se dépose que très-lentement lorsque l'eau est bouché et que l'air ne s'évapore que lentement; que *ce même principe contient aussi une portion de la terre ferrugineuse qui existe dans ces eaux*, puisque lorsqu'elles sont dépouillées de leur air et qu'elles ont formé leur dépôt, on ne remarque plus aucun indice de matière ferrugineuse; qu'on doit encore à ce même air mêlé avec la terre et le bitume, et qu'on peut en cet état regarder, suivant la pensée de Lister, comme une espèce d'esprit, la saveur acidule qu'ont ces eaux à leur source et qu'elles perdent avec leur air sur-abondant; enfin que ce même principe aérien est la cause d'une partie de l'effervescence qui ces eaux font avec tous les acides."—*Hist. de l'Académie*, 1753, p. 174.

\* Black also ascertained that the peculiar matter of fixed air combines with other bodies in more proportions than one; and Cavendish, subsequently, that *that gas* combines in proportions of which one was about *double* the other,—a fact which proved of great importance to chemical theory.

† *Essays, Phys. and Lit.*, vol. ii. p. 198, 1765; *Experiments on Magnesia, &c.*, 1777.

of the demesne of experimental philosophy with more deliberation than your leisure seems usually to allow you, but even ventured on searching some of the inner chambers of the art of experiment, I must appeal to you, not in the style of arch solemnity with which your "illustrious colleague" addressed you in the chamber of the Institute, as having weighed the evidence in the case of Watt *versus* Cavendish—"Avec le scrupule en quelque sorte judiciaire qu'on pouvoit attendre de l'ancien *Lord Chancellor* de la Grande Bretagne\*,"—but I appeal to you, as ever you have learnt the laws of evidence from the only Chancellor of England who is of authority in philosophical questions, as ever you have listened to, and comprehended, that pupil of Bacon and Newton, the beauty of whose lectures you have so vividly described,—to take some shame to yourself, for having perused, by your own confession, the notes of Cavendish, without perceiving that all which I have said of the experiments of Black, as being so connected as clearly to manifest the whole train of the experimenter's thoughts, is still more clearly true of *these*.

You know what the problem was, on the investigation of which Cavendish was intent when he made the discovery in question. You know his aim to have been to find out what was become of "*the air lost*" in the combustion of hydrogen with common air. And what were the preliminary trials by which he searched for the lost gases? He tried—1. whether they were "*changed*" into carbonic acid;—2. whether they were "*changed*" into nitric acid;—3. whether they were *changed*

\* *Annuaire*, 1839, Note, p. 361. Lord Brougham, out of court, deals I fear as hastily with literature as with science; and *there* also sometimes makes the facts on which he reasons. Thus he criticises as "*unintelligible*" the condensed sense of that well-known line, in which Johnson, in his imitation of the Tenth Satire of Juvenal, speaks of "*patience*" as "sovereign o'er transmuted ill:" but he first makes it unintelligible, by substituting from his own poetical mint—"nature," where Johnson had written "*patience*." (Life of Johnson, p. 76.) Again, he animadverts severely on Johnson for "roaring out, 'No, Sir!'" in the presence of Hume, on being asked by a common friend to let him present the Historian to the Moralist" (Life of Hume, p. 223); and he adds, "above all we have a right to complain that the associate of Savage, the companion of his debauches, should have presumed to insult men of such *pure minds* as David Hume and Adam Smith, rudely *refusing to bear them company, but for an instant*." (Life of Johnson, p. 22.) It is curious to compare this with Johnson's own account: "I was but once in Hume's company; and then his only attempt at merriment consisted in his display of a drawing too indecently gross to have delighted even in a brothel." (Hawkins.) The *real man* from whom Johnson turned on his heel, was one who added to the moral *purity* of the school of Voltaire the garb of an *ecclesiastic*,—a circumstance which perhaps may abate something of Lord Brougham's indignation at the ill-manners of Johnson.

into sulphuric acid: he *negatived* by conclusive experiments these three suppositions of *condensation*: at this crisis of his inquiry Warltire burnt inflammable gas and common air in a vessel which he imagined to be close, and finding a very sensible loss of weight, concluded, with Scheele, that ponderable matter had passed through the vessel's pores in the form of heat: at the same time he repeated an observation, made also by others, that in the combustion *water* was deposited from the air, in which he supposed it *to have been contained*: to the mind of Cavendish, deeply meditating what might be the form of matter into which the *lost airs* could have been *condensed*, this inaccurate experiment immediately suggested the light for which he was watching. Such is his own description of the manner of the discovery; and the course of experiments recorded in his note-book leave no shadow of a doubt that he has described it with truth.

In the progress of these experiments, that is to say, in his *fourth* experiment of exploding the gases, in an apparatus like Warltire's but *really close*, on the 5th day of July 1781, Cavendish arrived at the *exact volume of hydrogen* which destroyed in combustion the whole of *its own elasticity*, with the whole elasticity of the *exact volume of oxygen found* to exist in the common air with which he exploded it: in the same experiment he ascertained that the vessel, which was of such capacity as to hold 24,000 grains of water, had lost scarcely one-fifth of a grain in weight: he then varied his apparatus in such a manner as to collect a sufficient quantity of the liquid formed in these experiments, for chemical examination, and before the end of the month demonstrated it to be *pure water*.

And now point out to me, if you can, in the whole range of experimental science *three facts* the ascertainment of which was more *obviously and indubitably conclusive* of the point in question? Is there any alternative left for scepticism? The total weight undiminished—three volumes of elastic matter gone—in its place pure water—could *any man* draw *any conclusion* from such experiments but *one*? could anything but *one foregone conclusion* have led a man to institute such a course of experiment? Does *the man* who *has* instituted, and made, such experiments, want any one to come to him two years afterwards with an idle *doctrine*, or *hypothesis*, as you call it, about the connection of *water* with some *undefined kind of phlogiston*, by way of explaining to him his own investigation? What is the use of a doctrine, or a hypothesis *after an inductive demonstration*, even if the hypothesis had had any real substance or distinctness of meaning? Are we to

deny the author of the demonstration the credit of understanding it, for no better reason than that in the private notes of his chain of proofs we find no shout of *εὕρηκα*?

I omit here all the multiplied precautions to ensure the most perfect accuracy in regard to every elementary material of these experiments—I omit the singular caution and sagacity which, on the unexpected intrusion of a minute quantity of nitric acid in one of his varied trials, induced Cavendish to wait till he had obtained evidence that *this* was the product of the *other ingredient* in atmospheric air, before he would publish his experiments: I put the question in a shape so simple that a child may understand it; and I ask you once more,—ought you not, with all this, clearly stated, before you, to feel some compunction for having admitted a suspicion of the good faith of Cavendish, or made a question of his having been the sole discoverer?

Again,—I have shown you, that though these experiments were communicated to Priestley as soon as they were made, and by Priestley mentioned to the public in express terms as —“*Mr. Cavendish’s experiment on the re-conversion of air into water,*” Priestley understood them no better than the communication which I have before mentioned of the discovery of nitrogen, and subsequently, with the aid of Watt’s opinion, concluded that “water by exposing it to heat in porous earthen vessels is capable of being converted into *respirable air* by the influence of heat:” I have shown you out of that very letter of Watt, communicated to the Royal Society, on which the only real question rests—whether he understood the consequences of Cavendish’s experiments nearly two years after they were finished—that Watt’s *doctrine* about water and *phlogiston* was built on *this false supposition*, and that he adhered to it after Priestley had communicated to him *that* experiment which was designed to be a repetition of Cavendish’s\*: I have shown you that in Priestley’s repetition the inflammable gas which he used cannot have contained more than one-fifteenth of its weight of hydrogen, and if it had proved anything, would have proved that water consists chiefly of *carbon*†: lastly, I have shown you that both Priestley and Watt were entirely ignorant of the distinction between *hydrogen* and the *inflammable gases on which they experimented and reasoned*; and until at a later time they were taught that distinction by Cavendish, and thus learnt what the *real basis* of water is—were obviously as incompetent to *understand*, as to *discover* its composition‡.

\* Report of the British Association, Postscript to Address, p. 24.

† Ibid. p. 27.

‡ Ibid. p. 25.

With these things before the world, you even now venture to reiterate, as your final conclusion, this most unjustifiable judgement—"It is undeniable that from less elaborate experiments Mr. Watt had before Cavendish drawn the inference then so startling, that it required all the boldness of the philosophic character to venture upon it,—the inference that water was not a simple element, but a combination of oxygen with *inflammable air thence called hydrogen gas*. That Mr. Watt first generalised the facts so as to arrive at this great truth, I think has been proved as clearly as any position in the history of physical science. It is equally certain from the examination of Mr. Cavendish's papers, and from the publication lately made of his journals, first, that he never so clearly as Mr. Watt drew the inference from his experiments; and secondly, that though those experiments were made before Mr. Watt's inferences, yet Mr. Cavendish's conclusion was not drawn privately even by himself till after Mr. Watt's inference had been made known to many others."—!!!

What friend of yours, my dear Lord, but must regret to see a great man trifling with his own reputation by interfering in subjects of which he thus betrays but too superficial a knowledge? I sincerely lament, for my own part, that having once been honoured by your regard, and having always respected your talents, it should have fallen to me to presume in this manner to rectify your misapprehensions. I declined to enter into controversy with *you*, partly for old acquaintance sake, and partly because I thought you on this question less *responsible* than the official writer of the Institute of France. But you *would* do battle with me; and your weapons were none of the fairest: for instead of replying to my arguments, you did me the injustice, without provocation, to compare the abilities of the obscurest lover of science in England with one of the most eminent of its cultivators in France. I know not that I shall even now have convinced you that the meanest of our philosophical chemists, in his own art, and in a just cause, may be more than a match for the most learned judge of *Patents*, or even for the ablest member of the "*Institut*," whose studies have lain in another direction. A judge in a patent cause may see his way well enough, no doubt, through intricate scientific questions, if he is but prudent in his selection of authorities: but I do not perceive that in this case you have abided by any authority better than *your own in 1803\**. You are even bold enough, on the strength of such authority, to differ from a deceased Secretary of the "*Institut*" itself, than whom few men were bet-

\* "I first stated that opinion in a published form in 1803-04," Edinburgh Review, vol. iii. Life of Lavoisier, p. 253.

ter acquainted with sciences not peculiarly his own: but though the subject is chemistry, though *you* have attended Black's lectures, and though Black's own discoveries are in question, I greatly fear that on almost every point in which you differ from Cuvier you are yourself in the wrong.

Thus, you are certainly wrong in *denying* Cuvier's assertion—that permanently elastic fluids were *measured* by Hales; and you have only to consult the 'Analysis of the Air,' to be convinced of your mistake.

Again, you are wrong in *denying* Cuvier's assertion, that "no one before Cavendish had distinguished fixed air as a separate *aëriiform* substance:" and you need only look at Black's treatise to assure yourself that he declined to decide, for want of evidence, whether it was an *aëriiform* substance, or not; and left it among the class of—"bodies of which it is difficult to say, whether they are really *combined* with the *aërial* particles, or are merely *suspended* in the fluid, in consequence of their being of the *same specific gravity*\*."

But above all, you are *most wrong* in reprehending the former Secretary of the Institute, for "*making no mention whatever of Watt in connection with the discovery of the composition of water*"—for not confounding, that is, the rights of discovery—for not falsifying the history of chemistry in one of its most material parts—for not representing Watt as the claimant of a merit to which he had not the smallest pretensions—and thus degrading, with intent to exalt, the venerable name of one who has entitled himself to the admiring gratitude of ages, by realising, beyond any other man, the vision which Bacon saw—of experimental *works of fruit*.

You have no sufficient ground, I think, for imputing to—"a person of M. Cuvier's eminent attainments, filling the high office of '*Secrétaire perpétuel*,' and charged with the delicate and important duty of recording the history of science yearly"—that "he has not read Mr. Cavendish's paper †, or Dr.

\* Cavallo on Air, p. 361. 1781. Hawksbee approached the nearest to the discovery of the different density of gases, as early as 1707;—"whether," said he, "the space deserted by the water [after an explosion of gunpowder in a close vessel] is possessed by a body of the *same weight and density*, or is of the same quality, as common air, I dare not determine; *since an experiment I have lately made seems to conclude it otherwise*." He observed likewise "a loss, or absorption of this air, after it had reached its former temperature;" and suggested that a temporary distension of the springs or constituent parts, of the ambient air, as well as of those contained in the body of the gunpowder, may account for "*this odd phænomenon*."—*Phil. Trans.* vol. xxv. p. 2409.

† One of Lord Brougham's reasons for thinking that Cuvier had never read Cavendish's paper is, that he says,—"*Cavendish unfolded his discoveries in a manner even more striking than the discoveries themselves*"—

Black's treatise." And certainly you have no ground to "lament," with respect to him, "that the history of science should be written with such *remarkable carelessness* and such *manifest inattention to the facts*,"—however true it may be, were the censure justly pointed—"that to find mistakes so very gross in the works of ordinary writers might excite little surprise; but when they are embodied in the history of the *National Institute*, and when they come to us under the name, among the very first in all sciences, of Cuvier, we may at once wonder and mourn\*."

I only trust that the stone which you have so rashly cast at Cuvier will not recoil on any other head. I still trust sincerely that so severe a reproof will not *permanently* rest on the *present* "Perpetual Secretary" of the Institute of France; and that conformably to the known manliness of his character, and clearness of his understanding, M. Arago will yet rectify, as he knows how, the inadvertence into which he has fallen.

It now only remains for me to remark on your last words in reply to one who has supported with far greater ability than myself the same opinions which I have expressed.

I have known you, my dear Lord, more strenuously and skilfully employed than in deciding these questions for chemists; and think I remember it to have been one of the arts of a dexterous *advocate*, with which you were then familiar, to speak somewhat *largely* in an opening speech, of evidence which yet it might not be discreet to bring into court: and so I suppose it is now; for in animadverting upon the ignorance of this enemy in ambush, whom however you seem to suspect of being no ordinary man, I perceive you affirm, that you "have lying before you *fifteen pages* of statements of chemical errors in the thirty-four pages of his paper, and as these corrections are the work of a *most experienced, learned and practical chemist* whom you consulted, you have entire reliance on his report and opinion." It was some disappointment to me, at first, to find that you kept the *fifteen pages* in your pocket; but I remembered, how it happened not unfrequently of old, that in the torrent of that forensic eloquence which

an assertion which will scarcely be disputed by any competent judge who compares the brief perspicuity of expression, and the select sequence of most exact experiment, which shines in every page of Cavendish, with the rambling, inconsequent manner of thinking and writing, general in his time, and I fear not infrequent in our own. Lord Brougham also accuses Cuvier of stating, that Cavendish established in his paper of 1764 these propositions—"l'eau n'est pas un élément; il existe plusieurs sortes d'air essentiellement différentes." But is not 'l'eau', in this paragraph, merely a misprint for 'l'air'?

\* Brougham's Lives, vol. ii. p. 507.



so often dazzled and delighted your hearers, something that should have been kept back would occasionally slip out, of which an astute adversary did not fail to make his advantage. And even so it is still: from the *fifteen* critical pages you have allowed *one* criticism to creep out, as too good to be suppressed. And here it is:—

“I leave him” (the author of this heap of errors) “in the hands of M. Arago, who will observe with some wonder that he has been accused, and judged, and condemned, by a chemist so well-versed in that science, and so reflecting, as to announce the astonishing novelty—that the exhibition of sulphur to sulphuric acid reduces that acid, and restores it to its primitive state of sulphur! The writer had probably read somewhere that sulphuric acid is reduced to sulphurous by the process; for he is assuredly the first that had ever hit upon the acid’s reduction by sulphur to ‘its primitive state’\*.”

Now we will at least give credit to the *present* perpetual Secretary of the Institute, to whose scorn you devote the unhappy Reviewer, for having read the papers of Cavendish; and he would no doubt recollect this remarkable passage in the “experiments on factitious airs” (1764) to which the Reviewer should seem to be referring—“Sulphur is allowed by chemists to consist of the plain vitriolic acid united to phlogiston; the volatile sulphureous acid appears to consist of the same acid united to a less proportion of phlogiston than what is required to form sulphur; a circumstance which I think shows the truth of this is, that if oil of vitriol be distilled from sulphur, the liquor which comes over will be the volatile sulphureous acid.” M. Arago might perhaps compare these early notions of Cavendish with the Reviewer’s account of the phlogistic opinions, not in your interpolated words, but in his own—“It was concluded therefore that it was the *same* phlogiston which was derived from all those substances (charcoal, sugar, metallic bodies, &c.), however different in their nature: a similar succession of phænomena is presented by sulphur: if it be burnt, it forms sulphuric acid; but if the acid thus formed be heated with *phosphorus*, or *charcoal*, or *sugar*, or even *sulphur* itself, it is equally restored to its primitive state †,”—and having read this account, supposing M. Arago for a moment to be only as experienced, as learned, and as practical a chemist, as he whom you have consulted out of court, and no more—supposing him, that is, to believe, with your anonymous friend and yourself, that the total reduction of sulphuric acid by sulphur is a laughable absurdity—M. Arago would yet see,

\* Brougham’s Lives, vol. ii. p. 511.

† Quarterly Review, Dec. 1845, p. 106.

that there is nothing laughable, or ignorant, in the statement of the Reviewer; though in strictness of language it might have been more correct to say, that—‘sulphuric acid heated with *phosphorus*, or *charcoal*, or *sugar*, is reduced to its *primitive* state, and even heated with *sulphur* is reduced to its *previous* state of sulphurous acid’: and I think M. Arago might *wonder* a little at finding how much you, and your learned, experienced, and practical friend, have made of a mere *verbal* slip.

But *if*, as I am apt to suspect, the *perpetual Secretary* is a *better* chemist than yourself, or knows better, this time at least, on whose *information* to rely, then he will whisper to you privately, that sulphuric acid *really is reduced*, astonishing novelty as it seems to you, even by *sulphur itself*: and he will doubtless proceed to explain to you how this marvel comes to pass: he will remind you, that the great chemist of our time whose life you have written, when attempting to decompose sulphur, found it so closely united with a very considerable quantity of hydrogen, that he remained for some time in the belief, that he had effected its decomposition; and M. Arago will add, that since *hydrogen* decomposes *sulphurous acid*, it follows, that sulphuric acid cannot but be reduced by sulphur, in all the forms in which sulphur is commonly experimented with, to its *primitive* state, and that the Reviewer therefore is literally right.

And now, retaining the very sincere respect which I have always felt for one who has so laudably devoted the leisure hours of a busy life of public service to the promotion of literature and science, I hope I may have persuaded you, that it is at once more safe, and more just, for those who have not had leisure to pursue chemical studies to their foundation, to leave chemistry and chemists to themselves—at least so far as regards the minutiae of the science, and arbitrations of the rights of discovery; and I take the license of old acquaintance to advise, that if you *will* venture on such dangerous ground, you should at least learn how to choose your authorities; and when you find even Robison, and Watt, deserting you, and *the perpetual Secretary* so tardy in coming to the rescue, you should not think it enough to reflect that in 1803—1839—1845 and 1846—you yourself stated and re-stated an opinion contrary to the public\* voice of the chemists of England.

I have the honour to remain, with undiminished regard,

My dear Lord, your faithful Servant,

W. VERNON HARCOURT.

N.B. An oversight with respect to a date in the part of

\* Lord Brougham expresses more surprise than is just, that I take no

this letter previously published (p. 116) requires to be thus corrected. "Priestley addressed this paper to the Royal Society on the 21st of April 1783: and therefore the communication of Cavendish's experiments, acknowledged in it as having suggested his own, must have been prior to the speculations founded thereon which Watt addressed to Priestley on the 26th of the same month, as well as to Lavoisier's experiments which followed in June."

---

LXXIX. *Observations on Mr. Strickland's Article on the Structural Relations of Organized Beings.* By Prof. OWEN, F.R.S.

*To the Editors of the Philosophical Magazine and Journal.*

GENTLEMEN,

THE author of the interesting paper "On the Structural Relations of Organized Beings," in your last Number, appears,—in recommending the introduction of "the adjective *affine* or *homologous* in place of *analogous*, when referring to structures which essentially correspond in different organic beings" (p. 358),—not to have been aware that the term 'homologous' had been used in the sense he recommends, by comparative anatomists both in this country and abroad for some years past.

In the article *Marsupialia*, for example, Cyclopædia of Anatomy, part 21, April 1841, p. 283, he will find—"With reference to the interesting question,—What is the *homology* or essential nature of the *ossa marsupialia*?"—and their homologies discussed. In No. XXII. of the same Cyclopædia, article *Monotremata*, p. 375: "The interposed cartilages, which thus form a third element in the costal arch, repeat a structure common in Crocodiles, and may be regarded as the *homologues* of the costal appendages in the ribs of birds." And

notice of his having *quoted* a private letter to the son of Mr. Watt on the subject of his father's claims. I am aware that Lord Brougham says he has seen such a letter, and says also that the opinion expressed in it respecting Watt's MSS. is different from the opinion attributed to Dr. Henry by me: but I am not aware that Lord Brougham has given *any quotation* from this letter; nor if he had, would any *partial quotation* have satisfied me, that Dr. Henry's opinion was at any time different from that which he expressed to me, when I mentioned to him the sentiments which I had heard fall from M. Arago concerning the MSS. at Aston, and the insincerity of Cavendish. Dr. Henry then said, that he had seen nothing in those MSS. either to justify that impression, or to alter the received opinion respecting the discovery of the composition of water. Who indeed can doubt but that the MSS., had they contained any evidence to support an object which has been so long urged by private solicitation, would have been made public long ago?

again, p. 377:—"These clavicles are the *homologues* of the os furcatorium in the Bird." And in the note, same page:—"For a full and elaborate discussion of the various opinions which have been offered respecting the *homology* or signification of the complicated apparatus of the shoulder," &c. I could easily multiply instances in which the term *homology* has been applied, both substantively and adjectively, in the sense recommended by Mr. Strickland. I have been in the habit of defining, in my Introductory Lecture, the terms *homology* and *analogy*, as in the Glossary appended to the Lectures on Invertebrata, published in 1843, and of illustrating their meaning in comparative anatomy, by reference to the skeletons of the *Bird* and the *Draco volans*. The fore-limb of the Dragon being composed of essentially the same parts as the wing of the Bird, is *homologous* with it; but the wing or parachute of the Dragon, having a similar relation of function, is *analogous* to the wing of the Bird. But in thus illustrating the term *homology*, I have always felt and stated that I was merely making known the meaning of a term introduced into comparative anatomy long ago, and habitually used in the writings of the philosophical anatomists of Germany and France. Geoffroy St. Hilaire also, in defining the term, acknowledges its source:—"Les organes sont *homologues* comme s'exprimerait la Philosophie Allemande; c'est à dire qu'ils sont *analogues* dans leur mode de développement," &c.—*Annales des Sciences*, tom. vi. 1825, p. 341.

I have gone perhaps a little further than Oken and Geoffroy in defining the different kinds of 'homology,' which appear to me to be three, viz. 'general,' 'serial,' and 'special.' General homology is the relation in which a part or series of parts stands to the ideal or fundamental type; and thus, when the basilar process of the occipital bone in Anthropotomy is said to be the 'centrum' or 'body of the last cranial vertebra,' its *general* homology is enunciated. When it is said to repeat, in its vertebra or natural segment of the skeleton, the body of the sphenoid bone, the body of the atlas, and the succeeding vertebral bodies or centnums, its *serial* homology is indicated. When the essential correspondence of the basilar process of the occipital bone in Man with the distinct bone called 'basi-occipital' in a crocodile or a fish is shown, its *special* homology is determined.

Vicq d'Azyr began the study of 'serial homologies' in his ingenious memoir on the parallelism of the fore and hind limbs, in the Memoirs of the French Academy, 1788.

'Homologous parts' are, indeed, in one sense, 'analogous parts,' having like relations, as being repetitions of the same

parts of the body, to different animals; but I have been in the habit for some years past, of expressing this kind of analogy by the term 'homology;' and I heartily join in the recommendation of your ingenious correspondent, that *all* writers on comparative anatomy and zoology should use the word in that sense, whether it be or be not coupled with the likeness of function performed by such parts; to signify which relation alone, the term 'analogy' should be restricted. As instances of parts both homologous and analogous, may be cited the pectoral limb of the Porpoise and that of the Fish: they are homologous as being constituted of essentially the same or corresponding parts; they are analogous as having the same relation of subserviency to swimming. So likewise the pectoral fin of the flying-fish is analogous to the wing of the bird; but, unlike the wing of the Dragon, it is also homologous with it. Some organs are analogous, but only partially homologous: thus the Monkey's foot is analogous to the Man's hand, as having the functions of the opposable thumb: it is also homologous with it *generally*, as being part of the radiated appendage of a hæmal arch; and *serially* as being the terminal segment of that appendage; but it is not *especially* homologous with it. The thumbless hand of *Ateles* is especially homologous with the perfect hand of Man, but the pollicate foot of *Ateles* is not so. I offer these as examples of the mode in which 'homology\*' is illustrated in my own Lectures on Comparative Anatomy, in addition to those cited from my printed works.

I am, Gentlemen,

Your most obedient Servant,

College of Surgeons,  
May 9, 1846.

RICHARD OWEN.

LXXX. *Observations on Messrs. Lyon Playfair and Joule's Memoir on Atomic Volume and Specific Gravity. By Prof MARIGNAC of Geneva* †.

THE authors in this paper have determined the density of a large number of bodies, and have arrived, by a comparison of their atomic volumes, at laws which would be rather curious if they could be regarded as proved. Their investigations have been principally directed to the soluble salts, and they have sought to determine not only the atomic volume of

\* In geometry those sides of similar figures which are opposite to equal and corresponding angles are sometimes said to be *homologous*, as being proportional to each other.

† Translated from the *Bibliothèque Universelle*. Feb. 15, 1846. The memoir referred to will be found at p. 453 of the previous volume of this Journal. [ED. *Phil. Mag.*]

these salts in the solid state, but also their volume in the state of solution, that is to say, the increase in volume which the water in which they are dissolved undergoes. Their method consists in introducing a determined weight of salt into water contained in a glass bulb, capable of holding 1000 to 4000 grains of water, surmounted by a graduated stem, in which they measure the increase in volume of the water after the solution of the salt. The same apparatus served to measure the volume of the salts in the solid state, either by employing a saturated solution of the salts or a liquid in which they were insoluble, such as the oil of turpentine.

We shall examine successively the results relative to the volumes occupied by the salts in a state of solution and in a solid state.

In 1840 Dalton observed that sugar and certain hydrated salts, on solution in water, increased its volume by a quantity precisely equal to the volume of water they themselves contained. He generalised this observation, extending it to all the hydrated salts, and he thence concluded that the anhydrous salts did not increase the volume of the water in which they were dissolved.

Messrs. Playfair and Joule confirmed this interesting result for a sufficiently large number of salts, but not for all\*. Those which obey this law are in general the salts containing a large proportion of water of crystallization, as for instance—

The sulphates of the magnesian group with .....	5, 6 or 7 equiv. water
The chlorides of calcium, strontium and magnesium with...	6 equiv. water
The alums † with .....	24 equiv. water
The phosphates or arseniates of soda, neutral or basic, with	24 equiv. water
The carbonate, borate, sulphate, pyrophosphate of soda with	10 equiv. water
The sulphate of alumina with.....	18 equiv. water

Lastly, cane-sugar, considering as water the 11 equiv. oxygen and hydrogen which it contains.

For all these compounds the increase in volume of the water in which they are dissolved is precisely equal to the volume of water they contain; so that if these salts be used in the anhydrous state, they dissolve in the water without causing any change of volume.

Several other salts, both anhydrous and hydrated, follow different laws. Messrs. Playfair and Joule advance the fol-

\* This had been previously done to a certain extent by Mr. Holker; see his paper in vol. xxvii. p. 207 of this Journal.—ED. PHIL. MAG.

† There is however an exception, for the ammoniacal alums possess a volume equal to 25 equiv. water and not 24 like the potash alums, although they contain like them only 24 equiv. water of crystallization.

lowing law in respect to them:—The volume occupied by an equivalent of any salt whatever in solution in water, is always an exact multiple of the number 9 representing the atomic volume of water.

It is difficult to conceive whence this simple relation between the volume of the salts and the volume of the water arises; nevertheless, if by adopting this hypothesis we were led to represent by similar formulæ the volume of analogous compounds this law would be interesting, but the following examples will suffice to show how many anomalies we meet with. We will here compare the number of volumes of water which some groups of analogous compounds occupy.

Bisulphate of soda . . .	2	Nitrate of copper . . .	2
... of potash . . .	4	... of soda . . .	3
... of ammonia . . .	5	... of potash . . .	4
Bicarbonate of soda . . .	2	... of ammonia . . .	5
... of potash . . .	4	Chloride of potassium . . .	3
... of ammonia . . .	4	Iodide of potassium . . .	5

But let us pass over these objections and see whether really the circumstance indicated as fact is sufficiently established by experiment. The atomic volume of a salt in solution is not a constant magnitude; it varies with the temperature and with the relative proportions of the salt and of the water; we will examine successively these two influences. That of the temperature is very considerable; this is proved by the experiments of Messrs. Playfair and Joule, who measured the volume occupied by a salt at different temperatures. The following are some of their results:—

	Volume of one equivalent of salt in solution.		
	at 0°	60·88	at 29° 63
Sulphate of magnesia with 7 equiv. water	at	0	56·12
... of zinc	...	0	61
... of iron	...	2 $\frac{3}{4}$	14·4
Anhydrous sulphate of potash	...	0	65·2
Sulphate of potash and copper	...	0	61·8
... of potash and magnesia	...		

Now at what temperature should these volumes be compared with one another? We are totally unable to answer this question, but certainly nothing authorizes us to choose for each salt a different and arbitrary temperature, as Messrs. Playfair and Joule have done, by taking only the numbers inscribed in the second column, because these alone satisfied the law which they wanted to prove. For other bodies, however, they have admitted experiments made at low temperatures; thus for the alums of iron and chromium the experiments were made at 2 $\frac{3}{4}$ °, for the sulphate of alumina at 10°.

When it is seen that the atomic volume of sulphate of potash and magnesia may vary more than 10 between  $0^{\circ}$  and  $27^{\circ}$ , it will readily be conceived that it will never be a difficult task to make the volume of each compound an exact multiple of 9; for this purpose it is merely requisite to make the experiment at a suitable temperature.

The relative proportions of water and of salt likewise cause the volume occupied by the latter in solution to vary. Messrs. Playfair and Joule are fully aware of this; they made experiments on the very subject, and proved this influence by the following results relative to the volume occupied by an equivalent of sugar in solution in different quantities of water:—

Relative proportions of sugar and water.	Temperature.	Volume of one equivalent of sugar in solution.
1 : 120	$15^{\circ}\cdot5$	99
1 : 10	11	105·89
1 : 1	11	107·01
3 : 1	11	108·06

The difference between the first number and the following would have been still greater if the experiments had been made at the same temperature.

What then are the relative proportions of salt and water which should be employed in order that the results may be capable of comparison? We know not, and the authors leave us in total ignorance of the subject; we only find that in their experiments they have not restricted themselves to uniform conditions; thus they employed

	Salt.	Water.	Proportion.
for the sulphate of alumina and potash .....	59	1000	1 : 17
... .. of iron and ammonia .....	30·06	1000	1 : 33
... .. of oxide of chrome and potash ...	32	4100	1 : 129
... .. of alumina and ammonia.....	20	4100	1 : 205

Evidently, if under such circumstances they have found an agreement between the volumes of these salts, it may be concluded that had they operated under uniform conditions they would not have found the least trace of one. In short, the atomic volume of a salt in solution depends both on the temperature and the relative proportions of water and salt, and these circumstances cause the volume to vary within considerable limits. As long as these influences are neglected, or we do not operate so that they act in all cases in the same manner, the results obtained cannot be compared in any possible way with one another.

Let us now pass to the atomic volume of the salts in the solid state. Messrs. Playfair and Joule have advanced for these the following law:—The atomic volume of any salt



whatever (anhydrous or hydrated) is a multiple of 11, or of a number near to 11, or a multiple of 9·8 (the atomic volume of ice); or again, the sum of a multiple of 11 or of 9·8.

This law appears to us to resemble very much the preceding, except that the indecision as to the choice between the multiples of two different numbers renders it still less probable.

We do not in this case meet in the same degree with the objections above set forth for the case of dissolved salts; the temperature cannot cause any great variation in their density, and the experiments were made at temperatures varying too little to have any separate influence,—if it be admitted, which however is far from being proved, that with respect to the solid bodies their densities should be compared at the same temperatures.

We will however make one remark relative to the process by which the densities were determined; it appears to us little suited to give accurate results. It is not stated what was the volume of the liquid to which the salt whose density was to be determined was added; but as it was the same apparatus which had served for the preceding experiments, we may suppose that it contained at least 1000 grains of water. The quantity of salt employed in each experiment was from 40 to 60 grains, and there thence resulted an increase in the volume of the liquid corresponding to about 20 to 40 grains of water; in a great number of cases indeed we find an increase of only 10 to 20 grains, that is to say, of from 1 to 2 per cent. of the total volume. It is evident that by this process it is extremely difficult to avoid serious errors produced by the slightest variations of temperature, which tend to alter the volume of so large a liquid mass, and of errors probably still more important, which might result either from the solution of a portion of the salt in the liquid, if this was not accurately saturated, or from the precipitation of a portion of the salt contained in the liquid, if it were more than saturated. The experiments of Gay-Lussac prove indeed that both these circumstances may readily occur.

These causes of error might perhaps be neglected if the volume of the liquid were very inconsiderable; but when, on the contrary, its proportion is so large relatively to the solid salt, they become too serious for any confidence to be placed on the densities obtained by this process.

We should add, that on reviewing the formulæ which Messrs. Playfair and Joule have established, based on the preceding law, they do not appear to us to indicate any very great probability for this law. Along with certain analogies which

do not surprise us, for they result simply from the fact of the equality of the atomic volumes with respect to the isomorphous compounds, we meet with a number of most striking anomalies. For the chlorides of calcium, strontium and magnesium, the atomic volume is equal to 11 multiplied by 6, *i. e.* the number of equivalents of water of crystallization of those salts, but for the alums it is  $11 \times 25$ , while there are only 24 equiv. water. For the sulphate and borate of soda with 10 equiv. water the volume =  $11 \times 10$ , but for the pyrophosphate with 10 equiv. water it is  $11 \times 11$ ; and for the carbonate likewise, with 10 equiv. water, it is  $9.8 \times 10$ ; for the anhydrous carbonate of soda the factor 11 is taken, and for the hydrated carbonate 9.8; on the contrary, for the anhydrous sulphate of soda the authors prefer 9.8, and for the hydrated sulphate 11. The bromide of potassium =  $4 \times 11$ , the bromide of sodium =  $5 \times 11$ , the chloride of potassium =  $4 \times 9.8$ , the chloride of sodium =  $3 \times 9.8$ .

These instances we think will suffice to show that the hypothesis of Messrs. Playfair and Joule is not confirmed by an analogy of formulæ such as ought to be expected, and that the coincidence which does exist between the calculated and the observed densities merely result from the easy way in which the authors select at will the factor 9.8 or the factor 11, or even of combining them for one and the same body, as they have done in a large number of cases.

LXXXI. *Remarks on Dr. Faraday's Paper on Ray-vibrations.*  
By G. B. AIRY, Esq., *Astronomer Royal.*

*To the Editors of the Philosophical Magazine and Journal.*

GENTLEMEN,

THE communication which accompanies this was sketched before my attention was called to Dr. Faraday's leading paper in your Number for the present month. I need not to say that I read that paper with great interest and great pleasure. Yet I will ask your indulgence, and I am sure that I shall receive the forgiveness of Dr. Faraday, while I comment on the principal points of that paper somewhat critically. I am desirous of examining, or of suggesting grounds for examination by others, as to how far the fundamental suppositions of Dr. Faraday are necessarily limited by recognised phænomena, and as to how far the subject is metaphysical or physical.

The paper, as I understand, treats of two subjects:—

1. The possibility of explaining phænomena of radiation,

more especially of light, by supposing that when there is no body obviously occupying the path of the light, &c., the vibrations which are assumed as the foundation of the undulations producing the phænomena are transmitted on the *lines of force* by what (for want of a received term) may be called *lateral shakes*.

2. The possibility of removing the idea of *substance* and substituting for it that of *centres of force*.

I shall treat of these in the order in which I have written them above.

1. With regard to the transmission of light through the planetary spaces.

Dr. Faraday and myself agree in receiving the undulatory theory of light with transversal vibrations, as applicable to those phænomena which present themselves in ordinary optical experiments. Without any wish therefore to dogmatize on this matter, I shall assume the undulatory theory in all the following remarks.

It is admitted that vibrations forming progressive undulations are required for the explanation of certain crystalline and other phænomena. But I must claim somewhat more. Progressive undulations (leaving the nature of their vibrations undetermined) are required to explain the phænomena of *diffraction*; and these progressive undulations must not be of the nature of radial shakes, where each shake derives its virtue or existence from the momentary influence of the distant origin, but they must be true waves, of which the mechanical characteristic is that the motion of a succeeding set of particles is determined by the relative motion of the preceding set of particles; the order of "preceding" and "succeeding" not being confined to a radial line or to any lines whatever, but being such that the motion of particles may be origin of motion to other particles extending round them through a very large solid angle. I defy any one to put together a theory of radial lines subject to lateral shakes which shall explain diffraction; and I say that it will be found absolutely necessary to admit, in the theory explanatory of diffraction, that each disturbance of particles produces a swell (to use language derived from the motion of water), which swell is propagated in all directions through at least a very large solid angle. Now the consequences of this are very important. Diffraction takes place in air; therefore the vibrating medium exists in air, and the undulations are transmitted by *it*, and not by radial shakes. As far as we can perceive air in its utmost degree of tenuity, it produces *refraction*; refraction inexorably requires for its explanation a

change of velocity of the undulations, and a power at the same time of changing the direction of progress in a degree exactly corresponding to the change of velocity: these changes are in the simplest and most natural manner possible explained by the theory of true waves in which the swell produced by every particle is propagated in all directions through a very large angle, while (as I apprehend) it will be found somewhat difficult to modify a theory of radial shakes so as to explain them; therefore I conceive it demonstrated that the propagation of true waves takes place through the air to its utmost borders. Beyond the existence of sensible air we can make no experiments; and I am free to concede that if we supposed the air and its accompanying æther [if different] to terminate at a distinct frontier, and if we supposed the transversal shakes to be propagated radially through the planetary spaces to that frontier, and then supposed each shake, as it presented itself, to be the origin of a spreading swell through the æther, the phænomena of light would be explained. But here a remarkable circumstance forces itself on our minds. A moment's consideration will show that at this frontier the course of the light will be subject to refraction, in just the same way as if the incident light had consisted of waves, and following the same law as depending on the velocity of propagation. Now it is abundantly established that at the boundary of our air there is no sensible refraction, that is, that the velocity of the propagation is not sensibly altered. Now is it not a very curious circumstance that there should be a system of radial shakes outside and a system of true waves inside which propagate the undulations with *exactly* the same velocity? Is there any philosopher who would be inclined to receive as true this suggestion of two independent causes of velocity, and this exact adjustment of independent velocities, when the adjustment will *necessarily* exist if the same vibrating medium or æther occupies all space? Not I, certainly. However well-disposed I might be to admit any such *saltus* of nature at the surface of glass or crystal where the phænomena of light are totally changed, I cannot bring myself to believe in it as existing either *through the air* where the change of phænomena is gradual, or *at the limits of the air* where there is no change at all. In a word, I must have the same theory of light for the planetary spaces as for the air in which our experiments on diffraction are made; and that theory must be the theory of true waves.

I do not insist on the novelty of the conception, that lateral influences take place in a travelling succession along a radial line, in a manner different from anything whatever that we

know, I am perfectly aware that the theory is merely sketched by Dr. Faraday as the result of hasty thought, and that it might be in some measure modified in its details on further consideration by its author. But while the distinctive features of the theory are retained, it will be, for the reasons which I have given, inadmissible to me. The theory is however, in my opinion, a fair subject for the consideration of the natural philosopher.

2. With regard to the substitution of centres of force for matter.

This speculation, in its general character, differs little from the celebrated inquiry regarding Substance and Accidents. In the latter the question is, whether, when we have found a lump of matter to possess certain form, colour, weight, and other properties, we can satisfy ourselves by saying that this lump of matter is a combination of such a form, such a colour, such a weight, &c.? And the answer has usually been that the mind is not satisfied unless we describe the lump of matter as *something* possessing the properties of such a form, such a colour, such a weight, &c. In the speculation before us, the question is, whether instead of matter which exerts certain actions upon other matter, we may assume that there is nothing but a number of centres of force producing these actions? I think that most persons would say that the mind is not satisfied with this assumption, and that it requires the idea of a *something* as foundation for these centres of force. But this question, in my opinion, is purely a metaphysical question, entirely removed from the province of the natural philosopher.

To a great extent I am willing to admit that the supposition of centres of force is satisfactory. Mechanical attraction or repulsion (including weight under the former term), colour, radiation of every kind where the existence of something intermediate between the radiating body and the body receiving the radiation is not apparently demonstrated; all these may, I think, be received without scruple as the results of mere centres of force. But there is one property, to which by chance Dr. Faraday has not alluded in his paper, that appears to me irreconcilable with the notion of centres of force; I mean the property of *inertia*. And I believe that the general notion of *substance* is really founded upon the perception of *inertia*. Construct for any one a mass of matter possessing invariable form, colour, and other attributes, even attraction; if he finds that this mass yields to muscular or other force without perceptible resistance (it matters not whether it continually retain the same velocity or not), he will

scarcely scruple to admit that there is no *substance*. While the resistance to force remains, it seems scarcely possible to get rid of the idea of *substance*.

Perhaps it may be said that even inertia may be represented by centres of force, only supposing the development of the force to be dependent in some way upon time. Such, however, is not the character of forces that we know best; and the introduction of this idea appears to give greater complexity to the force-centre-theory than is given by the idea of substance in the material theory.

Now I say that, in the wave-theory of light, and in all theories of waves where the amplitude of the vibrations does not diminish transcendently with relation to the distance passed over by the wave, the supposition of inertia (or something equivalent) is absolutely necessary. This will be evident to any mathematician who compares the results obtained from the different suppositions of inertia or no inertia. For instance; in the theory of the transmission of heat by conduction, no inertia is supposed; the equation then has the form  $\frac{dh}{dt} = A \cdot \frac{d^2h}{dx^2}$ , of which the solution (supposed to be periodic) is,  $h = B \cdot \epsilon^{-ax} \cdot \cos(nt - \beta x)$ . But in the theory of the transmission of sound, where the vibrating particles are supposed to possess inertia, the equation is  $\frac{d^2X}{dt^2} = A \cdot \frac{d^2X}{dx^2}$ , of which the solution (similarly restricted) is  $X = B \cdot \cos(nt - \beta x)$ . The former result certainly does not represent anything like the law of diminution of light; the latter does represent its general constancy of intensity (the distance of the source being very great). I infer therefore that the supposition of inertia is absolutely necessary.

Combining this inference with that obtained above regarding the universality of undulations in space, I am led to the conclusion that all space with which we are acquainted contains something which exhibits the property that we call *inertia*. The reasons which have led me to this conclusion appear to me decisive, but I admit them to be fair subjects for doubt and discussion by natural philosophers. Whether we are to infer from this that there is *matter* through all space, is, in my opinion, a metaphysical question.

But the remarks that I have just made will enable me to answer one paragraph of Dr. Faraday's paper. "Perhaps I am in error in thinking the idea generally formed of the æther is that its nuclei are almost infinitely small, and that such force as it has, namely its elasticity, is almost infinitely

intense. But if such be the received notion, what then is left in the æther but force or centres of force?" To this I reply, that *almost infinitely* has no meaning but *finitely*, and therefore that the supposed æther, under this description, is precisely in the same category as all other fluids. But I add, in regard to the latter sentence, that the mathematical considerations which I have detailed above, show that there is something in the æther besides force or centres of force, namely *inertia*. And I repeat the expression of my own opinion, that it is easier to conceive this as indicating *substance* (however obscure the idea may be), than to frame a system of laws applying to centres of force which shall represent its effects equally well.

I am, Gentlemen,

Royal Observatory, Greenwich,  
May 12, 1846.

Your obedient Servant,  
G. B. AIRY.

---

LXXXII. *Explanation of the Vorticose Movement, assumed to accompany Earthquakes.* By ROBERT MALLETT, C.E., M.R.I.A., Ph.D., &c., Secretary of the Geological Society of Dublin\*.

IN our progress to the ascertainment of physical knowledge, the removal of error is next in importance to the discovery of that which is true, inasmuch as by the former the road is cleared, by which the difficult journey towards truth is to be accomplished. The substitution, therefore, of a true for a false explication of phænomena, however in themselves unimportant, is never to be neglected; and with this view it was that I some time since addressed myself to the discovery of what I believe to be the true explanation of a somewhat singular and heretofore puzzling circumstance attendant upon the effects of earthquakes upon buildings, which has been frequently observed, and has been hitherto explained, so far as it has been attempted to be explained at all, by the assumption of a vorticose or gyratory movement having been in some inexplicable way given to the ground. The phænomenon alluded to, is the displacement of the separate stones of pedestals or pinnacles, or of portions of masonry of buildings by the motion of earthquakes, in such a manner that the part moved presents evidence of having been *twisted* in its bed *round a vertical axis*.

The first notice I find recorded of such a peculiar motion, is in the Philosophical Transactions, in an account of the

\* Communicated by the Author.

earthquake at Boston, in New England, of November 18th, 1755, communicated by John Hyde, Esq., F.R.S. He says, "the trembling continued about two minutes; near one hundred chimneys were levelled with the roofs of the houses, and many more shattered. Some chimneys, though not thrown down, are dislocated or broken several feet from the top, and partly turned round as on a swivel. Some are shoved to one side horizontally, jutting over, and just nodding to fall," &c. This author does not seem to have been struck with this odd circumstance of the twisting round of the chimneys, and offers no explanation. The next instance that I have found is in the account of the great earthquake of Calabria, in 1783, as recorded by the Royal Academy of Naples, quoted by Mr. Lyell, in his *Principles of Geology*, vol. i. page 482. After describing several other remarkable phænomena, tending to show the great velocity of the shock, such as that many large stones were found, as it were, shot out of their beds in the mortar of buildings, so as to leave a complete cast of themselves in the undisturbed mortar; while in other instances the mortar was ground to powder by the transit of the stone, he says, "Two obelisks (of which he has given figures) placed at the extremities of a magnificent façade in the convent of St. Bruno, in a small town called Stephano del Bosco, were observed to have undergone a movement of a singular kind. The shock, which agitated the building, is described as having been horizontal and vorticose. The pedestal of each obelisk remained in its original place, but the separate stones above were turned partially round, and removed sometimes nine inches from their position without falling." This is all that Lyell says upon the subject; he contents himself apparently with the vorticose account of the Neapolitan Academy.

I have found some few other notices of similar phænomena in old books of travels. Two additional instances, however, will be sufficient. The first will be found in the quarterly journal of the Royal Institution, in a narrative of the earthquake in Chili, of November 1822, communicated by F. Place, Esq.

The church of La Morceda, at Valparaiso, built of burnt bricks, stood with its length north and south. [The houses are built of adobes, or sun-dried bricks.] "The church tower, sixty feet high, was levelled; the two side-walls, full of rents, were left standing, supporting part of the shattered roof, but the two end-walls were entirely demolished. On each side of the church were four massive buttresses, six feet square, of good brickwork; those on the western side were thrown down and broken to pieces, as were two on the eastern side.



The other two were twisted off from the wall in a north-easterly direction, and left standing." The direction of the shocks was thought to be either from the south-west, or from the north-west.

We shall see hereafter evidence in the twisting of the two remaining buttresses, that the former was the real direction of the shocks, and that there was no vorticose motion, (indeed, the idea of two vortices, with centres only a few feet apart, is absurd upon the face,) but that the twisting of the buttresses is accounted for simply by a straight line movement, in connection with the attachment of the buttresses at one side, to the flank wall of the church.

The last instance I shall quote is from the pages of the able and delightful Darwin, in his *Journal of a Naturalist's Voyage* (Colonial Library, edit. p. 308), in describing the effects of the great earthquake of March 1835, upon the buildings in the town of Conception; and after noticing, also, the evidences of immense velocity in the shock, by which the projecting buttresses from the nave walls of the cathedral had been cut clean off close to the wall, by their own inertia, while the wall, which was in the line of shock, remained standing; he proceeds,—“Some square ornaments on the coping of these same walls were moved by the earthquake into a diagonal position. A similar circumstance was observed after an earthquake at Valparaiso, Calabria and other places, including some of the ancient Greek temples” (for which he quotes Arago, in *L'Institut*. 1839, p. 337, and Miers's *Chile*, vol. i. p. 392).

“This twisting displacement,” he proceeds, “at first appears to indicate a vorticose movement *beneath each point*, thus effected; but this is highly improbable. May it not,” he adds, “be caused by a tendency in each stone to arrange itself in some particular position with respect to the lines of vibration, in a manner somewhat similar to pins on a sheet of paper when shaken?”

The sagacity of Darwin at once showed him that the vorticose hypothesis was most improbable, and that in order to its being able at all to account for the phænomenon, a separate vortex must be admitted for every separate stone found twisted, the axis of rotation of the vortex having been coincident with that of the stone: besides this paramount improbability, therefore, a little further reflection would have led either Lyell or Darwin to estimate the necessarily inconceivable velocity of motion, at the extremity of the radius of one of these vortices, even if assumed at no more than a few hundred feet, in order that its velocity, within a few inches of the centre;

should be so great as to wrench out of its mortared bed, and twist a block of masonry by merely its own inertia.

Considering these circumstances, on lately reading the foregoing passages of Darwin, I was soon led to see that the twisting phenomena observed could be readily accounted for upon the established principles of mechanics, without having recourse to either vortices or vibrations, arranging blocks of many hundred weights, after the manner of pins on paper, or sand on one of Chladni's acoustic plates,—an explanation which, with all my admiration of Darwin, appears quite as far from probability as its predecessor.

I assume, then, nothing more than what is universally admitted, that during earthquakes a motion of some sort takes place, by which the ground itself, and all objects resting upon it, are shaken or moved back and forwards, by an alternate horizontal motion, within certain narrow limits, which, for all present evidence to the contrary, may be a straight line motion, though possibly variable in direction at different, and sometimes closely successive times, and the velocity of which is sufficient to throw down or disturb the position of bodies supported by the earth, through their own inertia.

Let us now apply this to the cases described of stones twisted on their bases, and the explanation will at once come to light.

If a stone, whether symmetrical or otherwise, rest upon a given base, and that motion be suddenly communicated horizontally to that base in any direction, the stone itself will be solicited to move in the same direction, and the measure of force with which the movement of the base is capable of affecting the stone or other incumbent body, is equal to the amount of friction of the latter upon its base—a function of its weight which, without the intervention of cement, may be from one-fifth to one-tenth of the weight of the body, for cut stone resting on cut stone, but may be increased to any amount by the intervention of cement.

The stone, however, is possessed of weight, and therefore of inertia; that is to say, being at rest, its whole mass cannot be instantly brought into motion by the plane, and if the amount of adhesion between the stone and its bed be less than the inertia due to any given velocity of horizontal movement of the bed, the bed will move more or less from under the stone, or the stone will appear to move in a contrary direction to that of the motion of its bed.

Now the inertia of the stone, which is here the resisting force, may be considered to act at the centre of gravity of the body.

The impelling force is the grasp of the stone, which its bed holds of it by friction or adhesion; and this may also be referred to some one point in the surfaces of contact, which we might call *the centre of adherence*.

If, then, a stone or other solid body rest upon a horizontal plane, which is suddenly moved with sufficient velocity to effect motion in the incumbent body, three several conditions of motion of the body may occur, according to the respective position of the centre of gravity of the stone, and of the centre of adherence.

1st. The centre of gravity of the stone may be at such a height above the base, that it shall upset by its own inertia. This is the case with houses, towers, walls, &c., when they fall by earthquakes, accompanied also by dislocation of their parts.

2nd. The centre of adherence may be in a point of the base, plumb under the centre of gravity of the stone; or in a vertical plane, passing through the centre of gravity of the stone, and in the direction of motion of the base.

In this case, the stone will appear to move in the opposite direction to that in which the base has moved; that is to say, the stone may have acquired more or less the direction of motion of the base, according as the motion of the latter has been longer or shorter continued, or less or more rapid; but, in so far as the movement in the opposite direction has taken place, the base, in reality, has slipped from under the stone.

3rd. The centre of adherence may neither be plumb under the centre of gravity of the stone, nor in the plane of motion passing through its centre of gravity, but in some point of the base outside the line of its intersection by this plane; in which case, the effect of the horizontal rectilinear motion of the base will be to twist the stone round upon its bed, or to move it laterally, and twist it at the same time, thus converting the rectilinear into a curvilinear motion, in space; the relative amount of the two compounded motions being dependent upon the velocity and time of movement of the base, and upon the perpendicular distance measured horizontally at the surface of adherence, between the centre of adherence and the centre of gravity of the stone.

This latter case is that which applies to the twisted stones of Calabria, South America and Greece; and affords, as I feel assured, the true explanation of the phænomena.

The relation of these forces, which have taken so many words to state correctly, might, of course, have been expressed algebraically in three lines; but as this would not be universally intelligible, I have preferred the more tedious and in-

elegant statement of words; and to render the matter quite familiar, have prepared a model of one of the Calabrian pedestals, figured by Mr. Lyell, which will exhibit to the eye all the phænomena already adverted to, by giving by the hand a rectilinear horizontal motion to the base\*.

I have now proved that no vorticose motion is requisite to account for the twisting of obelisks, &c., as observed in earthquakes, and that nothing more than a simple horizontal rectilinear motion is demanded; but, it may be asked, if this rectilinear horizontal motion in earthquakes be an alternate one also—if the earth shake both back and forwards—how is it that these and other displaced bodies are not moved back into their places again by the reverse motion, by the same sort of motion, acting in the contrary direction?

This question is, I believe, fertile in consequences, and its consideration has led me to some further conclusions as to the nature of earthquake motions. The first reason obviously is, that as the forward movement has by displacement produced a new set of conditions as to the centres of gravity and of adherence of the stone and base, so it can scarcely by possibility ever occur that there shall be precisely such as to give rise to such a new form of twisting motion as shall neutralize that first produced, although it is quite probable that *some* second twisting may be produced by the backward stroke or motion; for this view I am indebted to my friend Dr. Apjohn. But this alone is not sufficient. After looking through a great number of authors, on earthquakes, I have not been able to find one that has endeavoured, far less succeeded, in shaping to himself any distinct notion as to what the precise nature of the earthquake movement is. The ancients, appealing to their senses, so far as these could guide them, thought that it was like the shaking of a sieve, as the word *σεισμός* tells us. The moderns in general are not more exact in their notions: a trembling, a vibration, a concussion, a movement, and so forth, are the words we find scattered through even scientific authors. Mitchell, Lyell and Darwin, with some others, although they obviously have formed no distinct idea on the subject, use the word “undulation,” and in so far, have come nearer to the truth; for it appears to me, that the fact, that displaced bodies are not occasionally replaced, in earthquakes, is conclusive evidence of either one of two things: either the motion is limited to horizontal direct movement, in one or more directions; and, if so, the whole mass of the disturbed country must be pushed bodily forward, and remain so, of which there

\* Exhibited at the Geological Society of Dublin, from whose Transactions this paper is extracted.—Read 8th December 1845.

is no evidence; and all bodies must, as the effect of one shock, fall in the one direction, and not in opposite directions, which is contrary to observed facts: or, on the other hand, if the movement be an alternate horizontal motion, as all observations go to prove it is, then the motion in one direction must be slower than in the other, or attended with other differences of circumstances. The backward motion must be different from the forward motion, or otherwise displaced bodies would be replaced by the recurrence, in the opposite sign of forces similar and equal to those that first set them in motion; but they are not found so replaced.

Now, of all conceivable alternate motions, the only one that will fulfill the requisite conditions observed, namely, that shall move with such an immense velocity as to displace bodies by their inertia, or even shear close off great buttresses from the wall, they sustained, (Darwin) or project stones out of their beds, by inertia; that shall have a horizontal alternate motion, either much quicker in one direction than in the other, or different in its effects; and that shall be accompanied by an upward and downward motion at the same time—a circumstance universally described as attendant on earthquakes—the only motion, I say, that will fulfill these conditions, is the transit of a great solitary wave of elastic compression, or of a succession of these, in parallel or in intersecting lines through the solid substance and surface of the disturbed country.

The general idea of the nature of earthquake motion, viz. that it consists of a wave of some sort, is not however new, although so entirely neglected by the mass of recent geological authors. To the Rev. John Mitchell, M.A., Fellow of Queen's College, Cambridge, the merit of this idea appears originally due. In a paper communicated to the Royal Society, read in 1760, and published in the 51st vol. of the Philosophical Transactions, part 2nd, he treats at length of the origin and phænomena of earthquakes, and distinctly enunciates the following view:—

That the motion of the earth is due to a wave, propagated along its surface, from a point where it has been produced by an original impulse. This impulse, he conceives, to arise from the sudden production or condensation of aqueous vapour, under the bed of the ocean, by the agency of volcanic heat, the supposed mechanism of which he minutely describes; but while he was so far right in his conception of an elastic wave of *some sort*, I expect to be able shortly to prove that he has wholly mistaken the nature of the wave that actually occurs, and that a wave, such as he assumes, can have no existence consistently with the physical structure of our globe, with the

observed facts of earthquake motions, or even with the conditions of his own hypothesis.

LXXXIII. *On the Causes to which Musical Sounds produced in Metals by discontinuous Electric Currents are attributable.*

By Prof. ÉLIE WARTMANN\*.

SINCE the discovery made in 1837 by Dr. Page, and verified the following year by Prof. Delezenne, of the possibility of producing a musical sound by electricity, this interesting phenomenon had scarcely been studied, when in 1844 MM. Marian, Beatson, Gassiot, and De la Rive all at once made known the various conditions of its production. The interesting memoir of the last gentleman, printed in vol. v. p. 500 of the *Archives de l'Electricité*, contains a great number of very valuable results. But the theoretical part of the subject has not yet been presented under a precise and general form, and it is with a view to supply, if possible, this gap, that I have undertaken the following experiments. I imagined them in the month of August 1845, in consequence of a meeting at which M. de la Rive exhibited his curious apparatus to Prof. Dove and myself.

A well-annealed soft-iron wire, 1<sup>m</sup>·7 long and 2<sup>mm</sup>·5 in diameter, was fixed in a horizontal position on a thick trencher of hard wood inserted into the wall. One of its extremities was held back by the jaws of a clamp, whilst the other supported a weight of 24 kilogrammes. Upon a cork, pierced by friction with the wire, I arranged a small plane mirror, with parallel faces, made at the Optical Institute of Munich and intended to reflect, into a telescope furnished with cross wires, the divisions of a scale placed at a distance of two metres. This arrangement, similar to that of the magnetometer, exhibits the least deviations of the reflecting surface, when it is not displaced parallel to itself. The iron wire passed through a wooden reel, the bore of which was five centimetres in diameter, and on which were rolled three copper wires enveloped with silk, 23<sup>m</sup>·6 long and 3 millimetres in diameter. I employed a Bunsen's battery of eleven pairs, and a mercurial rheotome or contact-breaker; these two instruments were enclosed in an ante-room adjoining the laboratory.

According to the place which the wire occupies, it becomes the seat of greater or less transversal vibrations, whose plane may be varied at will. In general, in any position of the wire, the intensity of the effect varies at different points of its length, as is perceived on bringing the mirror to such points. The

\* Communicated by the Author.

amplitude of the vibrations is not the same for different parts of the wire subjected similarly to the reel. M. de la Rive found this by the comparison of the sounds obtained. These phænomena result from the attraction exerted upon the wire by the parts of the coil which are the nearest to it: they cause a distinct class of sounds. But there exists another cause of vibrations in the wire, the effect of which is more or less independent of this lateral attraction. Longitudinal vibrations are produced in it, with which correspond sounds of a peculiar character. If the axis of the reel was identical with that of the wire, supposing it exactly rectilinear and cylindrical, a transverse deviation would no longer take place. But even then, the molecules on which the electro-magnetic action is exerted are attracted right and left of the centre of the axis of the reel towards this central point, as a steel needle is seen to be drawn into it as soon as it is introduced into the hollow of the helix. It is this internal vibration which, by the discontinuity of the electric current, is rendered periodical in two opposite directions, that determines the second class of sound.

Let us now pass to the case of the current transmitted by the wire. In order to study it, I substituted for the mirror the spherical and perfectly polished bulb of a small mercurial thermometer. The optical axis of the telescope, passing through the intersection of the crossed wires, was directed on the brilliant image of a luminous point reflected very obliquely at the upper part of the convexity. This arrangement discovers any change in form of the wire, even in the direction of its length. I was not able to perceive any elongation of the wire under the electric action, although it gave a very distinct sound. I attribute the principal cause of this sound to the polary arrangement which the molecules undergo in order to give passage to the electricity. This arrangement is manifest in many cases, and I have elsewhere pointed out a very great number of them\*. It is the result of a struggle between the molecular forces which constituted the primitive state of equilibrium of the body and the new activity which the dynamical condition of the fluid excites. If the flow of the latter is continuous, this struggle is instantaneous, and the noise which it occasions is null or nearly so; but it recommences with each closing of the circuit if the flow is periodical.

It is already known from the experiments of M. Peltier †,

\* Memoir on the Electric Diathermance of Voltaic Pairs: *Archives de l'Electricité*, vol. i. page 74.

† *Comptes Rendus des Séances de l'Acad. des Sciences de Paris*, Jan. 6th, 1845.

and of various scientific men, that the prolonged passage of the electricity by metallic wires alters essentially their tenacity. It seemed to me very probable that the elasticity of wires subjected for some time to the intermittence of currents which renders them sonorous, must be altered in a permanent manner.

Since the experiments just mentioned were made, M. Wertheim has published\* a very interesting note, in which he describes a process of observation analogous to mine, although less delicate, and indicates the causes to which he attributes the sounds produced. Although I agree with him on most points, I differ from him both as to what relates to the attraction exerted from the two sides of the centre of the helix, an attraction which he does not mention, and in the explanation of the case in which the wire is directly traversed by the discontinuous current. The skilful experimentalist whom I have just named attributes the sound produced to the heat engendered by the current. Nevertheless my wire indicated no perceptible heat. It results from the experiments of M. de la Rive and my own, that the sonorous state continues with more than 600 interruptions a second. How shall we admit that the elevation of temperature and the diminution of elasticity which accompany it can disappear in  $\frac{1}{600}$ th of a second? The current of a pile of eleven pairs certainly does not alter the thermal state of a bar of a centimetre square in section, as I have directly established †: nevertheless, if it is discontinuous, it renders it sonorous. I may add, finally, that this heating does not take place when the reel is employed, as any one may convince himself by placing a bismuth and iron pair in its hollow, connected with a very delicate rheometer. Nevertheless the sonorous property may be the same as with the wire directly subjected to the current.

Lausanne, March 16, 1846.

LXXXIV. *An Account of various Substances found in the Guano Deposits and in their Vicinity.* By E. F. TESCHEMACHER, Esq. †

REPORTS having been circulated that large quantities of saltpetre (nitrate of potash and nitrate of soda) were to be found of a very good quality in the neighbourhood of the guano deposits on the coast of Africa, numerous vessels were

\* *Comptes Rendus*, Feb. 23, 1846.

† *Phil. Mag.*, Oct. 1843; *Archives de l'Electricité*, vol. ii. page 601.

‡ Communicated by the Chemical Society; having been read December 1, 1845.



despatched both from London and Liverpool in search of those valuable substances, particularly as it was considered they might be obtained upon the same terms as Ichaboe guano, namely, for nothing but the labour and expense of fetching. No favourable accounts however have as yet been received as to the success of these undertakings. The evidence of such deposits existing there at all was very unsatisfactory; the circumstance much relied upon was the existence of large beds of nitrate of soda in the neighbourhood of the coast of South America, and large deposits of guano similar in many respects to the deposits of guano on the African coast: there was certainly an abundance of animal matter and ammoniacal salts to furnish the nitric acid, and a temperature high enough to effect the decomposition, but the source from whence the alkaline bases of potash and soda were to be derived was not very evident. The principal source of saltpetre in the East Indies is from numerous districts of nitrous earth found on the surface of the soil, which being compounds of lime and magnesià with nitric acid, they are dissolved out, and the saltpetre subsequently formed by the decomposition of these nitrous compounds by potash salts. The nitrate of soda saltpetre beds in the Province of Tarapaca near Iquiqua on the coast of South America, are the only instances known of the occurrence of saltpetre ready-formed in extensive beds, but even this deposit contains the salt in a state of great impurity.


These explorations, however, on the African coast have brought to light various other substances which have been found there, the details of which are more particularly the object of this communication.

The substances which I shall now describe are found in the guano beds, or in their vicinity, either in a crystalline state, or in distinct masses. The first substance is a crystalline salt, perfectly transparent, with a cleavage and brilliant faces in one direction only; it gives a yellow precipitate with nitrate of silver; gives off ammonia upon application of caustic potash, and when heated to redness loses about 50 per cent. of water and ammonia; I consider it therefore to be *phosphate of ammonia*. The portion of salt I examined consisted only of a few grains, and was consequently too small a quantity to analyse with exactness.

The next substance was also a crystalline salt a little mixed with guano in its cavities; it possessed a cleavage with brilliant planes in two directions: upon examination with the reflecting goniometer, it gave  $112^{\circ}$  as the measurement of the angle formed by the meeting of the adjacent planes. Upon analysis I found it to consist of—

21·0	parts of Ammonia.
55·50	... Carbonic acid.
23·50	... Water.
<hr/>	
100·00	

being nearly equivalent to 1 atom of ammonia, 2 atoms of carbonic acid, and 2 atoms of water.




and is consequently a *bicarbonate of ammonia*.

The third substance was found at Saldanha Bay on the coast of Africa, imbedded in patches in the mass of guano. It is found in distinct crystals with numerous modifications, many of the planes possessing sufficient brilliancy to enable me to measure the angles by the reflecting goniometer. I have given the measurements of one crystal, from which it appears the primary form is the right rhombic prism of  $57^\circ 30'$  and  $122^\circ 30'$ : it has a cleavage parallel to plane M\*. Upon analysis I find this substance to be composed of—

14·30	parts of Ammonia.
17·00	... Magnesia.
30·40	... Phosphoric acid.
38·10	... Water.
<hr/>	
99·80	

which is nearly equivalent to 1 atom ammonia, 1 atom magnesia, 1 atom phosphoric acid, 5 atoms water.



It is therefore the *ammonio-magnesian phosphate*. The specific gravity is 1·65, hardness 2; it falls to powder before the blowpipe, giving off water and ammonia. It occurs white, translucent, sometimes coloured brown by the guano; it readily dissolves in weak acids.

This substance is clearly derived from the guano; but being insoluble in water, it must have been held in solution by some of the organic acids of the guano, and deposited therefrom in large crystals, as they are found, but disseminated in patches only of the guano, in various parts of the beds.

This substance not having been found before in a native state, but hitherto only known as one of the artificial products of the laboratory, must be considered as a new mineral body; I therefore propose to give it the mineralogical name of *Guanite*, this name being derived from the circumstances and locality of its formation.

The source from which the first two substances, namely, the phosphate of ammonia and the bicarbonate of ammonia,

\* See the angular measurements subjoined.

are derived, is clearly the percolation of water through the guano beds dissolving out these salts, which running into lower situations may be detained in lagoons and hollows of rocks, where being subject to the high temperature of the climate they would be evaporated down, leaving these salts in the crystalline state described. As guano contains abundance of these two salts, it is possible there may exist considerable masses of them; should this be the case, it is evident that to the chemist in particular it would be of great interest as an additional source of these valuable salts.

The chance of finding any considerable quantity of guanite in the state of crystals is not great, but as it forms one of the ingredients of guano it is a substance of some importance. The application of it as a manure in combination with other ingredients is likely to be highly beneficial, it being a compound containing two important substances in an insoluble state, namely, ammonia and phosphoric acid; these may be taken up by plants only as they may be required, and not be liable to be dissolved out of the soil or evaporated like other ammoniacal salts.


The last substance which I shall describe was also found imbedded in the guano from Saldanha Bay; it consists of small globular particles composed of concentric laminae slightly adhering together, of a yellowish white colour, containing in places portions of a similar nature, which on fracture have appearances of an organic structure like bone, but on examination by the microscope proved to be portions of shells resembling Nummulites. On analysis I found the substance to be composed of—

37.50	parts	Carbonate of lime.
32.50	...	Carbonate of magnesia.
12.00	...	Phosphate of lime.
12.00	...	Water with a little ammonia and animal matter.
3.00	...	Sand.
2.50	...	Alkaline sulphates and chlorides.
99.50		

There does not appear to be any great quantity of this substance. How it has been formed it is difficult to imagine; the composition is so very different either from that of bones or shells, particularly in regard to the large quantity of carbonate of magnesia which it contains. It is however probable that both bones and shells form the base of this substance, and that partial decomposition having taken place, the magnesia may have subsequently entered into combination with the carbonate and phosphate of lime.

## Measurements of Guanite.

M on M'	57°·30
M on f	118°·30
M' on f	118°·30
M' on h	151°·00
f on h	89°·30
M on e	142°·10



M' on e'	142°·10
h on c	133°·20
e on e'	91°·50
e on f	112°·20
e' on f'	112°·20
e on c	142°·10

## LXXXV. Notes on the Preparation of Alloxan.

By WILLIAM GREGORY, M.D., F.R.S.E.\*

IN an interesting and able paper on alloxan and its derivatives, the first part of which appears in Liebig's *Annalen* for September 1845, Schlieper enters into minute details concerning the most advantageous method of preparing alloxan, and after describing the results which he obtained on repeating the process given by me, proposes a new method of his own, which he considers in every way preferable, as yielding, with greater facility and certainty, a larger proportion of alloxan. Professor Liebig in his Lectures (*Lancet* 1845) also recommends Schlieper's method as the best in every respect.

I am still, notwithstanding, inclined to give a decided preference to my own process, when carefully performed, and that on the grounds of its superior simplicity, facility and productiveness. A brief comparison of the two methods, with their results, will enable the reader to judge for himself.

I must first of all, however, observe, that Schlieper, in repeating my process, has not obtained results so favourable as I had formerly announced; so that, in his hands, his own method has been the more productive. I formerly obtained from 100 parts of uric acid 90 of crystallized (hydrated) alloxan, perfectly pure, not reckoning the portion of alloxan remaining in the mother-liquids. Schlieper, on the other hand, from 15 ounces of uric acid, treated by my process, obtained, including the contents of the mother-liquids, 8 ounces hydrated alloxan,  $1\frac{3}{4}$  ounce alloxantine (=  $2\frac{1}{4}$  ounces alloxan), and  $\frac{3}{4}$  ounce parabanic acid; in all equivalent to about  $11\frac{1}{4}$  ounces of alloxan. This only amounts to 75 per cent.; whereas I obtained 90 per cent., exclusive of the mother-liquids, which I find on an average to yield fully one-tenth more; in all, therefore, at least 100 per cent. I may here state that I have

\* Communicated by the Chemical Society; having been read December 15, 1845.

never failed to obtain this as an average result since my process was published, although I have very often repeated the process. Several of my pupils have been equally successful. I shall now, therefore, describe the process as I have for some time pursued it, and its simplicity will, I trust, be evident.

In my original account of this process, I recommended the use of nitric acid of sp. gr. 1.3 to 1.35, and it was with such acid, as I believed, that my results were obtained. But as Schlieper found it impossible to succeed with acid of less sp. gr. than 1.4 to 1.42, I suspect that I may have been mistaken as to the sp. gr. of my acid. This I cannot now ascertain; but it is rendered probable by the circumstance that, in the experiments about to be mentioned, I found an acid of 1.412 to answer my purpose perfectly, with the same appearances as I had formerly observed.

Schlieper having corrected this error proceeds to describe my process, as performed by him, with great accuracy and minuteness, and his description of the phenomena entirely agrees with my experience. I can only account for his not obtaining such favourable results as I have always done, to the circumstance of his acid being a little too concentrated. However this may be, on reading his paper I proceeded to repeat my process, and obtained the results to be hereafter stated.

The following is the process I now follow:—2 or  $2\frac{1}{2}$  fluid ounces of colourless nitric acid, sp. gr. 1.412, are placed in a flat-bottomed dish or beaker glass, and as much uric acid is introduced as will lie on the point of a small spatula. This is well-stirred in to prevent the formation of lumps, and in a few minutes effervescence commences, the liquid becomes slightly warm, and the powder dissolves. More uric acid is now added, taking care never to exceed a certain small quantity, and not to allow the liquid to become warm beyond a certain degree, which is easily judged of by laying the dish on the hand. If too hot when uric acid is added, or if too much acid be added at once, the uniform steady effervescence is changed into a violent and tumultuous action, after which no alloxan can be obtained. It is proper to have a plate with cold water at hand, in which to place the dish or glass if it should seem likely to become too warm. But a little practice enables us to regulate the operation so that no external cooling is required.

After several portions of uric acid have been added, crystals of alloxan begin to appear in the warm liquid, but the addition of uric acid is to be continued, with the same precautions, till so much alloxan has been formed, that on cooling the whole becomes nearly semisolid. When this point is reached

the liquid has become somewhat viscid, and this, along with the presence of the crystalline deposit of alloxan, gives a peculiar character to the effervescence toward the end of the operation. I commonly find that with  $2\frac{1}{2}$  fluid ounces of nitric acid the point above alluded to is reached when about 1200 grains of uric acid dried at  $212^{\circ}$  have been dissolved. It does not answer to operate on a much larger scale; it is better to use several dishes at once, each containing  $2\frac{1}{2}$  or at the most 3 fluid ounces of acid. For every 500 grains of uric acid 1 fluid ounce of nitric acid may be allowed.

The whole is now allowed to stand all night in a cool place, and next day the alloxan is collected on a funnel with the aid of a little asbestos. The mother-liquid drains off, and the last portions of it are cautiously displaced by ice-cold water, till the droppings are found to have only a moderately strong acid taste. The alloxan on the funnel, which is anhydrous, is then digested with just as much water at  $140^{\circ}$  or  $150^{\circ}$  F. as will dissolve it. The solution is filtered, and on cooling deposits a large crop of crystals of hydrated alloxan. [Should too much water have been added, the filtered liquid must be evaporated at from  $120^{\circ}$  to  $140^{\circ}$  F., till on cooling it crystallizes abundantly.] The mother-liquid of these crystals, evaporated at the same temperature, yields a second crop. The mother-liquor of this is added to the acid mother-liquor previously drained off, and the whole liquid treated, after the addition of two or three times its bulk of water, with sulphuretted hydrogen, till the alloxan present is reduced to the state of alloxantine. As a part is always reduced still further to dialuric acid, the liquid must be exposed to the air for a day or two, or until it deposits no more crystals. The alloxantine is purified by solution in boiling water, filtration to separate sulphur, and crystallization; and when dry three parts of it correspond to rather more than four of hydrated alloxan. If required, it may very easily be converted into alloxan; as Schlieper has described this process I need not repeat it here.

The mother-liquid of the alloxantine generally yields some parabanic acid; but very little if the process has been carefully performed.

I think it will be admitted that the above process is sufficiently simple. It will be observed that I no longer recommend the separation of the alloxan formed from the nitric acid in several successive portions, but that there is only one operation for all, in which the alloxan is collected on a funnel with asbestos. I used sometimes to divide the process into five successive operations, and generally made three of them: but I am now convinced that it is best to dissolve in the nitric

acid the whole of the uric acid that is to be dissolved before collecting the alloxan:

Let us now consider the productiveness of this method. I have already stated my average of former results to have been 90 per cent. of crystallized alloxan, exclusive of the mother-liquid, which corresponded to one-tenth more. As the process now stands we have—1. The first crop of crystals of alloxan, varying with the proportion of water used to dissolve the anhydrous alloxan. 2. The second crop of the same crystals. 3. The alloxantine from the mother-liquid converted into alloxan, or calculated in that form. I take no account of the parabanic acid.

*Experiment 1.*—Uric acid 2600 grains; alloxan, first crop, 1950 grains, second crop, 550 grains; alloxantine, 200 grains, equivalent to alloxan, 290 grains. In all, therefore, from 2600 grains of uric acid, 2790 grains of hydrated alloxan, or 107 per cent. nearly.

*Experiment 2.*—Uric acid, 1130 grains; alloxan, first crop, 800 grains, second crop, 140 grains; alloxantine, 80 grains, equivalent to alloxan, 116 grains. In all, therefore, 1056 grs. of alloxan from 1130 of uric acid, or 93 per cent.

*Experiment 3.*—Uric acid, 1500 grains; alloxan, first crop, 1150 grains, second crop, 270 grains; alloxantine, 120 grains, equivalent to alloxan, 174 grains. In all, therefore, from 1500 grains of uric acid, 1594 grains of alloxan, or 106 per cent.

The above results, averaging 102 per cent. of pure hydrated alloxan, were obtained without difficulty. Indeed the only delicate point in the process is the attention necessary to avoid too great a rise in temperature, alloxan being decomposed by heat even when simply dissolved in water, but still more when acid is present. A little experience however makes this quite easy; and besides, this difficulty attaches equally to Schlieper's new method, as we shall see.

The formula of uric acid being  $C_{10} N_4 H_4 O_6$ , while that of hydrated alloxan is  $C_8 N_2 H_4 O_{10} + 6 aq$ , it is obvious that 100 parts of uric acid can produce about 128 of alloxan. It is not likely that we shall ever obtain the full proportion without loss, but I consider my process, simple as it is, to furnish a very satisfactory approximation, considering the impossibility of separating the whole alloxan from the acid liquid in which it is formed.

If we now refer to Schlieper's account of his new method, we find that it includes the following operations:—1. The uric acid is acted on by hydrochloric acid and chlorate of potash, care being necessary, as in my process, to keep the

temperature below a certain point. 2. The *whole* of the alloxan is reduced by sulphuretted hydrogen to the state of alloxantine. 3. The alloxantine is reoxidized by nitric acid, and thus reconverted into alloxan. I cannot admit that this process is either more simple or more easy than my own. On the contrary, as I obtain nine-tenths of the whole alloxan, or 90 parts from 100 of uric acid directly as alloxan, and pure, in the first crystallizations, while Schlieper first converts all his alloxan into alloxantine, and then reconverts the alloxantine into alloxan; and further, as I use no other reagent but nitric acid in preparing these nine-tenths, the advantage of simplicity and facility is entirely on my side.

From 4 ounces of uric acid, Schlieper obtains by his own process 2 ounces 7 drachms and 20 grains of alloxantine, equivalent theoretically to 3 ounces and 7 drachms of alloxan, or nearly 97 per cent. But in reconverting this alloxantine into alloxan by nitric acid, it will be found impossible to obtain, practically, the whole alloxan, since some of it must remain in the mother-liquid; and moreover, in the process of oxidation by heating with nitric acid some alloxan is very likely to be converted into parabanic acid, and thus lost. Judging from experience, I should not expect the 97 per cent. of alloxan obtained in theory to yield, in crystals, more than 90 per cent.

As far as productiveness, therefore, is concerned, I may claim also a superiority for my method. It is true that it has not succeeded so well in the hands of Schlieper, but this must I think be attributed to accidental causes, and possibly to a want of perfect familiarity with the method on the part of Schlieper, who seems to be so good an operator, that I cannot doubt that he would, after a little practice, obtain the same results as I have always succeeded in obtaining.

Finally, I beg to remind those who may wish to try my process, that what Schlieper describes as a modification of my process is the process itself, unmodified; because the only change introduced by Schlieper consists in the use of acid at 1.4 or 1.42 instead of 1.3 or 1.35, as erroneously recommended in my original process. In point of fact, the acid which I have long used for the purpose has the sp. gr. 1.412, and for this number 1.3 or 1.35 was accidentally substituted in writing or printing my former notice. In common with all chemists I am much indebted to M. Schlieper for pointing out this oversight.



LXXXVI. *Intelligence and Miscellaneous Articles.*

## ON CHLOROAZOTIC ACID.

**M.** BAUDRIMONT remarks, that Mr. Edmund Davy published his researches on aqua regia in 1831, and concluded from them, that what he terms chloronitrous acid is composed of equal volumes of chlorine and nitric oxide gases, which combine without alteration of volume.

According to M. Baudrimont, the presence of [uncombined] chlorine in the product obtained by Mr. Davy prevented the product from being properly examined, and he therefore undertook fresh researches on the subject.

In order to prepare the active product [chloroazotic acid] of aqua regia, M. Baudrimont mixed three parts, by weight, of nitric acid, of specific gravity about 1.314, with five parts of hydrochloric acid, of specific gravity 1.156; this mixture yields a colourless liquid, which after an uncertain time becomes red, according to the temperature of the air and the intensity of the light to which it is exposed. If, however, the mixture be heated, it becomes red at about 187° Fahr., and yields vapour of the same colour; the temperature gradually increases to nearly 230° Fahr., and then remains so invariably during the whole time of the operation.

If the product of the distillation be received in a vessel properly cooled, a red liquid is obtained; but if the neck of the retort be simply passed into a receiver, a red vapour is formed which does not condense, and a colourless liquid is condensed.

This experiment shows that this distillation yields two distinct products—a red vapour which is very volatile, and a colourless product which is more fixed. It shows also that the temperature of 230° Fahr. does not indicate a boiling-point, but a fixed point of decomposition. By adopting the requisite arrangements, the red vapour may be condensed in tubes in the state of a red liquid, which boils at a very low temperature; it can be preserved only in tubes hermetically sealed. This is what the author terms chloroazotic acid; the properties of which are as follows:—

At a sufficiently low temperature it is a red limpid liquid, surmounted with vapour of the same colour; its boiling-point is about 20° Fahr.; from this it follows that it is gaseous at ordinary temperatures. In the state of gas it is red, and possesses a suffocating odour, analogous to that of chlorine, but still differs considerably from it.

The extreme volatility of chloroazotic acid presented almost insurmountable difficulties to the determination of its principal properties.

The elements of chloroazotic acid reduced to volumes, and the volumes multiplied by the corresponding specific gravities, give the following results:—

N or 2 volumes	.....	$2 \times 0.9727 = 1.9440$
O <sup>3</sup> or 3	... ..	$3 \times 1.1057 = 3.3171$
Cl <sup>2</sup> or 4	... ..	$4 \times 2.4216 = 9.6864$

9 elementary volumes give . . . . 14.9475

One experiment on the specific gravity of chloroazotic acid gave 2.49, and another 2.45, and lead to the same result,—

$$\frac{14.9475}{6} = 2.49.$$

Thus the 9 volumes of elementary gases which form chloroazotic acid are condensed into 6 volumes, and one volume of the acid contains  $\frac{1}{3}$ rd volume of nitrogen,  $\frac{1}{2}$  of oxygen, and  $\frac{2}{3}$ rds of chlorine. The specific gravity of the liquid acid was found to be 1.3677.

Chloroazotic acid consists of—

Nitrogen ..	12.6 or 1 eq.
Oxygen....	22.4 ... 3 ...
Chlorine....	65.0 ... 2 ...

100.

The extreme volatility of chloroazotic acid renders the examination of its chemical reaction extremely difficult, and it can be effected only at very low temperatures.

With phosphorus, the acid enters into ebullition, and disappears without acting sensibly upon it; arsenic in powder is acted upon, and yields a white product; silver in powder occasions deflagration, and the liquid disappears; gold is rapidly dissolved, but platina is acted upon with more difficulty; alcohol yields an æthereal odour, analogous to that of nitric æther.

Chloroazotic acid in the gaseous state appears to have no action on phosphorus at ordinary temperatures; the latter may be even liquefied by heat, without producing any more apparent action.

Pulverized arsenic and antimony burn vividly in the gas; bismuth is immediately attacked, yielding white vapours, but unaccompanied with light; potassium is slowly acted upon at common temperatures, but the reaction is violent when heated to its fusing-point; there occurs sudden increase of temperature, accompanied with vivid light; gold is acted upon, and a plate of copper, heated to dull redness, burns very vividly; tin heated nearly to its melting-point, does not appear to be immediately attacked, but in a little time it is tarnished and rendered white; mercury is immediately acted upon; one-half of the gas disappears, and the remainder is nitric oxide, entirely absorbable by solution of protosulphate of iron.—*Ann. de Ch. et de Phys.*, Mai 1846.

NOTICES OF NEW LOCALITIES OF RARE MINERALS, AND REASONS FOR UNITING SEVERAL SUPPOSED DISTINCT SPECIES. BY FRANCIS ALGER, MEMBER OF THE BOSTON SOCIETY OF NATURAL HISTORY\*.

*Phacolite from New York.*

This rare mineral, which comes to us principally from Bohemia and Ireland, I have discovered among a suite of specimens of various kinds found on New York Island, near Harlem, by Messrs. Mathews and Johnson, of New York city. The specimens, which eventually proved to be this mineral, were labelled stilbite; but their appearance was so peculiar, that I questioned at the time whether they had been correctly designated, and determined to examine them carefully at my earliest convenience. I have since received two other specimens, better characterized than the first, from Mr. Johnson. The crystals are in a geode form, implanted on calcareous spar, and associated with silver-coloured mica and a few scales of oligisto-magnetic iron ore. They are of a wax or honey-yellow colour, have a waxy lustre, and the smallest individuals are translucent. They are brittle, breaking with an uneven fracture, have none of the foliated structure of stilbite, and afford no indications of cleavage. Hardness superior to that of stilbite, and equal to that of chabasite. Their surfaces are roughened or pitted, so as to reflect no image by which they could be subjected to measurement by the goniometer. Before the blowpipe, a fragment of the mineral swells and intumesces slightly, like the Bohemian and Ferroe chabasite, and fuses into an opaline, blebby bead; at the moment of ignition, in the outer flame, it gives out a beautiful green phosphorescence, which I have also noticed, in a less degree, in the phacolite from Ireland. It is soluble in hydrochloric acid. The crystals, at first sight, appear to be rounded, and to have no determinate form; but, on closer examination, some of the smaller and more isolated ones are found to be nearly perfect double six-sided pyramids, precisely similar to the phacolite from Bohemia, differing from it only in colour and lustre. I cannot doubt that, like that mineral, they are secondaries to a primary rhombohedron, probably of the same measurements, and are also identical with it in composition. The absence of well-defined cleavage is unfortunate, but this is a defect which applies equally to the foreign mineral. Nor is the rhombohedral cleavage of ordinary chabasite, of which phacolite is by many supposed to be only a variety, by any means easily determined; in fact, Sir David Brewster has suggested, from optical investigations, whether the primary form of chabasite be not a prism.

*Is Phacolite a variety of Chabasite, or distinct from it?*

Tamnan of Berlin, in his very complete little essay on Chabasites, has given very good reasons for uniting the two; while Breithaupt has maintained them to be distinct. The primary rhombohedron of

\* From the Journal of the Boston Society of Natural History.

phacolite, according to Breithaupt, is P on P,  $94^\circ$ , that of chabasite P on P,  $94^\circ 24'$ . Phillips makes the last  $94^\circ 46'$ . The analyses of Anderson and Rammelsberg would seem at first to show a marked difference in their composition, a difference which is also shown by the different analyses of common chabasite, resulting in varieties having different formularic expressions. For example, acadiolite contains three per cent. more of silicic acid than common chabasite, and is a tersilicate of lime and the other isomorphous bases, instead of a bisilicate of the same bases. The mineralogical formula of acadiolite is  $3Al Si^2 + (Ca, N, K,) Si^3 + 6 Aq$ , while that of chabasite is  $3Al Si^2 + (Ca, N, K,) Si^3 + 6 Aq$ . Rammelsberg is inclined to regard phacolite as a mixture of acadiolite and scolecite (lime mesotype), the latter containing an additional atom of water\*. By uniting the atoms of both, he thus states the chemical formula for phacolite:  $2RO SiO^3 + 2Al^2O^3 3Si^3 O^3 + 10HO$ . As the analyses stand (compare Berzelius's and Thomson's with the two just referred to), phacolite differs from chabasite in containing three per cent. less of silicic acid, and three atoms less of water. Now it is obvious that these differences are insufficient to authorize a separation of the two minerals, unless there be a want of agreement in crystallographical and other characters, greater than that as yet pointed out. An equally valid reason could be urged for the separation of acadiolite from chabasite, on the ground of a difference in their composition, had not the examinations of Prof. G. Rose proved an exact agreement in the angles of their primary crystals. So, also, of levyne and gmelinite, which are now admitted to be only varieties of chabasite, their occurring forms all being secondaries to the same primary rhombohedron. The evidence of the identity of any two minerals is best shown by the incipient or intermediate passages of one into the other, in the same specimen. I am not aware that, in the case of the Irish or Bohemian phacolite, such evidence has been adduced; no tendency of the sort is shown in the specimens I have examined from those countries. Now one of my specimens from New York has the distinct form of chabasite (the perfect rhombohedron) and of phacolite (perfect double six-sided pyramids). The first form, however, is rare; the incipient replacements are also shown; but these crystals have not the full perfection of waxy lustre reflected by the ultimate form of phacolite,—a singular effect, attributable, probably, to the nature of the solvent in which the molecules were suspended.

#### *Approach of twin-crystals to the Phacolite form.*

These, as they are sometimes presented, would, unless carefully examined, be mistaken for the true form of phacolite. The most perfect specimens I have seen are from Nova Scotia. They consist of two rhombohedrons united in the usual manner, each crystal turned half round, but having their superior edges and lateral angles

\* See First Suppl. to his *Handwörterbuch*, p. 112. It was on these grounds that Hoffmann proposed to separate acadiolite, as well as the Gustafsberg variety, from chabasite.—Poggendorff's *Annalen*, xxv. 495.

deeply replaced. The approach to the form of phacolite is thus produced; the edges and angles not standing out in relief, as they ordinarily do in these twin forms. The striæ, parallel with the edges of the two rhombohedrons, so intersect as to show the compound nature of the crystals. Dr. C. T. Jackson has a fine specimen of this variety from the Two Islands, in Nova Scotia, of a wine-yellow colour; I have another pure white, from the same place.

#### *Ytthro-cerite.*

This rare mineral is found, associated with brucite, in rolled masses of limestone, in the town of Amity, Orange county, New York. I have as yet seen but two specimens of it, which I found among some fragments of limestone containing brucite and mica, in the duplicate collections belonging to the late Dr. Horton of Edenville. It attracted my attention as being unlike fluor spar, which it was supposed to be at the time, and I have now satisfied myself that it is ytthro-cerite, though I have not gone so far as to detect the yttria, the presence of which in the mineral cannot be indicated by mere blowpipe experiments alone. It has no crystalline structure, but appears in thin layers or seams, which sometimes amount to scarcely anything more than peach-blossom or purple stains, penetrating the seams of the limestone: precisely the character of this mineral in the specimens I have of it from Finbo in Sweden. With this it also agrees in hardness and colour. When heated in a glass tube, it slightly decrepitates, shows no phosphorescence, gives out moisture, and becomes milk-white; at the same time there is a perceptible burnt smell. When its powder, moistened with sulphuric acid, is placed in a platinum-crucible, hydrofluoric acid is given out by the application of heat, and the usual reaction on glass is produced. The pulverized mineral, heated with fused salt of phosphorus in an open glass tube, also shows the same reaction, the glass losing its polish where the moisture is deposited. In these experiments I was careful to separate the mineral entirely from the brucite; but I have not been able to obtain fragments sufficiently free from carbonate of lime, to enable me to give its blowpipe characters in detail, or subject it to any other trials. I hope to be able to obtain better specimens at an early day, and then to complete its examination. The mineral is very characteristic, and, in the hand specimen, cannot be distinguished from the Finbo variety.

#### *Ottrelite identical with Phyllite.*

The name of phyllite, from φύλλον, a leaf, was given by Dr. Thomson to a mineral which was discovered and sent to him for analysis by Prof. Nuttall. It comes from Sterling, Massachusetts, and is disseminated in small thin plates through what appears to be an argillo-micaceous slate. Some of these plates are angular and others rounded, not appearing to have any regular crystalline form; yet in a few instances they present the distinct form of rhomboidal tables. Colour brownish-black, or grayish-black: lustre, shining and semi-metallic; opaque; fracture uneven. The knife makes a faint

impression upon them. In strong transmitted light, the thinnest discs present a greenish colour. Before the blowpipe, on charcoal, it becomes magnetic, but does not fuse even on the edges; with double its bulk of borax, it slowly dissolves into a dark iron-green glass. Its composition, as stated by Dr. Thomson, is as follows:—

Silica . . . . .	38.40
Alumina . . . . .	23.68
Peroxide of iron . . . . .	17.52
Magnesia . . . . .	8.96
Potash . . . . .	6.80
Water . . . . .	4.80
	100.16

Ottrelite was discovered by M. Desclozeaux, and analysed by M. Damour in 1842. A full description of it is given in the *Annales des Mines* for that year, vol. ii. p. 357. It occurs in small discs or plates, of a grayish-black or greenish-black colour, with considerable metallic lustre, disseminated through a gangue which appears like a greenish argillaceous slate. These discs present no distinct form in the specimens I have examined, their edges being rounded, as in the case of the phyllite; but Desclozeaux has referred them to a hexagonal prism, or to an acute rhomboid deeply truncated by a plane perpendicular to the axis, or deeply compressed in that direction. He also obtained a cleavage parallel with that plane. Minute fragments are translucent, and show a greenish colour by transmitted light. Before the blowpipe, it fuses, alone, with difficulty, *on the edges*, into a black, magnetic globule. It dissolves slowly in borax, giving the reaction of iron, and with carbonate of soda, shows the presence of manganese.

Its constituents are as follows:—

	Oxygen.	Ratio.	Formulae.
Silica . . . . .	43.34 22.51	4	2Al Si + (Fe <sup>3</sup> , Mn <sup>3</sup> .) Si <sup>2</sup> + Aq.
Alumina . . . . .	24.63 11.50	2	
Protox. of iron . . . . .	16.72 3.80	} 5.63 1	
Protox. of manganese ..	8.18 1.83		2Al <sup>2</sup> O <sup>3</sup> SiO <sup>3</sup> + (Fe <sup>3</sup> MnO <sup>3</sup> .)
Water . . . . .	5.66 5.03	1	2SiO <sup>3</sup> + 3H O <sup>3</sup>
	98.53		

Dr. Thomson's analysis affords a different formula, and, according to his method of determining the atomic proportions, phyllite is a simple silicate (the atoms of silica and bases being equal), consisting of nine atoms silicate of alumina, three atoms silicate of peroxide of iron, three atoms silicate of manganese, and one atom silicate of potash\*. The occurrence of so large a proportion of potash in the mineral is not a little remarkable, and I would suggest whether it

\* Outlines of Mineralogy, &c., vol. i. p. 384. Dr. Thomson's atomic weights, founded upon the idea of Prout, that they are all multiples of the atomic weight of hydrogen vary somewhat from Berzelius's.

may not have been derived from the gangue of slate, from which it is difficult to obtain the mineral entirely free. Its infusibility before the blowpipe would seem to show this. It has been suggested, also, that a part of the iron may have been in the state of protoxide. It seems impossible, without some such supposition, that substances so closely resembling each other in all their physical characters, should differ so much in chemical composition. Now, if the potash be left out, and the peroxide of iron be changed into protoxide, the ratio between the atoms of acid and bases is nearly the same as in ottrelite, if we unite the atoms of magnesia and iron as isomorphous with each other. Ottrelite, also, is not easily separated from its matrix, but the larger size of its plates would seem to render it more easy to obtain pure specimens for analysis; and it is to be observed that Damour repeated his analysis, and obtained precisely the same result. It is remarkable that Rammelsberg has alphabetically inserted phyllite, but has given no formula for its constitution. It seems proper that the name of phyllite, on the ground of its priority, and because it expresses so well the ordinary appearance of the mineral, should stand, and that of ottrelite be abandoned\*.

*Dysluite identical with Automalite.*

I am satisfied, from recent observations, that these two minerals, as they occur in New Jersey, should form but one species. The difference in hardness, colour, specific gravity and pyrognostic characters, can be accounted for by the well-established fact of the isomorphous replacement among the constituents of certain minerals which do not differ in crystalline form. In dysluite we have but thirty per cent. of alumina, the acting acid principle in the mineral, while in automalite we have sixty per cent. But the peroxide of iron, which is isomorphous with the alumina, amounts to nearly forty-two per cent. Now, if we suppose about thirty per cent. of this peroxide of iron to have replaced the same number of atoms of alumina in automalite, and the eight per cent. of protoxide of manganese to have replaced so much of the oxide of zinc, we make up very nearly the essential constituents as shown in the analyses of automalite by Ekeberg and Abich. It is to be observed that the latter chemist puts down the iron as *protoxide* in the Franklin automalite. If it should prove that the iron exists in dysluite in both states of oxidation, the twelve per cent. remaining out of the forty-two may be protoxide, replacing so much oxide of zinc. So that in this view of the case, the 17 per cent. oxide of zinc + 11 per cent. protoxide of iron + 7 per cent. protoxide of manganese = 35 per cent. oxide of zinc, which is nearly the exact quantity found by Abich in the crystals from Franklin. We may then state the constituents as follow:—

\* Brooke has supposed phyllite to be identical with gigantolite. If we compare the analysis of gigantolite with Damour's analysis above, the evidence of their identity (supposing ottrelite to be a purer variety of phyllite) is much more marked, and the ratio between the atoms of acid and bases is nearly the same in each.

		Oxygen.	Ratio.
Alumina . . . . .	30·49	14·24	} 23·43    3
Peroxide of iron . . . . .	30·00	9·19	
Protoxide of iron . . . . .	11·93	2·72	} 7·76    1
Protoxide of manganese ..	7·60	1·70	
Oxide of zinc . . . . .	16·80	3·34	

Here it is evident that the atoms of acid and bases are to each other as three to one, which is the case also with automalite, taking Abich's analysis, and grouping the isomorphous bases, thus :

		Oxygen.	Ratio.
Alumina . . . . .	57·09	26·66	3
Oxide of zinc . . . .	34·80	6·92	} 8·72    1
Magnesia . . . . .	2·22	·76	
Protoxide of iron ..	4·55	1·04	

Dr. Thomson, the only chemist who has analysed dysluite, reckons all the iron as peroxide, and as the principal basic constituent of the mineral, which, in his view, consists of the aluminates of iron, zinc and manganese. Rammelsberg, in stating the analysis, has given both oxides, and the atoms of alumina and peroxide of iron, as put down by him, are 22·80, and those of the isomorphous bases—protoxide of iron, protoxide of manganese and oxide of zinc—are 7·83 (7·89 ?); thus giving the same ratio as that above stated.

But other reasons may be urged why dysluite should be regarded only as a variety of automalite. I have seen specimens on which there were crystals well claiming the name of dysluite, as well as others equally entitled to the name of automalite; while there were yet others evidently passing from one into the other,—the bright and perfect crystals of automalite gradually losing their lustre, becoming porous, comparatively brittle and soft. I think if these circumstances had been attended to in the early history of the mineral, the name dysluite would long since have departed from the catalogue of mineral species.

#### *Polyadelphite.*

As Dana, in the new edition of his Mineralogy, has very properly included this mineral under the species garnet, I merely refer to it, to give further evidence of the correctness of his opinion from circumstances connected with its occurrence at the locality. It is evidently a granular, imperfectly crystallized yellow garnet, and the specimen which I received ten years ago from Prof. Nuttall, contains mechanical mixtures which it would be impossible to separate from it, so as to give us entire confidence in its analysis. To these, I believe, we may attribute its departure in composition from the common brown or yellow garnet, though it does not differ much from the brown garnet of Franklin, analysed both by Dr. Thomson and Mr. Seybert.

#### *Beaumontite* of Levy, and *Lincolnite* of Hitchcock.

In a paper read before the Boston Society of Natural History, and since published in their Transactions, and in the American Journal



of Science (vol. xlv. p. 235), I gave my reasons for classing these two minerals with heulandite. That beaumontite is heulandite, I believe is no longer doubted in this country or Europe. An analysis of the mineral by M. Delesse, has appeared since the publication of my paper\*, and it agrees with all the other analyses of heulandite, excepting in the slight excess of silicic acid. In this respect it offers an example analogous to that of the variety of chabasite called acadolite, in which the silicic acid forms a larger atomic proportion of the mineral, without causing any appreciable variation in the angles of the crystals. As to lincolnite, I must think that the various papers that have been called forth in relation to it since my first communication appeared, have established its indisputable identity with heulandite.

Peculiarities in the modifying planes † have given rise to a secondary form, rarely observed in heulandite. These consist in the enlargement of the planes *f* (Phillips), or *ë* (Dana), so as nearly to obliterate the primary planes *M*; being, in fact, the reverse of what we usually observe in heulandite from other localities. In the measurements by Prof. Hitchcock and Prof. Shepard, the angle of *f* on *T* was mistaken for that of *M* on *T*, and in the figure given by Prof. Hitchcock, it is evident that the planes lettered *M* should be *f*. The true value of *f* on *T* is 115° 10' (Dana); Prof. Shepard's last measurements made it 116° 17'.

*Ledererite.*


I am compelled, at last, to declare my conviction that the specific nature of this mineral can no longer be maintained. Connell's analysis of an Irish gmelinite, which agrees with ledererite in all its physical and crystallographical characters, has shown also an identity in chemical composition. The phosphoric acid detected by Mr. Hayes must be viewed as an accidental constituent, varying probably in different crystals, or in some of them not existing at all. Some of the zeolites, in the Nova Scotia trap, have been found associated with small crystals of phosphate of lime, and it is not impossible that some of the minutest of these may have intercrystallized with the ledererite. We regret that we have not been able to obtain other specimens to enable Mr. Hayes to give it a re-examination. For comparison, I subjoin the analyses of ledererite and gmelinite.

	Ledererite.	Gmelinite.
Silica.....	49·47	48·56
Alumina .....	21·48	18·05
Lime.....	11·48	6·13
Soda.....	3·94	3·85
Phosphoric acid ..	3·48	Potash 0·39
Protoxide of iron ..	0·14	0·11
Water .....	8·58	21·66
	98·56, Hayes.	98·75, Connell.

\* *Ann. de Chim. et de Phys.* for 1843, t. ix. p. 395. Phillips's *Min.* p. 627.

† For the figures see *Amer. Journ. of Science*, vol. xlv. p. 234, and vol. xlvii. p. 416. Corroborative evidence of the correctness of my opinions

Now, if the phosphoric acid in ledererite is united with lime as an accidental mixture,  $2\frac{1}{2}$  per cent. of the lime should be taken from the 11.48 per cent. found in the mineral: this brings the proportion down nearly to that obtained by Connell. Mr. Hayes was not able to determine the weight of the water with accuracy, owing to the small quantity of the mineral operated upon. As the loss (1.44 per cent.) was mostly water, we may suppose, with Rammelsberg, that ledererite is gmelinite containing  $\frac{1}{3}$  ( $\frac{1}{2}$ ?) its quantity of water. The chemical formula for gmelinite and chabasite is thus:



Excepting the absence of striæ, and the shorter dimensions of the prismatic planes of its crystals, the Irish gmelinite precisely resembles ledererite; their hardness, lustre, colour and blowpipe characters are the same. The appearance of hexahedral cleavage, on which Dr. Jackson originally founded the chief claim of the latter to the character of a new species, was only imperfectly produced by heating the crystals, and not by ordinary mechanical cleavage. This could not be effected, the mineral breaking in all directions with a vitreous fracture. Dr. Jackson agrees with me that it can no longer be retained as a distinct species.

While preparing my edition of Phillips's Mineralogy, I requested Mr. Hayes and Dr. Jackson to make several analyses for me with particular reference to that work. As some of these have not appeared in any other form, I wish now to make a permanent record of them, in order that they may be seen where they might not otherwise reach. The first are of the Nova Scotia chabasite (acadiolite), which Hoffmann has distinguished from common chabasite, by its containing three per cent. more silica, and for which Rammelsberg has given a formula differing somewhat from that of chabasite. (See first part of this article.)

Silica . . . . .	52.02	52.20
Alumina ..	17.88	18.27
Lime . . . . .	4.24	6.58
Potash . . . . .	3.03	2.12
Soda . . . . .	4.07	
Water . . . . .	18.30	20.52
	99.60, Hayes.	99.69, Hayes.

These results agree with those obtained by Hoffmann† in his analysis of the same mineral, the specimens of which were presented to him by Charles Cramer, Esq. of St. Petersburg.

by the editors of the Amer. Journ. of Science, may be seen at the pages here referred to.

\* *Handwörterbuch*, i. 150. Rammelsberg unites chabasite and gmelinite, the first as soda chabasite, the last as lime chabasite. This is in accordance with Tamnau, who has established their identity on crystallographical grounds. The close relation of the two minerals was, however, first shown by Prof. Mohs. See his Mineralogy, vol. ii. p. 105.

† Amer. Journ. of Science, vol. xxx. p. 366.

Washingtonite of Shepard, analysed by Mr. J. S. Kendall under the direction of Dr. Jackson, gave these results:—

		Oxygen.	Ratio.
Titanic acid. . . . .	25·28	4·82	1
Peroxide of iron . . . .	51·84	10·36	2
Protoxide of iron ..	22·86	5·08	1
	<u>99·98</u>		

The atomic proportions are thus nearly one atom titanic acid, two atoms peroxide of iron, one atom protoxide of iron; or, a trititanate of iron, consisting of two atoms trititanated peroxide and one atom trititanated protoxide. If we unite the magnesia and lime with protoxide of iron in the following analysis of an ilmenite from Arendal\*, by Mosander, we obtain precisely the same result. The crystalline form of the two varieties is also the same, and there can be no doubt of their identity as one species †.

Titanic acid. . . . .	24·19	
Peroxide of iron . . . . .	53·01	
Protoxide of iron . . . .	19·91	} 20·92
Magnesia and lime ..	1·01	

By referring to the analyses of ilmenite from other localities, it will be seen that the essential constituents, titanic acid and the two oxides of iron, so interchange with each as to produce different varieties, but all having the same crystalline form.

**NOTICE ON CERTAIN IMPURITIES IN COMMERCIAL SULPHATE OF COPPER. BY MR. S. PIESSE.**

One source of the sulphate of copper of commerce is the treatment of brass and German silver articles technically called dipping, which consists in plunging them for a short time into a mixture of nitric and sulphuric acids, an operation which removes the coat of oxide from the surface of the metal, and leaves the latter in a clean state proper for the reception of varnish or other finishing. In time this dipping liquid becomes in great measure saturated, and after neutralization with old copper yields on evaporation in leaden pans a large quantity of sulphate of copper in crystals. According to the author not less than 100 tons of dipping liquid are thus disposed of annually at Birmingham by the makers of buttons and other arti-

\* The hystatite of Breithaupt.

† An acute rhombohedron, P on P 86° 10', for the ilmenite. Shepard, employing varnished planes of the washingtonite, makes P on P 86°. Prof. Shepard finds the distinction on other than crystallographical characters; for, he says, it is not thus "shown to be distinct, in any essential manner, from the axotomous iron ore of Mohs, or from crichtonite (including ilmenite): indeed, it appears most probable that all these minerals are not only identical in their angles, but are isomorphous with specular iron."—Amer. Journ. vol. xliii. p. 365. The analysis, now, would seem to destroy the groundwork for any distinction.

cles. The crystallized sulphate of copper so obtained is often largely contaminated with sulphate of zinc, which may sometimes be seen in the form of slender white needles on the surface of the dark blue crystals, and in some of the applications of this salt may prove injurious. Sulphate of nickel, sulphate of lead, arsenic, and chlorides are also sometimes present.—*From the Proceedings of the Chemical Society.*

---

ON A NEW EUDIOMETRIC PROCESS. BY PROF. GRAHAM.

Professor Graham described a new eudiometric process for the rapid absorption of oxygen gas from atmospheric air and other gaseous mixtures containing oxygen. It consists in the employment of a solution in ammonia of a sulphite of the suboxide of copper and ammonia. This salt falls as a granular powder, when a stream of sulphurous acid gas is conveyed into a cold solution of the ammoniacal sulphate of copper. When dissolved in ammonia it absorbs oxygen with singular avidity, and when employed in this form in eudiometry gives results of considerable uniformity.—*From the Proceedings of the Chemical Society.*

---

EQUIVALENT OF CHLORINE.

M. Gerhardt observes, that M. Marignac has made some observations and experiments tending to show that the atomic weight of chlorine is not thirty-six times that of hydrogen, as he (M. Gerhardt) had concluded, but somewhat less. M. Marignac's conclusions are derived from the weight of chloride of potassium yielded by the calcination of chlorate of potash; to these results M. Gerhardt makes the following objections:—

It is the residue of chloride of potassium obtained, and not that of the oxygen gas, which is weighed; and from the following causes the quantity of chloride might be too small, and would diminish the atomic weight of chlorine:—a trace of moisture on the salt; a portion of chlorate or chloride carried off by the current of oxygen gas; if the oxygen were impure, and contained, as M. Marignac has stated, a trace of chlorine.

Thus, observes M. Gerhardt, all the errors which can be committed in these determinations are referrible to the chlorine, and give it in too small quantity.

From his experiments M. Gerhardt concludes, in opposition to those of M. Marignac, that the equivalents of chlorine, silver and potassium, are exact multiples of the equivalent of hydrogen, that is to say,—

Chlorine . . . . .	36
Silver . . . . .	108
Potassium . . . . .	39

ON HIPPURIC ACID, BENZOIC ACID, AND THE SUGAR OF GELATINE.

M. Dessaignes remarks, that hippuric acid has already been the subject of numerous researches; its metamorphoses are nevertheless so interesting as to leave something for those to glean who will study it. M. Liebig has shown, that when it is dissolved in boiling hydrochloric acid, it crystallizes on cooling without having been altered; but if the ebullition be prolonged for about half an hour, it is decomposed, and yields, according to M. Dessaignes, benzoic acid equal nearly in quantity to that indicated by theory. The benzoic acid was separated by the filter, and the filtered liquor gave by evaporation long, acid, nitrogenous prismatic crystals, into the composition of which hydrochloric acid entered as a constituent part. These crystals were neutralized by carbonate of soda or carbonate of lead; and after getting rid of the solution of chloride of sodium or chloride of lead, fresh crystals of a very saccharine and azotized matter were obtained; these were neutral to reagents, and formed crystalline compounds with oxide of silver, and with nitric, sulphuric, and oxalic acids.

M. Dessaignes soon found out that he had thus produced, by a metamorphosis which might have been foreseen, the sugar of gelatine discovered by M. Braconnot.

Reckoning C=150, H=6.25, and N=17.5, if from



to which it is sufficient to add 1½ equivalent of water to obtain sugar of gelatine. M. Dessaignes is inclined to the opinion, that 2 equivalents of water should be added, and that the true equivalent of sugar of gelatine is C<sup>4</sup> H<sup>10</sup> N<sup>2</sup> O<sup>4</sup>, as already indicated by M. Gerhardt.

All the reactions and beautiful crystallizations which M. Dessaignes obtained with the saccharine azotized matter from hippuric acid, convinced him of the identity of this substance with the sugar of gelatine obtained from isinglass; but in order to convince chemists of this fact, he thinks it requisite to analyse the sugar of hippuric acid. The metamorphosis which gives rise to this body is very distinct; no gas is evolved during the reaction; the only two products are benzoic acid and hydrochlorate of sugar. From 100 of dry hippuric acid M. Dessaignes obtained,—

Benzoic acid (dry) .....	67.49
Hydrochlorate of sugar (dried over sulphuric acid)	59.08
	126.57

Nitric acid boiled for twenty minutes with hippuric acid, converts it into benzoic acid, and nitro-saccharic acid, which crystallizes in magnificent truncated tables. Nitro-saccharic acid, prepared with the sugar of isinglass, yielded precisely similar crystals.

Sulphuric acid diluted with twice its volume of water also effects the metamorphosis of hippuric acid without the disengagement of gas, and without colouring the solution. The benzoic acid obtained is very easily purified, and also a compound, from which, by means of chalk or carbonate of lead, sugar of gelatine may be procured.

M. Dessaignes combined, equivalent to equivalent, sulphuric acid  $\text{SO}^3 \text{H}^2 \text{O}$ , and the sugar obtained from hippuric acid, giving as the formula of the latter  $\text{C}^4 \text{H}^{10} \text{N}^2 \text{O}^4$ ; and he obtained a solution which crystallized in large prisms of great splendour, to the last drop.

A very concentrated solution of oxalic acid boiled for two hours with hippuric acid, converts it into benzoic acid and oxalate of sugar, which crystallizes in fine prisms. Lastly, an excess of potash or soda, boiled for half an hour with hippuric acid, converts it into alkaline benzoate and sugar, which was obtained in the form of hydrochlorate, after having treated the mixture of benzoate and sugar with hydrochloric acid.—*Ann. de Ch. et de Phys.*, Mai 1846.

---

COMPARATIVE ANALYSES OF ORIENTAL JADE AND TREMOLITE.  
BY M. DAMOUR.

The jade selected for analysis had been worked in India; it was of a milk-white colour and semi-transparent, and had the appearance of white wax, or perhaps rather of spermaceti. Its fracture is splintery; it scratches glass, but feebly. Its specific gravity was found to be 2.970. Its tenacity is very great; when reduced to powder and heated in a glass tube, its appearance was not altered, and it yielded no water. In the flame of the blowpipe it swells up, and fuses slowly into a milk-white enamel. Borax dissolves it without colour; the salt of phosphorus dissolves it, leaving a skeleton of silica. It is not sensibly acted upon by hydrochloric acid.

Two analyses gave the following results:—

Silica .....	58.46	58.02
Lime.....	12.06	11.82
Magnesia.....	27.09	27.19
Protoxide of iron ..	1.15	1.12
	<u>98.76</u>	<u>98.15</u>

M. Damour having observed that this is precisely the composition of tremolite (white amphibole), submitted this substance to the same process of analysis as that adopted with the jade. The specimen which he selected was from St. Gothard, and in colourless crystals, very perfect and associated with granular dolomite, which was separated by hydrochloric acid previously to analysis.

It yielded,—

Silica.....	58.07
Lime.....	12.99
Magnesia.....	24.46
Protoxide of iron ..	1.82
	<u>97.34</u>

From the similarity of these results, M. Damour is of opinion that this jade may be ranked with tremolite; and if this opinion should be adopted, he observes, that in collections oriental jade will hereafter be classed as *compact tremolite*.—*Ann. de Ch. et de Phys.*, Avril 1846.

METEOROLOGICAL OBSERVATIONS FOR APRIL 1846.

*Chiswick*.—April 1. Fine. 2. Cloudy: showery. 3. Clear and windy: cloudy and fine. 4. Hazy: heavy rain. 5. Heavy rain: clear. 6. Heavy rain: cloudy. 7. Slight rain: densely overcast. 8—10. Fine. 11. Dry haze. 12, 13. Cloudy and fine. 14. Clear: dry haze: overcast. 15. Densely clouded: dry haze: densely overcast. 16. Slight dry haze. 17. Foggy. 18. Rain. 19. Cloudy and cold: clear. 20. Showery: frosty at night. 21. Foggy: cloudy and fine. 22. Foggy. 23. Heavy clouds: rain. 24. Rain: dark haze: cloudy. 25. Hazy and damp: showery: hazy: foggy. 26. Extraordinary fall of rain early A.M.: dense clouds: overcast at night. 27. Clear and fine. 28. Very fine. 29, 30. Clear: very fine: overcast.

Mean temperature of the month .....	47°·36
Mean temperature of April 1845 .....	48 ·41
Average mean temperature of April for the last twenty years	47 ·19
Average amount of rain in April .....	1 ·47 inch.

*Boston*.—April 1. Fine. 2. Rain. 3. Windy: rain P.M. 4. Cloudy: rain P.M. 5. Cloudy: rain A.M. and P.M. 6. Cloudy. 7. Rain. 8. Cloudy: rain early A.M. 9, 10. Fine. 11. Cloudy: rain P.M. 12. Fine. 13. Fine: rain early A.M. 14. Fine: rain P.M. 15. Cloudy: rain early A.M. 16—18. Cloudy. 19. Cloudy: rain A.M. 20—22. Fine. 23. Fine: rain A.M. 24, 25. Fine. 26. Cloudy: rain A.M. 27. Fine: rain A.M. 28. Cloudy. 29. Fine: ice this morning. 30. Cloudy.

*Sandwich Manse, Orkney*.—April 1. Snow: clear. 2. Showers: clear. 3. Snow-showers. 4. Snow-showers: frost: snow-showers. 5. Showers. 6. Showers: clear: aurora. 7. Clear: drops. 8. Cloudy: clear. 9. Bright: cloudy. 10. Bright: showers. 11. Bright: rain. 12. Fog: damp. 13. Damp: drizzle. 14. Clear. 15. Fog: cloudy. 16. Cloudy. 17. Cloudy: damp: fog. 18. Rain: clear. 19—21. Fine: clear. 22. Clear: cloudy. 23, 24. Clear. 25. Cloudy. 26. Sleet-showers: hail-showers. 27. Bright: cloudy. 28. Hail-showers: cloudy. 29. Snow-showers: clear. 30. Cloudy: clear.

*Applegarth Manse, Dumfries-shire*.—April 1. Wet. 2. Wet A.M.: cleared and fine. 3. Wet A.M.: cleared. 4. Slight showers: frost A.M. 5. Fair, but chilly. 6. Fair, but very bleak. 7. Fair. 8. Fair: frost A.M. 9. Fair: frost: fine. 10. Fine. 11. Rain all day. 12. Rain P.M.: thunder. 13. Frequent heavy showers. 14. Frequent heavy showers: hail: fine P.M. 15. Frequent heavy showers: rain all day. 16. Very fine spring day. 17, 18. Dropping day. 19. Fair, though chilly. 20. Frost, slight: fine. 21. Hoar-frost: rain P.M. 22. Slight showers. 23, 24. One slight shower. 25. Heavy shower: fair P.M. 26. Slight shower: fine. 27. Slight shower: frost A.M. 28. Frost A.M.: fine. 29. Frost A.M.: a slight shower. 30. A dropping day.

Mean temperature of the month .....	45°·6
Mean temperature of April 1845 .....	48 ·2
Mean temperature of April for twenty-three years	44 ·2
Mean rain in April for eighteen years .....	1 ·69 inch.





## INDEX TO VOL. XXVIII.

- ACIDS**:—ellagic, 41; margaric, 68; chloro-acetic, 154; phosphoglyceric, 158; resino-bezoardic, 192; valerianic, 234; hyponitric, 432; cinnamic, 442; carbazotic, 443; chloroazotic, 555; hippuric, 567.
- Airy (G. B.) on the equations applying to light under the influence of magnetism, 469; on Dr. Faraday's paper on ray-vibrations, 532.
- Albumen, observations on, 369.
- Alexander (Prof.) on the spots of the sun, 230.
- Algebraic equations of the fifth degree, on the resolution of, 63.
- Alger (F.) on new localities of rare minerals, and reasons for uniting several supposed distinct species, 557.
- Alloxan, on the preparation of, 550.
- Alumina, analysis of phosphate of, 68.
- Ammonia, on the decomposition and analysis of the compounds of, 222.
- Animal concretions, new species of, 36, 192.
- Atomic volume and specific gravity, observations on, 527.
- Aurora borealis, account of an, 70.
- Automalite, on the identity of, with dysluite, 561.
- Barometer, on the causes of the semi-diurnal fluctuations of the, 166, 416; on the oscillations of the, 241.
- Baudrimont (M.) on the preparation and composition of chloroazotic acid, 555.
- Beaumontite, observations on, 562.
- Beck (Mr.) on the nerves of the uterus, 408.
- Beluga stones, examination of the, 38.
- Bessel (Prof.), notice of the late, 343.
- Bezoar, oriental, examination of the, 41.
- Biela's comet, notice of, 238.
- Birt (W. R.) on the storm-paths of the eastern portion of the North American continent, 379.
- Black, remarks on certain statements in Lord Brougham's life of, 106, 478.
- Blood, on the cause of the circulation of the, 178.
- Bodies, solid, on the temperature and conducting power of, 161.
- Boracic aether, on the preparation and properties of, 337.
- Bronwin (Rev. B.) on the determination of the motion of a disturbed planet, 20; on certain definite multiple integrals, 373.
- Brougham's (Lord) Lives of Black, Watt and Cavendish, remarks on certain statements in, 106, 478.
- Brown (W.) on the oscillations of the barometer, 241.
- Calculi, chemical examination of some new species of, 36, 192.
- Cane-sugar, conversion of, into a substance isomeric with cellulose and inuline, 12.
- Capillarity, observations on, 341.
- Cassini (J. D.), notice of the late, 412.
- Cavendish, remarks on certain statements in Lord Brougham's life of, 106, 478.
- Cayley's (C. B.) inquiries in the elements of phonetics, 47.
- Ceradia furcata* resin, analysis of, 422.
- Chabasite, observations on, 557.
- Challis (Rev. J.) on the aberration of light, 90, 176, 393; on Biela's comet, 238.
- Chamærops*, on the wax of the, 350.
- Cheese, on the volatile acids of, 234.
- Chemistry:—influence of magnetism on crystallization, 1, 94; conversion of cane-sugar into a substance isomorphous with cellulose, 12; solubility of oxide of lead in pure water, 17; new species of animal concretions, 36, 192; action of nitric acid on wax, 66; dry distillation of wax, 67; phosphate of alumina, 68; preparation of chloro-acetic acid, 154; composition of phosphate of ammonia and magnesia and of the phosphate of soda, 155; on some double oxalates, 156; test for sulphurous acid, 157; analysis of the yolk of eggs, 158; on the ferrocyanide of potassium, 211; on the decomposition and analysis of compounds of ammonia and cyanogen, 222; method of obtaining pure oxide of uranium, 232; new double haloid salts, *ib.*; volatile acids of cheese, 234; amount of water in the

- double salts of the magnesian group, 235, 289; preparation of hypophosphites, 236; boracic aether, 337; action of boracic acid on pyroxylic spirit, 339; on the wax of *Chamærops*, 350; on pegmine and pyropine, 368; description of a new mercurial trough, 406; new compounds of perchloride of tin, 416; analysis of two species of epiphytes, 420; resin of *Ceradia furcata*, 422; on the constitution of the internal gas of plants, 426; relation of ozone to hyponitric acid, 432; composition of the fire-damp of the Newcastle coal-mines, 437; resin of *Xanthoræa hastilis*, 440; on atomic volume and specific gravity, 527; preparation of alloxan, 550; preparation of chloroazotic acid, 555; on certain impurities in commercial sulphate of copper, 565; on a new endiometric process, 566; on the equivalent of chlorine, *ib.*; decomposition of hippuric acid into benzoic acid and sugar of gelatine, 567.
- China, on the anthracite and bituminous coal-fields of, 204.
- Chlorine, on the equivalent of, 566.
- Chloro-acetic acid, preparation of, 154.
- Chloroazotic acid, on the preparation of, 555.
- Christie (J. R.) on the use of the barometric thermometer for the determination of heights, 220.
- Circulation of the blood, on the cause of the, 190.
- Coal-fields of China, observations on the, 204.
- Cobalt ore, analysis of a, from Western India, 352.
- Cockle (J.) on a proposition relating to the theory of equations, 132; on the finite solution of equations of the fifth, sixth and higher degrees, 191, 395.
- Cohesion of liquids, observations on the, 293.
- Collen (H.) on the application of the photographic camera to meteorological registration, 73.
- Comet, on a direct method of determining the distance of a, 226; Biela's comet, 238.
- Connell (Prof.) on the composition of the Elie pyrope or garnet, 152.
- Crystalline particles, on certain molecular actions of, 1, 94.
- Cyanogen, on the decomposition and analysis of the compounds of, 222.
- Daguerreotype process, on the cause of the fixation of mercurial vapours in the, 94.
- Damour (M. A.) on the composition of diaspore from Siberia, 336; on the composition of oriental jade and tremolite, 568.
- Dana (J. D.) on the origin of the constituent and adventitious minerals of trap and the allied rocks, 49.
- Daniell (J. F.), notice of the late, 409.
- Delesse (M. A.) on the composition of native phosphate of alumina, 68; analysis of a substance occurring with disthene, 150; on a double hydrated silicate of magnesia, 152.
- Dell (T.) on the transit of Mercury, 224.
- De Morgan (A.) on the invention of fluxions, 222; on the derivation of the word theodolite, 287; on the first introduction of the words tangent and secant, 382.
- Dessaigues (M.) on the decomposition of hippuric acid into benzoic acid and sugar of gelatine, 567.
- Diamagnetic bodies, action of magnets on, 403.
- Diaspore, analysis of, 336.
- Differentiation as applied to periodic series, observations on, 213.
- Donny (M. F.) on the cohesion of liquids, 293.
- Draper (Dr. J. W.) on the circulation of the blood, 178.
- Dysluite, on the identity of, with automalite, 561.
- Earth, on the connexion between the rotation of the, and the geological changes of its surface, 106.
- Earthquakes, on the vorticose movement, assumed to accompany, 537.
- Ebelmen (M.) on boracic aether, 337; on the action of boracic acid on pyroxylic spirit, 339.
- Eggs, on the composition of the yolk of, 158.
- Electric currents, on the production of musical sounds in metals by, 544.
- Electricity, experimental researches in, 64, 147, 294, 324, 455.
- Electro-magnetism, experiments on the mechanical powers of, 448.
- England, on the cause of remarkably mild winters which occasionally occur in, 317.
- Epidermis, on the development and growth of the, 82.
- Epiphytes, analysis of two species of, 420.
- Equations, on a proposition relating to the theory of, 132; of the fifth, sixth and higher degrees, on the existence of finite algebraic solutions of, 190, 395.
- Endiometric process, description of a new, 566.

- Faraday's (Prof.) experimental researches in electricity, 64, 147, 294, 324, 396, 455; on ray-vibrations, 345, 532.
- Ferrocyanide of potassium, on the conversion of, into the sesqui-ferrocyanide, 211; on the decomposition of, by solar light, *ib.*
- Fire-damp of the Newcastle coal-mines, on the composition of the, 437.
- Fluxions, on the invention of, 222.
- Forbes (J. D.) on the viscous theory of glacial motion, 219.
- Fox (R. W.) on certain pseudomorphous crystals of quartz, 5.
- Fresenius (M.) on the composition of the phosphate of ammonia and magnesia, 155.
- Gardner (Dr. D. P.) on the function of plants, 425.
- Garnet, analysis of the, 152.
- Gas lighting in China, notice respecting the antiquity of, 209.
- Gerhardt (M.) on the action of nitric acid on wax, and on the dry distillation of wax, 66; on the equivalent of chlorine, 566.
- Glacier motion, on the viscous theory of, 219.
- Glass, action of magnets on, 399.
- Gobley (M.) on the composition of the yolk of eggs, 158.
- Goodsir (J.) on the supra-renal, thymus and thyroid bodies, 220.
- Graham (Prof. T.) on the proportion of water in the magnesian sulphates and double sulphates, 289; on the composition of the fire-damp of the Newcastle coal-mines, 437; on a new eudiometric process, 566.
- Gregory (Prof. W.) on the preparation of alloxan, 550.
- Guanite, description and analysis of, 548.
- Guano deposits, account of various substances found in, 546.
- Gulf-stream, on the influence of the, in reference to the mild winters which occasionally occur in England, 317.
- Hail, on the formation of, 104.
- Harcourt (Rev. W. V.) on certain statements in Lord Brougham's Lives of Black, Watt and Cavendish, 106, 478.
- Heberden (Dr. W.), notice of the late, 408.
- Heights, on the use of the barometric thermometer for the determination of, 220.
- Heintz (M.) on a reaction for the discovery of sulphurous acid, 157.
- Hennessy (H.) on the connexion between the rotation of the earth and the geological changes of its surface, 106.
- Henry (Prof.) on the spots of the sun, 230; on a simple method of protecting from lightning buildings with metallic roofs, 340; on capillarity, 341.
- Henwood (W. J.), abstract of meteorological observations made in the interior of Brazil, 364.
- Hippuric acid, on the decomposition of, into benzoic acid and sugar of gelatine, 567.
- Hopkins (T.) on the causes of the semi-diurnal fluctuations of the barometer, 166, 416.
- Horses, experiments on the mechanical power of, 448.
- Hunt (R.) on the influence of magnetism on molecular arrangement, 1.
- Hypophosphites, preparation of, 237.
- Iguana, examination of a urinary calculus from the, 36.
- Iljenko (M.) on the volatile acids of cheese, 234.
- Infinite geometrical series, general expression for the sum of an, 10.
- Integrals, on certain definite multiple, 373.
- Jerrard (G. B.) on the resolution of algebraic equations of the fifth degree, 63.
- Jesuiticus on Fresnel's theory of double refraction, 144, 215.
- Jones (Dr. C. H.) on the secretory apparatus and function of the liver, 223.
- Joule (J. P.) on the mechanical powers of electro-magnetism, steam and horses, 448; on atomic volume and specific gravity, 527.
- Kepler's works, observations on a collection of, 387.
- Langberg (Chr.) on the determination of the temperature and conducting power of solid bodies, 161.
- Laskowski (M.) on the volatile acids of cheese, 234.
- Lassell (W.) on the solar eclipse of 1845, and on the transit of Mercury, May 8th, 1845, 223.
- Lead, on the solubility of the oxide of, in pure water, 17.
- Ledererite, analysis of, 563.
- Lewy (M.) on some new compounds of perchloride of tin, 416.
- Lhotsky (Dr. J.) on Kepler's works, 387.
- Light, on the aberration of, 15, 76, 90, 176, 335, 393; on the magnetization of, 64, 294, 324; action of electric currents on, 303; on the equations applying to, under the influence of magnetism, 469.
- Lightning, on a simple method of protecting buildings from, 340.
- Lincolnite, observations on, 562.


- Liquids, on the cohesion of, 293.
- Lithofellinic acid calculi, examination of, 192.
- Liver, on the secretory apparatus and function of the, 223.
- Loomis (Prof.) on the winter storms of the United States, 200.
- Louyet (Prof.) on a new mercurial trough, 406.
- Maclagan (Dr.) on the conversion of sugar into a substance isomeric with cellulose and inuline, 12.
- Magnesia, analysis of the hydrated silicate of, 152.
- Magnesian sulphates, on the proportion of water in the, 289.
- Magnetic lines of force, on the illumination of, 64, 294.
- Magnetism, influence of, on molecular arrangement, 1; on the equations applying to light under the influence of, 469.
- Magnets, action of, on metals, 455.
- Malaguti (M.) on chloro-acetic acid, 154.
- Mallet (R.) on the vorticose movement, assumed to accompany earthquakes, 537.
- Marignac's (Prof.) observations on Messrs. Playfair and Joule's memoir on atomic volume and specific gravity, 527.
- Mathematical Society, notice respecting the late, 225.
- Matter, on the magnetic condition of, 147, 396; on the constitution of, 443.
- Mercurial trough, description of a new, 406.
- Mercury, observations on the transit of, 224.
- Metals, on the action of magnets on, 455.
- Meteoritic iron, analysis of, 154.
- Meteorological observations, 71, 159, 239, 343, 423, 569.
- made at Gongo Soco in the interior of Brazil, abstract of, 364.
- phenomena of 1842, observations on the, 241.
- registration, on the application of the photographic camera to, 73.
- Meteorology of Bombay, on some points in the, 24.
- Meyer (E. I.), analysis of the molares of a fossil rhinoceros, 158.
- Micrometer, description of a new, 229.
- Middleton (J.) on a cobalt ore found in Western India, 352.
- Mineralogy:—influence of magnetism on crystallization, 1, 94; pseudomorphous crystals of quartz, 5; origin of the constituent and adventitious minerals of trap and allied rocks, 49; phosphate of alumina, 68; disthene, 150; kerolite, 152; Elie pyrope or garnet, *ib.*; diaspore, 336; analysis of a cobalt ore from Western India, 352; guanite, 548; phacolite, 557; ytthro-cerite, 559; ottrelite, *ib.*; dysluite, 561; polyadelphite, 562; beaumontite, *ib.*; ledere-rite, 563; oriental jade and tremolite, 568.
- Minerals of trap and the allied rocks, on the origin of the, 49.
- Molecular arrangement, influence of magnetism on, 1, 94.
- Moon (R.) on Fresnel's theory of double refraction, 134; on the evaluation of the sums of neutral series, 136; reply to Jesuiticus, 215.
- Musical sounds produced in metals by discontinuous electric currents, on the, 544.
- Optics, physical, contributions to, 212.
- Ottrelite, on the identity of, with phyllite, 559.
- Owen (Prof.) on the structural relations of organized beings, 525.
- Oxalates, on several new series of double, 156.
- Ozone, relation of, to hyponitric acid, 432.
- Pegmine, observations on, 368.
- Phacolite, observations on, 557.
- Phonetics, inquiries in the elements of, 47.
- Phosphate of ammonia and magnesia, on the composition of, 155.
- Phosphate of soda, composition of, 155.
- Photographic camera, on the application of the, to meteorological registration, 73.
- Phyllite, on the identity of, with ottrelite, 559.
- Pierre (M. J. J.) on the double salts of the magnesian group, 235.
- Piesse (S.) on certain impurities in commercial sulphate of copper, 565.
- Planet Astræa, notice respecting the, 69.
- Planet, equations for the determination of the motion of a disturbed, 20.
- Plants, researches on the functions of, 425; constitution of the internal gas of, 426; on the absorption of gases by, 429.
- Playfair (Dr. L.) on atomic volume and specific gravity, 527.
- Poggiale (M.) on some new double haloid salts, 232.
- Polyadelphite, notice respecting, 562.
- Potter (Prof.) on physical optics, 212.
- Pouillet (M.) on the recent researches of Prof. Faraday, 324.
- Powell (Prof.) on a new double-image micrometer, 229.
- Pyrope, observations on, 368.

- Pyroxylic spirit, action of boræic acid on, 339.
- Quartz, on certain pseudomorphous crystals of, 5.
- Ray-vibrations, thoughts on, 345, 532.
- Redfield (Mr.) on the storm-paths of the North American continent, 379.
- Reece (M. R.) on several new series of double oxalates, 156.
- Refraction, on Fresnel's theory of double, 48, 134, 144, 215.
- Resin of *Xanthoræa hastilis*, examination of, 440.
- Resino-bezoardic acid calculi, examination of, 192.
- Rhinoceros, fossil, analysis of the molares of, 158.
- Rockwell (C. H.) on meteoric iron from Burlington, 154.
- Ronalds (Mr.) on the application of the photographic camera to meteorological registration, 73.
- Royal Astronomical Society, proceedings of the, 223.
- Royal Society, proceedings of the, 64, 147, 219, 408.
- Sabine (Lieut.-Col.) on some points in the meteorology of Bombay, 24; on the winter storms of the United States, 200; on the cause of mild winters, 317.
- Salts, on some new double haloid, 232.
- Saussure (Th. de), notice of the late, 413.
- Schœnbein (Dr. C. F.) on the conversion of the solid ferrocyanide of potassium into the sesqui-ferrocyanide, 211; on the decomposition of the yellow and red ferrocyanides of potassium by solar light, *ib.*; on the relation of ozone to hyponitric acid, 432.
- Schumacher (H. C.) on a new planet, 69.
- Scoresby (Rev. W.) on the mechanical powers of electro-magnetism, steam and horses, 448.
- Secant, on the first introduction of the word, 382.
- Sloggett (H.) on the constitution of matter, 443.
- Smith (A.) on Fresnel's theory of double refraction, 48.
- Smyth (Dr. R.) on the decomposition and analysis of the compounds of ammonia and cyanogen, 222.
- Solar light, on the decomposition of the yellow and red ferrocyanides of potassium by, 211.
- Somerville (Mrs.) on the action of the rays of the spectrum on vegetable juices, 66.
- Spectrum, action of the rays of the, on vegetable juices, 66.
- Steam, experiments on the mechanical power of, 448.
- Stenhouse (Dr.) on the yellow gum-resin of New Holland, 440.
- Stokes (G. G.) on the aberration of light, 15, 76, 335.
- Storms of the United States, on the, 200.
- Storm-paths of the North American continent, observations on the, 379.
- Strickland (H. E.) on the structural relations of organized beings, 354, 525.
- Structural relations of organized beings, on the, 354, 525.
- Sturgeon (W.) on an aurora borealis seen at Manchester, 70.
- Sulphurous acid, test for the discovery of, 157.
- Sun, experiments on the spots on the, 230.
- Tangent, on the first introduction of the word, 382.
- Taylor (R. C.) on the anthracite and bituminous coal-fields in China, 204.
- (T.) on some new species of animal concretions, 36, 192.
- Teschemacher (E. F.) on various substances found in the guano deposits, 546.
- (J. E.) on the wax of the *Chamærops*, 350.
- Theodolite, on the derivation of the word, 287.
- Thermometer, barometric, on the use of the, for the determination of heights, 220.
- Thomson (J.), analysis of two species of epiphytes, or air-plants, 420.
- (Dr. R. D.) on pegmine and pyropine, 369; analysis of *Ceradia furcata* resin, 422.
- Tilley (Dr.) on the conversion of cane-sugar into a substance isomeric with cellulose and inuline, 12.
- Tin, on some new compounds of the perchloride of, 416.
- Tremolite, analysis of, 568.
- Uranium, mode of purifying the oxide of, 232.
- Vegetable juices, action of the rays of the spectrum on, 66.
- Waller (Dr. A.) on certain molecular actions of crystalline particles, &c., 94.
- Wartmann (Prof. E.) on the causes to which musical sounds produced in metals by electric currents are attributable, 544.
- Waterston (J. J.) on a direct method of determining the distance of a comet, 226.

- Watt, remarks on certain statements in Lord Brougham's life of, 106, 478.
- Wax, on the action of nitric acid on, 66 ; dry distillation of, 67.
- Wilson (E.) on the development and growth of the epidermis, 82.
- Wöhler (Prof.), method of purifying oxide of uranium from nickel, cobalt and zinc, 232.
- Wurtz (A.) on the preparation of the hypophosphites, 236.
- Yorke (Lieut.-Col. P.) on the solubility of the oxide of lead in pure water, 17.
- Young (Prof. J. R.) on the general expression for the sum of an infinite geometrical series, 10 ; on differentiation as applied to periodic series, 213.
- Yttrite, on a new locality for, 559.

END OF THE TWENTY-EIGHTH VOLUME.

PRINTED BY RICHARD AND JOHN E. TAYLOR,  
RED LION COURT, FLEET STREET.




BOMBAY 1843.

*Diurnal Variations.*

*Gaseous Pressure* —————

*Tension of Vapour* - - - - -




*Annual Variations*

*Gaseous Pressure* —————

*Barom<sup>c</sup> Pressure* - - - - -

*Tension of Vapour* - - - - -



LONDON  
Richards  
PRINTED & SOLD





*Handwritten: Magnet-Crystallization*

Fig. 1.




Fig. 3.




Fig. 2.




Fig. 4.




Fig. 5.




Fig. 8.




Fig. 7.




Fig. 6.




Fig. 10.




Fig. 9.

